arXiv:1208.2726v1 [math.AP] 13 Aug 2012

WELL-POSEDNESS OF THE FREE-BOUNDARY COMPRESSIBLE 3-D EULER EQUATIONS WITH SURFACE TENSION AND THE ZERO SURFACE TENSION LIMIT

DANIEL COUTAND, JASON HOLE, AND STEVE SHKOLLER

Abstract. We prove that the 3-D compressible Euler equations with surface tension along the moving free-boundary are well-posed. Specifically, we consider isentropic dynamics and consider an equation of state, modeling a liquid, given by Courant and Friedrichs [8] as $p(\rho) = \alpha\rho^\gamma - \beta$ for consants $\gamma > 1$ and $\alpha, \beta > 0$. The analysis is made difficult by two competing nonlinearities associated with the potential energy: *compression* in the bulk, and *surface area dynamics* on the free-boundary. Unlike the analysis of the incompressible Euler equations, wherein boundary regularity controls regularity in the interior, the compressible Euler equation require the additional analysis of nonlinear wave equations generating sound waves. An existence theory is developed by a specially chosen parabolic regularization together with the vanishing viscosity method. The artificial parabolic term is chosen so as to be asymptotically consistent with the Euler equations in the limit of zero viscosity. Having solutions for the positive surface tension problem, we proceed to obtain a priori estimates which are independent of the surface tension parameter. This requires choosing initial data which satisfy the Taylor sign condition. By passing to the limit of zero surface tension, we prove the well-posedness of the compressible Euler system without surface on the free-boundary, and without derivative loss.

Contents

1. Introduction 2
2. Preliminaries 7
3. A parabolic κ-approximation of the surface tension problem (1.7) 13
4. A priori estimates for the κ-problem (3.1) 15
5. Construction of solutions to the κ-problem (3.1) 26
6. Well-posedness of the surface tension problem (1.7) 39
7. The asymptotic limit as surface tension tends to zero 41
Appendix A. Constructing C^∞-class initial data 57
Appendix B. Solutions to the μ-problem (5.52) 63
Appendix C. Equivalence of the $\kappa\epsilon$-problem (5.6) and the heat-type $\kappa\epsilon$-problem (5.49) 68
List of Notation 71
References 72

1991 *Mathematics Subject Classification.* 35L65, 35L70, 35L80, 35Q35, 35R35, 76B03.

Key words and phrases. Compressible flow, vacuum, free boundary problems, surface tension.

1. Introduction

1.1. **The compressible Euler equations in Eulerian variables.**

The compressible Euler equations with moving free-boundary are given by the following system:

$$\begin{aligned}
\partial_t(\rho u) + \operatorname{div}(\rho u \otimes u + p\,\mathrm{Id}) &= 0 \qquad \text{in } \Omega(t), && (1.1a)\\
\partial_t \rho + \operatorname{div}(\rho u) &= 0 \qquad \text{in } \Omega(t), && (1.1b)\\
p &= \sigma H(t) \quad \text{on } \Gamma(t), && (1.1c)\\
\mathcal{V}(\Gamma(t)) &= u \cdot n(t) && (1.1d)\\
(u, \rho) &= (u_0, \rho_0) \quad \text{on } \Omega(0), && (1.1e)\\
\Omega(0) &= \Omega\,, && (1.1f)
\end{aligned}$$

where $\Omega(t)$ denotes an open and bounded subset of $\mathbb{R}^3$, $\Gamma(t) = \partial\Omega(t)$ is the moving free-boundary, and $t \in [0, T]$ denotes time. We use the notation $\mathcal{V}(\Gamma(t))$ for the normal velocity of boundary $\Gamma(t)$, which is equal to the normal component of the fluid velocity $u \cdot n$, where $n(t)$ is the outward-pointing unit normal to $\Gamma(t)$, $u = (u_1, u_2, u_3)$ denotes the velocity field, p denotes the pressure, and ρ denotes the density.

The first two equation are conservation laws for momentum and mass. The boundary condition (1.1c) is often referred to as the Laplace-Young condition, stating that the fluid stress is proportional to the mean curvature $H(t)$ of the moving surface, the proportionality constant defining the surface tension parameter σ. The last two equations provide the initial conditions for the dynamics.

In order to model the motion of a compressible liquid, we use the equation-of-state given by Courant and Friedrichs [8] as

$$p(x,t) = \alpha \rho(x,t)^\gamma - \beta \quad \text{for } \gamma > 1, \tag{1.2}$$

where $\alpha > 0$ and $\beta > 0$. For convenience, we set $\alpha = 1$. [1]

Using the equation of state (1.2), the momentum equations (1.1a) and Laplace-Young boundary condition (1.1c) are equivalently written as

$$\begin{aligned}
\rho[\partial_t u + (u \cdot D)u] + D\rho^\gamma &= 0 \qquad \text{in } \Omega(t), && (1.3a)\\
\rho^\gamma &= \beta + \sigma H \quad \text{on } \Gamma(t). && (1.3b)
\end{aligned}$$

We assume that the initial density function is strictly positive and that

$$\rho_0 \geq \lambda > 0 \text{ in } \overline{\Omega}.$$

In the absence of surface tension, we further require the initial pressure function p_0 to satisfy the *Taylor sign condition* (see, for example, [30] and [26]), given by

$$0 < \nu \leq -\frac{\partial p_0}{\partial N} \text{ on } \Gamma\,,$$

where N denotes the outward unit normal to Γ. This is equivalent to

$$0 < \nu \leq -\frac{\partial \rho_0^\gamma}{\partial n} \text{ on } \Gamma\,. \tag{1.4}$$

1.2. **Prior results on the Euler equations with moving free-boundary.**

---

[1] Using (1.2), liquid water is modeled using the values $\gamma = 7$, $\alpha = 3001$ and $\beta = 3000$.

1.2.1. *The incompressible setting.* There has been a recent explosion of interest in the analysis of the free-boundary incompressible Euler equations, particularly in irrotational form, that has produced a number of different methodologies for obtain- ing a priori estimates. The accompanying existence theories have relied mostly on the NashMoser iteration to deal with derivative loss in linearized equations when arbitrary domains are considered, or on complex analysis tools for the irrotational problem with infinite depth. We refer the reader to [3, 11, 14, 21, 22, 27, 35, 36, 39] for a partial list of papers on this topic.

1.2.2. *The compressible setting.* The mathematical analysis of moving hypersurfaces in the multidimensional compressible Euler equations began with the existence and stability of the shock-front solution initiated in [24] and extensively studied by [16–18, 25] (see the references in these articles for a thorough review of the literature in this area.) More delicate than the non-characteristic case of the shock-front solution, the characteristic boundary case is encountered in the study of vortex sheets or current vortex sheets. This class of problems has been studied by [4, 6, 7, 32, 34] (and see the references therein); the linearization of the vortex-sheet problem produces derivative loss, similar to that experienced by many authors in the incompressible flow setting (both irrotational flows and flows with vorticity).

The problem of the expansion of a compressible gas with the so-called physical vacuum singularity has been studied in [9, 12, 13, 19, 20], and is degenerate because of the vanishing of the density function on the moving free-boundary.

For the model of a compressible liquid, considered in this paper, the Euler equations are uniformly hyperbolic thanks to the equation-of-state (1.2). In the absence of surface tension, an existence theory was given in [23] using Lagrangian coordinates and a Nash-Moser construction, but the estimates had derivative loss. Using the theory of symmetric hyperbolic systems, the paper [33] gave a different proof for the existence of solutions, also with derivative loss. We prove well-posedness for the motion of a compressible liquid with and without surface tension, and with no derivative loss. We also establish asymptotic limit of zero surface tension.

1.3. **Fixing the domain and Lagrangian variables.** To transform the system (1.1) into Lagrangian variables, we let $\eta(x,t)$ denote the flow of a fluid particle x at time t. Thus,

$$\partial_t \eta = u \circ \eta \text{ for } t > 0 \quad \text{and} \quad \eta(x,0) = x$$

where $\circ$ denotes composition, so that $[u \circ \eta](x,t) = u(\eta(x,t),t)$. The *flow map* η induces the following Lagrangian variables on Ω:

$$\begin{aligned} v &= u \circ \eta && (\textit{Lagrangian velocity}), \\ f &= \rho \circ \eta && (\textit{Lagrangian density}), \\ A &= [D\eta]^{-1} && (\textit{inverse of the deformation tensor}), \\ J &= \det D\eta && (\textit{Jacobian determinant}), \\ a &= JA && (\textit{cofactor matrix of the deformation tensor}). \end{aligned}$$

Using the notation defined below in Section 2.1.2, the Lagrangian version of equations (1.1) on the fixed domain Ω is given by

$$f v_t^i + A_i^k f^\gamma,_k = 0 \qquad \text{in } \Omega \times (0,T], \tag{1.5a}$$

$$f_t + f A_i^j v^i,_j = 0 \qquad \text{in } \Omega \times (0,T], \tag{1.5b}$$

$$f^\gamma = \beta + \sigma H(\eta) \qquad \text{on } \Gamma \times (0,T], \tag{1.5c}$$

$$(\eta, v, f)|_{t=0} = (e, u_0, \rho_0) \qquad \text{on } \Omega, \tag{1.5d}$$

where $\Gamma = \partial\Omega$ and $e(x) = x$ denotes the identity map on Ω.

Since $J_t = a^j_i v^i,_j$ by (2.5), we multiply (1.5b) by J and integrate in time for the identity

$$f = \rho_0 J^{-1}. \tag{1.6}$$

For $\sigma \geq 0$ and $\gamma = 2$, we use the identity (1.6) for the Lagrangian density f and equivalently write the compressible Euler equations (1.5) as

$$\rho_0 v^i_t + a^k_i(\rho_0^2 J^{-2}),_k = 0 \qquad \text{in } \Omega \times (0,T], \tag{1.7a}$$

$$\rho_0^2 J^{-2} = \beta + \sigma H(\eta) \qquad \text{on } \Gamma \times (0,T], \tag{1.7b}$$

$$(\eta, v)|_{t=0} = (e, u_0) \qquad \text{on } \Omega. \tag{1.7c}$$

For $\sigma > 0$, we shall refer to the equations (1.7) as the *surface tension problem.*

For reference, we record that the momentum equations (1.7a) are equivalent to

$$v^i_t + 2A^k_i(\rho_0 J^{-1}),_k = 0 \qquad \text{in } \Omega \times (0,T]. \tag{1.8}$$

Remark 1. The identity (1.6) establishes that in Lagrangian variables the initial density ρ_0 is a parameter in the system of compressible Euler equations.

1.4. **The higher-order energy functions** $E(t)$ **and** $\mathcal{E}(t)$**.**

While the physical energy $\int_\Omega[\rho_0 \frac{1}{2}|v|^2 + \rho_0^2 J^{-1} + \beta J] + \sigma\mathcal{A}(t)$, $\mathcal{A}(t)$ denoting the surface area of $\Gamma(t)$, is a conserved quantity, it is far too weak for the energy estimates methodology that we employ. We instead define the higher-order energy functions $E(t)$ and $\mathcal{E}(t)$ to respectively correspond with the surface tension problem and the zero surface tension limit. Although neither $E(t)$ or $\mathcal{E}(t)$ is conserved, we will establish that each of $\sup_{t\in[0,T]} E(t)$ and $\sup_{t\in[0,T]} \mathcal{E}(t)$ is bounded on a sufficiently small time-interval of existence $[0,T]$.

1.4.1. *The higher-order energy function for* $\sigma > 0$*.* We define the energy function $E(t)$ as

$$E(t) = 1 + \sum_{a=0}^{5} \|\partial_t^a \eta(t)\|^2_{5-a} + |v_{ttt} \cdot n(t)|^2_1 + \sum_{a=0}^{2} |\bar\partial^2 \partial_t^a v \cdot n(t)|^2_{2.5-a}. \tag{1.9}$$

We let $M_0 \geq 0$ denote a generic constant given by a polynomial function P of $E(0)$:

$$M_0 = P\left(E(0)\right). \tag{1.10}$$

1.4.2. *The higher-order energy function for* $\sigma = 0$*.* We define the energy function $\mathcal{E}(t)$ as

$$\mathcal{E}(t) = 1 + \sum_{a=0}^{7} \|\partial_t^a \eta(t)\|^2_{4.5-\frac{1}{2}a} + \sum_{a=0}^{5} \|\partial_t^a J(t)\|^2_{4.5-\frac{1}{2}a} + \|\partial_t^6 J(t)\|^2_1. \tag{1.11}$$

We let $\mathcal{M}_0 \geq 0$ denote a generic constant given by a polynomial function P of $\mathcal{E}(0)$:

$$\mathcal{M}_0 = P\left(\mathcal{E}(0)\right). \tag{1.12}$$

Section 2.1 explains the notation in (1.9) and (1.11).

Remark 2. Corresponding with $\sigma = 0$, time-derivatives in the higher-order energy function $\mathcal{E}(t)$ scale like one-half a space-derivative. Since the scalar product of the momentum equations (1.7a) and the tangential vector $\eta,_\alpha$ yields the identity $\rho_0 v_t \cdot \eta,_\alpha = -J(\rho_0^2 J^{-2}),_\alpha$ in Ω, the Laplace-Young boundary condition (1.7b) with $\sigma = 0$ provides that

$$v_t \cdot \eta,_\alpha = 0 \quad \text{on } \Gamma.$$

Differentiating this identity with respect to time shows that ∂_t scales like $(\partial_x)^{1/2}$ in the zero surface tension limit of (1.7).

1.5. **The initial data (ρ_0, u_0, Ω).** We assume that ρ_0 and u_0 are given and sufficiently smooth. We require that the initial density ρ_0 satisfy

$$\rho_0 \geq 2\lambda > 0 \quad \text{in } \overline{\Omega}. \tag{1.13}$$

Using the identities $\eta(0) = e$ and $J_t = a^s_r v^r,_s$, we let J_1 be defined by

$$J_1 = \partial_t(J^{-2})(0) = -2 \operatorname{div} u_0.$$

We let v_1 and v_2 be the vectors respectively given by

$$v_1^i = -2\partial_i \rho_0 \quad \text{and} \quad v_2^i = -\rho_0^{-1}[\partial_i(\rho_0^2 J_1) + \partial_t a^k_i(0)(\rho_0^2),_k\,],$$

where $\partial_t a(0)$ is a smooth function of Du_0. We define, as a function of ρ_0 and u_0,

$$J_a = \partial_t^a (J^{-2})|_{t=0} \quad \text{for } a = 0, 1, 2, 3.$$

1.5.1. *The case of positive surface tension.* For $\sigma > 0$, we assume that Ω is an H^5-class domain and that ρ_0 and u_0 are in $H^4(\Omega)$. We define, as a function of ρ_0 and u_0,

$$H_a = \partial_t^a H(\eta)|_{t=0} \quad \text{for } a = 0, 1, 2, 3.$$

We require that the initial data satisfy the following compatibility conditions:

$$\rho_0^2 J_a = \partial_t^a \beta + \sigma H_a \quad \text{on } \Gamma \text{ for } a = 0, 1, 2, 3. \tag{1.14}$$

For β as in (1.2) and λ as in (1.13), we note that $a = 0$ in (1.14) yields

$$\sigma H_0 \geq 4\lambda^2 - \beta.$$

1.5.2. *The case of zero surface tension.* For $\sigma = 0$, we assume that Ω is an $H^{4.5}$-class domain and that ρ_0 is in $H^{4.5}(\Omega)$ and u_0 is in $H^4(\Omega)$.

We require that ρ_0 and Ω satisfy the Taylor sign condition:

$$0 < 2\nu \leq -\frac{1}{\sqrt{g}} N^j a^j_i a^k_i (\rho_0^2 J^{-2}),_k \Big|_{t=0} \quad \text{on } \Gamma. \tag{1.15}$$

We require that the initial data satisfy the following compatibility conditions:

$$\rho_0^2 J_a = \partial_t^a \beta \quad \text{on } \Gamma \text{ for } a = 0, \ldots, 6. \tag{1.16}$$

Remark 3. The twice-mean-curvature function $H(\eta)$ is not present in the zero surface tension limit of (1.7). Accordingly, there is no restriction on the curvature of the initial surface Γ.

1.6. **Main Results.** The main results of this paper are the existence and uniqueness of solutions to the surface tension problem and its zero surface tension limit.

Theorem 1.1 (Existence and uniqueness for $\sigma > 0$)**.** *Suppose that the initial data* (ρ_0, u_0, Ω) *verify*

1. $M_0 = P(E(0)) < \infty$,
2. *the lower-bound condition* (1.13), *and*
3. *the compatibility conditions* (1.14).

Then for some $T > 0$, *there exists a solution to* (1.7) *on the time-interval* $[0, T]$ *such that* $\rho(t) \geq \lambda$ *in* $\overline{\Omega}(t)$, $\sigma H(t) > -\beta$ *on* $\Gamma(t)$, *and*

$$\sup_{t \in [0,T]} E(t) \leq 2M_0.$$

Furthermore, the solution is unique if the initial data is such that

$$\sum_{a=0}^{6} \|\partial_t^a \eta(0)\|^2_{6-a} + |v_{tttt} \cdot n(0)|^2_1 + \sum_{a=0}^{3} |\bar{\partial}^2 \partial_t^a v \cdot n(0)|^2_{3.5-a} < \infty.$$

Theorem 1.2 (The zero surface tension limit)**.** *Suppose that the initial data* (ρ_0, u_0, Ω) *verify*

1. $\mathcal{M}_0 = P(\mathcal{E}(0)) < \infty$,
2. *the lower-bound condition* (1.13),
3. *the Taylor sign condition* (1.15), *and*
4. *the compatibility conditions* (1.16).

Then for some $T > 0$, *there exists a solution to* (1.7) *with* $\sigma = 0$ *on the time-interval* $[0, T]$ *such that* $\rho(t) \geq \lambda$ *in* $\overline{\Omega}(t)$, $-\frac{\partial \rho^2(t)}{\partial n(t)} \geq \nu$ *on* $\Gamma(t)$, *and*

$$\sup_{t \in [0,T]} \mathcal{E}(t) \leq 2\mathcal{M}_0.$$

Furthermore, the solution is unique if the initial data is such that

$$\sum_{a=0}^{9} \|\partial_t^a \eta(0)\|^2_{5.5-\frac{1}{2}a} + \sum_{a=0}^{7} \|\partial_t^a J(0)\|^2_{5.5-\frac{1}{2}a} + \|\partial_t^8 J(0)\|_1^2 < \infty.$$

Remark 4. The proofs of Theorems 1.1 and 1.2 do not rely on our choice of $\gamma = 2$ and as such are valid for general $\gamma > 1$, since by introducing new enthalpy-type variables for $\rho_0^{\gamma-1}$ and $J^{\gamma-1}$ in (1.5), we can reduce the case of general $\gamma > 1$ to the case that $\gamma = 2$.

1.7. **Structure of the proofs of Theorems 1.1 and 1.2.**

1.7.1. *Existence for the surface tension problem* (1.7). An existence theory is obtained via the vanishing viscosity method, but with a very special choice of artificial viscosity which does not alter the transport-type structure of vorticity. This is introduced in Section 3. The Euler equations (1.7a) yield

$$\operatorname{curl}_\eta v_t = 0,$$

where the Lagrangian curl operator curl_η is defined below in Section 2.1.3. By defining a parabolic approximation of the Euler equations (1.7a) which preserves the homogeneous vorticity equation, we ensure that the vorticity estimates for the approximate problem are independent of the parabolic approximation parameter κ. The parabolic approximate κ-problem defined in Section 3 maintains a homogeneous vorticity equation.

The structure of the Euler equation (1.7a) shows that an a priori estimate for v_t provides an estimate for the gradient of J, which then yields an estimate for the gradient of $\operatorname{div} \eta$. As such, yet another constraint on the choice of parabolic regularization operator is that its presence must still permit estimates for $D \operatorname{div} \eta$ whenever estimates for v_t are known. We choose our κ-parabolic operator so that the parabolically-regularized momentum equations have the form

$$f + \kappa f_t = g,$$

for f equalling the gradient of J. Such a structure allows us to obtain κ-independent estimates for f, given estimates for g. We must further add parabolic regularization to the surface tension boundary conditions (1.7b) which also has the same structure, in order to infer the maximal regularity of the free-boundary.

Our strategy is to obtain energy estimates for the highest number of time-derivatives in the problem, and then use the structure of the momentum equations to infer estimates for the divergence. In conjunction with curl-estimates and boundary regularity, we boot-strap the full regularity of the velocity, its time-derivatives, and the flow map η.

In Section 4, we perform energy estimates in the fourth (highest) time-differentiated approximate κ-problem. From these energy estimates, we are able to close estimates for

$v_{tttt}(t)$ in $L^2(\Omega)$ and $v_{ttt}(t)$ in $H^1(\Omega)$. We then infer the optimal regularity of v_{tt} and v_t. This, in turn, enables us to get the optimal regularity for v and η. All estimates are found to be independent of the artificial viscosity parameter κ.

In Section 5, we construct smooth solutions the approximate κ-problem. We utilize (the Lagrangian variables version of) the basic vector identity $-\Delta = \operatorname{curl}\operatorname{curl} - D\operatorname{div}$ in order to replace gradient of the divergence of v with the Laplacian of v, modulo lower-order terms. Additionally, we use that a sufficiently regular vector $\xi \in \mathbb{R}^3$ satisfies

$$N^j A^j_r A^k_r \xi,_k = \sqrt{g} J^{-1} (\operatorname{curl}_\eta \xi) \times n + \sqrt{g} J^{-1} (\operatorname{div}_\eta \xi) n + b(\xi, \eta),$$

where $b(\xi, \eta)$ is a first-order differential boundary operator with respect to ξ and a second-order differential boundary operator with respect to η. Since the regularity of the flow map η is dictated by the regularity of v in our construction scheme, we employ a horizontal convolution-by-layers operator (first introduced in [11]) in order to view $b(v, \eta)$ as a lower-order term; with the horizontal convolution-by-layers approximation in place, solutions can be found via the existence theory for uniformly parabolic second-order equations. We must then find estimates for such solutions which are indeed independent of the convolution parameter and pass to the limit.

In Section 6, we use the κ-independent a priori estimates established in Section 4 and the construction of solutions to our approximate κ-problem in Section 5 to establish the existence and uniqueness of solutions to the surface tension problem (1.7).

1.7.2. *The zero surface tension limit.* In Section 7, we establish our existence theory for the zero surface tension limit of (1.7) via σ-independent a priori estimates. For initial data satisfying the Taylor sign condition (1.15), we have that solutions to the surface tension problem (1.7) satisfy

$$0 < \nu \int_\Gamma |\eta \cdot n|^2 \leq - \int_\Gamma \frac{1}{\sqrt{g}} N^j a^j_\ell a^k_\ell (\rho_0^2 J^{-2}),_k |\eta \cdot n|^2,$$

on a sufficiently small time-interval $[0, T]$.

2. Preliminaries

2.1. **Notation.**

2.1.1. *The three-dimensional gradient vector.* Throughout this paper the symbol D will be used to denote the three-dimensional gradient vector

$$D = \left(\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \frac{\partial}{\partial x_3} \right).$$

2.1.2. *Notation for partial differentiation and Einstein's summation convention.* The kth partial derivative of F will be denoted by $F,_k = \frac{\partial F}{\partial x_k}$. Repeated Latin indices i, j, k, etc., are summed from 1 to 3, and repeated Greek indices α, β, γ, etc., are summed from 1 to 2. For example, $F,_{ii} = \sum_{i=1}^3 \frac{\partial^2 F}{\partial x_i \partial x_i}$, and $F^i,_\alpha I^{\alpha\beta} G^i,_\beta = \sum_{i=1}^3 \sum_{\alpha=1}^2 \sum_{\beta=1}^2 \frac{\partial F^i}{\partial x_\alpha} I^{\alpha\beta} \frac{\partial G^i}{\partial x_\beta}$.

2.1.3. *The divergence and curl operators.* We use the notation $\operatorname{div} u$ to denote the divergence of a vector field u on Ω:

$$\operatorname{div} u = u^1,_1 + u^2,_2 + u^3,_3 ,$$

and we use the notation $\operatorname{curl} u$ to denote the curl of a vector u on Ω:

$$\operatorname{curl} u = \left(u^3,_2 - u^2,_3 , u^1,_3 - u^3,_1 , u^2,_1 - u^1,_2 \right).$$

We recall that the permutation symbol ε_{ijk} is defined by

$$\varepsilon_{ijk} = \begin{cases} 1, & \text{even permutation of } \{1,2,3\}, \\ -1, & \text{odd permutation of } \{1,2,3\}, \\ 0, & \text{otherwise.} \end{cases}$$

This allows for the curl of a vector field u to be expressed as $\operatorname{curl} u = \varepsilon_{\cdot jk} u^k,_j$. Letting $v = u(\eta)$ for a given flow map η, we use the notation $\operatorname{div}_\eta v$ to denote the Lagrangian divergence of v on Ω:

$$\operatorname{div}_\eta v = A^s_r v^r,_s ,$$

and we use the notation $\operatorname{curl}_\eta v$ to denote the Lagrangian curl of v on Ω:

$$\operatorname{curl}_\eta v = \varepsilon_{\cdot jk} A^s_j v^k,_s .$$

2.1.4. *The scalar- and cross-product of vectors in* $\mathbb{R}^3$. Let u and v be vectors in $\mathbb{R}^3$. The scalar-product of u and v, denoted $u \cdot v$, is defined as

$$u \cdot v = u^1 v^1 + u^2 v^2 + u^3 v^3. \tag{2.1}$$

The cross-product of u and v, denoted $u \times v$, is defined as

$$u \times v = \varepsilon_{\cdot jk} u^j v^k. \tag{2.2}$$

2.1.5. *Local coordinates near* Γ. We let $\Omega \subset \mathbb{R}^3$ denote an open, bounded subset of class H^s for $s \geq 4$, and we let $\{U_l\}_{l=1}^K$ denote an open covering of $\Gamma = \partial\Omega$, such that for each $l \in \{1, 2, \ldots, K\}$, with

$$\begin{aligned} \mathcal{B}_l &= B(0, r_l), \text{ denoting the open ball of radius } r_l \text{ centered at the origin,} \\ \mathcal{B}_l^+ &= \mathcal{B}_l \cap \{x_3 > 0\}, \\ \mathcal{D}_l &= \mathcal{B}_l \cap \{x_3 = 0\}, \end{aligned}$$

there exists an H^s-class chart θ_l satisfying

$$\begin{aligned} \theta_l \colon \mathcal{B}_l \to U_l \ &\text{ is an } H^s \text{ diffeomorphism,} \\ \theta_l(\mathcal{B}_l^+) &= U_l \cap \Omega, \\ \theta_l(\mathcal{D}_l) &= U_l \cap \Gamma. \end{aligned}$$

For $L > K$, we let $\{U_l\}_{l=K+1}^L$ denote a family of open balls of radius r_l properly contained



FIGURE 1. Indexing convention for the open cover $\{U_l\}_{l=1}^L$ of Ω.

in Ω such that $\{U_l\}_{l=1}^L$ is an open cover of Ω. We let

$\{\xi_l\}_{l=1}^L$ denote a C^∞ partition-of-unity subordinate to the open covering of Ω.

2.1.6. *Tangential derivatives.* On each $U_l \cap \Omega$, for $1 \leq l \leq L$, we let $\bar\partial_l$ denote the *local tangential-derivative*. That is, for a differentiable function f on Ω, the αth component of the local tangential-derivative of f is defined in $U_l \cap \Omega$ by

$$\bar\partial_{l,\alpha} f = \left(\frac{\partial}{\partial x_\alpha}[f \circ \theta_l]\right) \circ \theta_l^{-1} = \left((Df \circ \theta_l)\frac{\partial \theta_l}{\partial x_\alpha}\right) \circ \theta_l^{-1},$$

where for $K+1 \leq l \leq L$, we have set θ_l to be the identity map e.

We let $\bar\partial$ denote the *tangential-derivative* whose αth component is given by

$$\bar\partial_\alpha = \sum_{l=1}^L \xi_l \bar\partial_{l,\alpha}.$$

We use $\bar\partial_\alpha f$ or $f,_\alpha$ to denote the components of the tangential-derivative of f.

2.1.7. *Geometry of the moving surface* $\Gamma(t)$. The vectors $\eta,_\alpha$ for $\alpha = 1, 2$, span the tangent space to the moving surface $\Gamma(t) = \eta(\Gamma)$ in $\mathbb{R}^3$. The *surface metric* g on $\Gamma(t)$ has components

$$g_{\alpha\beta} = \eta,_\alpha \cdot \eta,_\beta\,.$$

We let g_0 denote the surface metric of the initial surface Γ. The components of the inverse metric $[g]^{-1}$ are denoted by $g^{\alpha\beta}$. We use $\sqrt{g}$ to denote $\sqrt{\det g}$; we note that $\sqrt{g} = |\eta,_1 \times \eta,_2|$, so that $n = [\eta,_1 \times \eta,_2]/\sqrt{g}$. Equivalently,

$$\sqrt{g} n = \eta,_1 \times \eta,_2\,. \tag{2.3}$$

The Laplace-Beltrami operator Δ_g is defined on Γ as

$$\Delta_g = \sqrt{g}^{-1} \bar\partial_\alpha[\sqrt{g} g^{\alpha\beta} \bar\partial_\beta].$$

2.1.8. *Sobolev spaces on* Ω. For integers $k \geq 0$ and a smooth, open domain Ω of $\mathbb{R}^3$, we define the Sobolev space $H^k(\Omega)$ ($H^k(\Omega;\mathbb{R}^3)$) to be the closure of $C^\infty(\overline{\Omega})$ ($C^\infty(\overline{\Omega};\mathbb{R}^3)$) in the norm

$$\|u\|_k^2 = \sum_{|a|\leq k} \int_\Omega |D^a u(x)|^2\,,$$

for a multi-index $a \in \mathbb{Z}_+^3$, with the convention that $|a| = a_1 + a_2 + a_3$. For real numbers $s \geq 0$, the Sobolev spaces $H^s(\Omega)$ and the norms $\|\cdot\|_s$ are defined by interpolation. We will write $H^s(\Omega)$ instead of $H^s(\Omega;\mathbb{R}^3)$ for vector-valued functions. We use $H^s(\Omega)'$ to denote the dual space of $H^s(\Omega)$.

2.1.9. *Sobolev spaces on* Γ. For functions $u \in H^k(\Gamma)$, $k \geq 0$, we set

$$|u|_k^2 = \sum_{|a|\leq k} \int_\Gamma \left|\bar\partial^a u(x)\right|^2\,,$$

for a multi-index $a \in \mathbb{Z}_+^2$. For real $s \geq 0$, the Hilbert space $H^s(\Gamma)$ and the boundary norm $|\cdot|_s$ is defined by interpolation. The negative-order Sobolev spaces $H^{-s}(\Gamma)$ are defined via duality. That is, for real $s \geq 0$,

$$H^{-s}(\Gamma) = H^s(\Gamma)'.$$

Remark 5. Throughout this paper, we suppress the Euclidean measure dx by letting $\int_\Omega$ represent $\int_\Omega dx$. Similarly, the notation $\int_\Gamma$ represents $\int_\Gamma dS_0$ where $dS_0 = \sqrt{g_0}dx_1dx_2$ is the surface measure of the initial surface Γ. Equally, the time integral $\int_0^t$ should be read as $\int_0^t ds$.

Remark 6. We let $|\cdot|_{s,\mathcal{D}_l}$ denote the $H^s(\mathcal{D}_l)$-norm.

2.2. **Differentiation and geometric identities and properties.**

2.2.1. *An identity for the Jacobian determinant J.* With $\dim\Omega = 3$, we have that the Jacobian determinant J is written as

$$J = \frac{1}{\dim\Omega}\left[a^s_r \eta^r,_s\right], \tag{2.4}$$

which follows from the definition of the cofactor matrix a in Section 1.3.

2.2.2. *Time-differentiating the Jacobian determinant J and the cofactor matrix a.* We record the following basic differentiation formulas:

$$\partial_t J = a^s_r v^r,_s\,, \tag{2.5}$$

$$\partial_t a^k_i = J^{-1}[a^s_r a^k_i - a^s_i a^k_r]v^r,_s\,. \tag{2.6}$$

Using (2.5) and (2.6) and the fact that $a = JA$, we have that

$$\partial_t A^k_i = -A^s_i A^k_r v^r,_s\,. \tag{2.7}$$

We note that the time-differentiation formulas (2.5)–(2.7) at once become formulas for tangential-differentiation by replacing $v^r,_s$ with $\bar\partial\eta^r,_s$ in the right-hand sides of (2.5)–(2.7).

The formulas (2.5) and (2.6) imply the following scaling relations:

$$\begin{aligned} \partial_t J &\sim aDv, & \partial_t a &\sim a^2 Dv,\\ \partial_t^2 J &\sim a^2(Dv)^2 + aDv_t, & \partial_t^2 a &\sim a^3(Dv)^2 + a^2 Dv_t. \end{aligned}$$

These scaling relations are particularly useful when estimating error-terms.

2.2.3. *The Piola identity.* Columns of every cofactor matrix are divergence-free. Thus,

$$a^k_i,_k = 0. \tag{2.8}$$

2.2.4. *Relating the normal vectors of Γ and $\Gamma(t)$.* With N as the outward unit normal to the reference surface Γ, the outward-normal direction of the moving surface $\Gamma(t)$ is

$$a^k_i N^k = |\eta,_1 \times \eta,_2|n^i.$$

The identity $\sqrt{g} = \sqrt{\det g} = |\eta,_1 \times \eta,_2|$ implies that

$$a^k_i N^k = \sqrt{g}n^i. \tag{2.9}$$

2.2.5. *Derivatives of the inverse metric $g^{\alpha\beta}$, Jacobian $\sqrt{g}$ and unit normal n.* A tangential-derivative of the inverse metric $g^{\alpha\beta}$, Jacobian determinant $\sqrt{g}$ and moving outward normal unit vector n are given by the formulas

$$\bar\partial g^{\alpha\beta} = -g^{\alpha\mu}\bar\partial g_{\mu\nu} g^{\nu\beta}, \tag{2.10a}$$

$$\bar\partial\sqrt{g} = \tfrac{1}{2}\sqrt{g} g^{\mu\nu}\bar\partial g_{\mu\nu}, \tag{2.10b}$$

$$\bar\partial n = -g^{\gamma\delta}[\bar\partial\eta,_\delta \cdot n]\eta,_\gamma\,. \tag{2.10c}$$

Also,

$$\partial_t g^{\alpha\beta} = -g^{\alpha\mu}\partial_t g_{\mu\nu} g^{\nu\beta}, \tag{2.11a}$$

$$\partial_t\sqrt{g} = \tfrac{1}{2}\sqrt{g} g^{\mu\nu}\partial_t g_{\mu\nu}, \tag{2.11b}$$

$$\partial_t n = -g^{\gamma\delta}[v,_\delta \cdot n]\eta,_\gamma\,. \tag{2.11c}$$

Remark 7. The right-hand side of (2.10c) and (2.11c) is a vector that is tangent to the embedded surface.

2.2.6. *Relating the Laplace-Beltrami operator Δ_g to the unit normal n.* With the formulas (2.10a) and (2.10b), we have that the Laplace-Beltrami operator $\Delta_g = \sqrt{g}^{-1}\bar\partial_\alpha[\sqrt{g}g^{\alpha\beta}\bar\partial_\beta]$ applied to the particle flow η decomposes into normal and tangential components as

$$\sqrt{g}\Delta_g(\eta) = \sqrt{g}\,\underbrace{(g^{\alpha\beta}\eta,_{\beta\alpha} - \eta,_{\mu\alpha}\cdot\eta,_\nu\, g^{\alpha\mu}g^{\nu\beta}\eta,_\beta\,)}_{g^{\alpha\beta}[\eta,_{\alpha\beta}\cdot n]n} + \sqrt{g}\eta,_{\mu\alpha}\cdot\eta,_\nu\,(g^{\alpha\beta}g^{\mu\nu} - g^{\alpha\mu}g^{\nu\beta})\eta,_\beta\,.$$

We therefore have the identity

$$\sqrt{g}\Delta_g(\eta) = \sqrt{g}g^{\alpha\beta}[\eta,_{\alpha\beta}\cdot n]n. \tag{2.12}$$

For reference, we recall the identity $\Delta_g(\eta) = -H(\eta)n$.

2.3. **Two identities for the Euler equations in Lagrangian variables.**

2.3.1. *The Lagrangian vorticity equation.* With the operator curl_η defined in Section 2.1.3,

$$\operatorname{curl}_\eta v_t = 0 \quad \text{in } \Omega. \tag{2.13}$$

The identity (2.13) is obtained by taking the Lagrangian curl of the Euler equations (1.8).

2.3.2. *A tangential identity for v_t on the boundary Γ.* Setting $\sigma = 1$,

$$v_t\cdot\eta,_\alpha = f^{-1}\bar\partial_\alpha(g^{\mu\nu}[\eta,_{\mu\nu}\cdot n]) \quad \text{on } \Gamma. \tag{2.14}$$

The identity (2.14) is established using the Euler equations (1.8), the Laplace-Young boundary condition (1.7b), and the formula (2.12).

2.4. **General inequalities.**

2.4.1. *Trace estimates.* For $s > \frac{1}{2}$ and some constant C independent of $w \in H^s(\Omega)$, the trace theorem [1] states that the trace of w is defined in $H^{s-\frac{1}{2}}(\Gamma)$ with the estimate

$$|w|_{s-\frac{1}{2}} \le C\|w\|_s.$$

Lemma 2.1. *Let $w \in L^2(0,T;H^1(\Omega)) \cap L^\infty(0,T;L^2(\Omega))$. Then*

$$\int_0^T |w|^2_{0.25} \le \delta\int_0^T \|w\|^2_1 + C_\delta\, T \sup_{t\in[0,T]}\|w(t)\|^2_0, \tag{2.15}$$

where the constant C_δ depends on $1/\delta > 0$.

Proof. Using interpolation and Young's inequality with $\delta > 0$, we have that

$$\int_0^T \|w\|_{0.75}^2 \le C \int_0^T \|w\|_0^{\frac{1}{2}} \|w\|_1^{\frac{3}{2}} \le \delta \int_0^T \|w\|_1^2 + C_\delta \, T \sup_{t\in[0,T]} \|w(t)\|_0^2 .$$

The proof is complete thanks to the trace theorem. ☐

Lemma 2.2 (Normal trace theorem)**.** *Let w be a vector field defined on Ω such that $\bar\partial w \in L^2(\Omega)$ and $\operatorname{div} w \in L^2(\Omega)$, and let N denote the outward unit normal vector to Γ. Then the normal trace $\bar\partial w \cdot N$ exists in $H^{-0.5}(\Gamma)$ with the estimate*

$$|\bar\partial w \cdot N|_{-0.5}^2 \le C \left[\|\bar\partial w\|_{L^2(\Omega)}^2 + \|\operatorname{div} w\|_{L^2(\Omega)}^2 \right], \tag{2.16}$$

for some constant C independent of w.

See [31] for the proof of Lemma 2.2 in the case that $\bar\partial w$ is replaced by w. For $\phi \in H^1(\Omega)$, $\int_\Gamma \bar\partial w \cdot n \phi dS = \int_\Omega \bar\partial w \cdot D\phi dx - \int_\Omega \operatorname{div} w \bar\partial \phi dx$ because we can integrate-by-parts with $\bar\partial$ without any boundary contributions. Thus, the identical proof given in [31] proves Lemma 2.2. Similarly, we have

Lemma 2.3 (Tangential trace theorem)**.** *Let w be a vector field defined on Ω such that $\bar\partial w \in L^2(\Omega)$ and $\operatorname{curl} w \in L^2(\Omega)$, and let τ_1, τ_2 denote the unit tangent vectors to Γ, so that any vector field u on Γ can be uniquely written as $u^\alpha \tau_\alpha$. Then the tangential trace $\bar\partial w \cdot \tau_\alpha$ exists in $H^{-0.5}(\Gamma)$ with the estimate*

$$|\bar\partial w \cdot \tau_\alpha|_{-0.5}^2 \le C \left[\|\bar\partial w\|_{L^2(\Omega)}^2 + \|\operatorname{curl} w\|_{L^2(\Omega)}^2 \right], \tag{2.17}$$

for some constant C independent of w.

See [5] for the proof of Lemma 2.3.

2.4.2. *An elliptic estimate which is independent of κ.* The following lemma is used to establish our κ-independent a priori estimates. The proof is given in Section 6 of [10].

Lemma 2.4. *Let $\kappa > 0$, $s \ge 0$ and $g \in L^\infty(0,T;H^s(\Omega))$ be given. Suppose that $f \in H^1(0,T;H^s(\Omega))$ satisfies*

$$f + \kappa f_t = g \quad \text{in } \Omega \times (0,T). \tag{2.18}$$

Then,

$$\|f\|_{L^\infty(0,T;H^s(\Omega))} \le C \max\{\|f(0)\|_s, \|g\|_{L^\infty(0,T;H^s(\Omega))}\}.$$

2.4.3. *A technical lemma.* The following technical lemma is established in [11].

Lemma 2.5. *There exists a constant C such that*

$$\|\bar\partial w\|_{H^{0.5}(\Omega)'} \le C \|w\|_{H^{0.5}(\Omega)} \quad \forall\ w \in H^{0.5}(\Omega).$$

2.4.4. *The Hodge decomposition elliptic estimates.* The following Hodge-type elliptic estimate is well-known and follows from the identity $-\Delta w = \operatorname{curl}\operatorname{curl} w - D \operatorname{div} w$, together with estimates divergence-form elliptic operators with Sobolev-class coefficients:

Proposition 2.1. *For an H^r-class domain Ω, $r \ge 3$, if $w \in L^2(\Omega;\mathbb{R}^3)$ with $\operatorname{curl} w \in H^{s-1}(\Omega;\mathbb{R}^3)$, $\operatorname{div} w \in H^{s-1}(\Omega)$, and $w \cdot N|_\Gamma \in H^{s-\frac{1}{2}}(\Gamma)$ for $1 \le s \le r$, then there exists a constant $C > 0$ depending only on Ω such that*

$$\|w\|_s \le C \left[\|w\|_0 + \|\operatorname{curl} w\|_{s-1} + \|\operatorname{div} w\|_{s-1} + |\bar\partial w \cdot N|_{s-\frac{3}{2}} \right],$$

where N denotes the outward unit-normal to Γ.

In fact, the well-known version of this elliptic estimate replaces $|\bar\partial w \cdot N|_{s-\frac{3}{2}}$ with $|w \cdot N|_{s-\frac{1}{2}}$, but this requires too much regularity for the unit normal N. It is easy to verify that having estimates for the divergence and curl of a vector field w only requires the slightly weaker norm $|\bar\partial w \cdot N|_{s-\frac{3}{2}}$ estimated in order to infer full regularity of w. Also, this elliptic estimate is usually stated for smooth domains; the Sobolev-class regularity follows from the Sobolev embedding theorem and somewhat standard elliptic estimates for second-order elliptic operators with Sobolev-class coefficients.

2.4.5. *A polynomial-type inequality.* For a constant $M \geq 0$, suppose $f\colon t \mapsto f(t) \geq 0$ continuously and satisfies

$$f(t) \leq M + t\, P(f(t)), \qquad t \geq 0, \tag{2.19}$$

where P denotes a generic polynomial function. Then for t taken sufficiently small,

$$f(t) \leq 2M.$$

3. A PARABOLIC κ-APPROXIMATION OF THE SURFACE TENSION PROBLEM (1.7)

In this section, we define a parabolic approximation of the surface tension problem (1.7), which we term the κ-problem. The κ-problem is defined by adding artificial viscosity terms to the Euler equations and the Laplace-Young boundary condition. The salient feature of the κ-problem is its compatibility with our energy-estimates methodology based on Proposition 2.1 in that (1) the transport structure of the Euler equations is maintained and (2) the momentum equations of the κ-problem are equivalently expressed in the form $f + \kappa f_t = g$ with f equalling the gradient of J. These structural properties of the κ-problem respectively yield the κ-independent curl- and divergence-estimates.

3.1. **Assuming C^∞-class initial data.**

In our construction of solutions to the surface tension problem (1.7), we assume that the initial data (ρ_0, u_0, Ω) is of C^∞-class and satisfy the conditions (1.13) and (1.14), as in Appendix A. Later, in Section 6.2, we will recover the optimal regularity of the initial data stated in Theorem 1.1.

3.2. **The parabolic approximation of the surface tension problem (1.7).**

We recall that an equivalent expression of the surface tension problem (1.7) is

$$\begin{aligned} v_t^i + 2A_i^k(\rho_0 J^{-1}),_k &= 0 && \text{in } \Omega \times (0,T], \\ \rho_0^2 J^{-2} &= \beta - \sigma g^{\alpha\beta} \eta,_{\alpha\beta} \cdot n && \text{on } \Gamma \times (0,T]. \end{aligned}$$

We have used the identity $H(\eta) = -g^{\alpha\beta}\eta,_{\alpha\beta} \cdot n$ in writing the boundary condition.

The variables in the following problem a priori depend on the parabolic parameter κ. To indicate this dependence, we place the symbol $\sim$ above each of the variables.

Definition 3.1 (The κ-problem)**.** *For $\kappa > 0$, we define $\tilde v$ as the solution of*

$$\tilde v_t^i + 2\tilde A_i^k(\rho_0 \tilde J^{-1}),_k - \kappa \tilde A_i^k(\rho_0 \tilde J_t),_k = 0 \qquad \text{in } \Omega \times (0, T_\kappa], \tag{3.1a}$$

$$\rho_0^2 \tilde J^{-2} - \kappa \rho_0^2 \tilde J^{-1} \tilde J_t = \tilde H_\kappa \qquad \text{on } \Gamma \times (0, T_\kappa], \tag{3.1b}$$

$$(\tilde\eta, \tilde v)|_{t=0} = (e, u_0) \quad \text{on } \Omega. \tag{3.1c}$$

The function $\tilde H_\kappa$ appearing in the right-hand side of (3.1b) *is defined as*

$$\tilde H_\kappa = \beta(t) - \sigma \tilde g^{\alpha\beta} \tilde\eta,_{\alpha\beta} \cdot \tilde n - \kappa \tilde g^{\alpha\beta} \tilde v,_{\alpha\beta} \cdot \tilde n, \tag{3.2}$$

where the function $\beta(t)$ appearing in the right-hand side of (3.2) *is defined as*

$$\beta(t) = \beta + \sum_{a=0}^{3} \frac{t^a}{a!}\partial_t^a \Big[\rho_0^2 \tilde{J}^{-2} - \beta + \sigma \tilde{g}^{\alpha\beta}\tilde{\eta},_{\alpha\beta}\cdot\tilde{n}\Big]|_{t=0} + \kappa \sum_{a=0}^{3} \frac{t^a}{a!}\partial_t^a \Big[- \rho_0^2 \tilde{J}^{-1}\tilde{J}_t + \tilde{g}^{\alpha\beta}\tilde{v},_{\alpha\beta}\cdot\tilde{n}\Big]|_{t=0}. \tag{3.3}$$

Remark 8. The particular artificial viscosity $-\kappa\tilde{A}_i^k(\rho_0\tilde{J}_t),_k$ appearing in (3.1a) preserves the transport structure of the Euler equations. In comparison, an artificial viscosity of the form $-\kappa\tilde{A}_r^j[\tilde{A}_r^k\tilde{v}^i,_k],_j$ would not preserve the transport structure of the Euler equations.

Remark 9. The initial data satisfy the compatibility conditions (1.14) so that definition (3.3) of $\beta(t)$ implies that $\beta(t)\to\beta$ in the limit as κ tends to zero. Formally, the κ-problem (3.1) is asymptotically consistent with the surface tension problem (1.7).

Remark 10. For $\phi \in L^2(0,T;H^1(\Omega))$ such that $\phi\cdot\tilde{n} \in L^2(0,T;H^1(\Gamma))$, the variational equation for the κ-problem (3.1) is

$$\int_0^T\int_\Omega \rho_0\tilde{v}_t\cdot\phi - \int_0^T\int_\Omega \rho_0^2\tilde{J}^{-2}\tilde{a}_i^k\phi^i,_k + \kappa\int_0^T\int_\Omega \rho_0\tilde{J}_t\tilde{a}_i^k(\rho_0\tilde{J}^{-1}\phi^i),_k + \int_0^T\int_\Gamma \beta(t)\sqrt{\tilde{g}}\phi\cdot\tilde{n} + \int_0^T\int_\Gamma [\sigma\tilde{\eta},_\beta + \kappa\tilde{v},_\beta]^i(\tilde{n}^i\sqrt{\tilde{g}}\tilde{g}^{\alpha\beta}\phi\cdot\tilde{n}),_\alpha = 0. \tag{3.4}$$

Since $\lim_{\kappa\to 0}\beta(t) = \beta$, we have that (3.4) with $\kappa = 0$ is the variational equation for the surface tension problem (1.7). Furthermore, for $\kappa > 0$ our choice of artificial viscosity $-\kappa\tilde{A}_i^k(\rho_0\tilde{J}_t),_k$ has the nice property of not introducing any nontrivial boundary integrals in the variational equation (3.4), whereas the use of the artificial viscosity $-\kappa\tilde{A}_r^j[\tilde{A}_r^k\tilde{v}^i,_k],_j$, for example, in (3.4) would require additional boundary conditions to account for the tangential components of $-\kappa N^j\tilde{A}_r^j\tilde{A}_r^k\tilde{v}^i,_k$.

3.3. **The constant-in-time vectors v_a for $a = 1, 2, 3, 4$.**

The vector field $\tilde{v}_t|_{t=0}$ is computed using the momentum equations (3.1a), as follows:

$$\tilde{v}_t|_{t=0} = \Big(\kappa\tilde{A}_\cdot^k(\rho_0\tilde{J}_t),_k - 2\tilde{A}_\cdot^k(\rho_0\tilde{J}^{-1}),_k\Big)|_{t=0} = D(\kappa\rho_0\operatorname{div}u_0 - 2\rho_0).$$

Similarly, for all $a\in\mathbb{N}$,

$$\partial_t^a\tilde{v}|_{t=0} = \frac{\partial^{a-1}}{\partial t^{a-1}}\Big(\kappa\tilde{A}_\cdot^k(\rho_0\tilde{J}_t),_k - 2\tilde{A}_\cdot^k(\rho_0\tilde{J}^{-1}),_k\Big)\Big|_{t=0} \quad \text{on } \Omega.$$

This formula makes it clear that each $\partial_t^a\tilde{v}|_{t=0}$ is a function of space-derivatives of the initial data u_0 and ρ_0. We define the constant-in-time vectors v_a as

$$\mathrm{v}_a = \frac{\partial^{a-1}}{\partial t^{a-1}}\Big(\kappa\tilde{A}_\cdot^k(\rho_0\tilde{J}_t),_k - 2\tilde{A}_\cdot^k(\rho_0\tilde{J}^{-1}),_k\Big)\Big|_{t=0}, \quad \text{for } a = 1,2,3,4. \tag{3.5}$$

Since $2\tilde{A}_i^k(\rho_0\tilde{J}^{-1}),_k = \rho_0^{-1}\tilde{a}_i^k(\rho_0^2\tilde{J}^{-2}),_k$, we have that $\mathrm{v}_a \to v_a$, $a = 1, 2$, as $\kappa\to 0$ where v_a are defined in Section 1.5. We use (3.1b) to compute the following identities: for $a = 0, 1, 2, 3$,

$$\partial_t^a[\rho_0^2\tilde{J}^{-2} - \kappa\rho_0^2\tilde{J}^{-3}\tilde{J}_t]\big|_{t=0} = \partial_t^a[\beta(t) - \sigma\tilde{g}^{\alpha\beta}\tilde{\eta},_{\alpha\beta}\cdot\tilde{n} - \kappa\tilde{g}^{\alpha\beta}\tilde{v},_{\alpha\beta}\cdot\tilde{n}]\big|_{t=0}. \tag{3.6}$$

4. A PRIORI ESTIMATES FOR THE κ-PROBLEM (3.1)

We establish our κ-independent a priori estimates in this section; the precise estimate is stated below in Lemma 4.1. Our existence of solutions to the κ-problem (3.1) is established in Section 5.

For $\kappa > 0$, we define the following higher-order energy function:

$$E^\kappa(t) = 1 + \sum_{a=0}^{5} \|\partial_t^a \tilde\eta(t)\|^2_{5-a} + |\tilde v_{ttt} \cdot \tilde n(t)|^2_1 + \sum_{a=0}^{2} |\bar\partial^2 \partial_t^a \tilde v \cdot \tilde n(t)|^2_{2.5-a}$$
$$+ \int_0^T |\sqrt{\kappa}\bar\partial \tilde v_{tttt} \cdot \tilde n|^2_0 + \int_0^T \|\sqrt{\kappa}\tilde v_{tttt}\|^2_1 + \sum_{a=0}^{3} \|\kappa \partial_t^a \tilde v(t)\|^2_{5-a} + \sum_{a=0}^{2} |\kappa \bar\partial^2 \partial_t^a \tilde v_t \cdot \tilde n(t)|^2_{2.5-a}. \tag{4.1}$$

Remark 11. The inequality stated in Theorem 5.1 ensures that $E^\kappa(t)$ is continuous in time.

We make the following definition to allow for constants to depend on $1/\delta$:

Definition 4.1 (Notational convention for constants depending on $1/\delta > 0$)**.** *We let $\mathcal{P}$ denote a generic polynomial with constant and coefficients depending on $1/\delta > 0$.*

We define the constant $\mathcal{N}_0 > 0$ by

$$\mathcal{N}_0 = \mathcal{P}(\|u_0\|_{100}, \|\rho_0\|_{100}). \tag{4.2}$$

We let $\mathcal{R}$ denote generic lower-order terms satisfying

$$\int_0^T \mathcal{R} \le \mathcal{N}_0 + \delta \sup_{t\in[0,T]} E^\kappa(t) + T\,\mathcal{P}(\sup_{t\in[0,T]} E^\kappa(t)).$$

The artificially high $H^{100}(\Omega)$-norm in defining $\mathcal{N}_0$ is acceptable as the initial data (ρ_0, u_0, Ω) is of C^∞-class. We shall assume that

$$\frac{1}{2} \le \tilde J \le \frac{3}{2} \quad \text{for all } t \in [0,T] \text{ and } x \in \Omega. \tag{4.3}$$

The bounds (4.3) are possible by taking $T > 0$ sufficiently small, since thanks to Theorem 5.1

$$\|\tilde J - 1\|_{L^\infty(\Omega)} \le C \|\int_0^t \partial_t \tilde J\|_2 \le C\sqrt{T}. \tag{4.4}$$

Lemma 4.1 (A priori estimates for the κ-problem)**.** *We let $\tilde v$ solve the κ-problem* (3.1) *on a time-interval* $[0,T]$*, for some $T = T_\kappa > 0$. Then independent of $\kappa > 0$,*

$$\sup_{t\in[0,T]} E^\kappa(t) \le \int_0^T \mathcal{R}. \tag{4.5}$$

We will establish Lemma 4.1 in the following six steps:

Step 1: The κ-independent curl-estimates. We follow [13] in establishing

Lemma 4.2 (The κ-independent curl-estimates)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{4} \|\operatorname{curl} \partial_t^a \tilde\eta(t)\|^2_{4-a} + \int_0^T \|\sqrt{\kappa}\operatorname{curl} \tilde v_{tttt}\|^2_0 + \sup_{t\in[0,T]} \sum_{a=0}^{3} \|\kappa \operatorname{curl} \partial_t^a \tilde v(t)\|^2_{4-a} \le \int_0^T \mathcal{R}.$$

Proof. The Lagrangian curl of (3.1a) yields $\operatorname{curl}_{\tilde\eta} \tilde v_t = 0$. Setting

$$B(\tilde A, D\tilde v) = \varepsilon_{\cdot ji} \tilde A_t{}^s_j \tilde v^i,_s \,,$$

we find that $(\operatorname{curl}_{\tilde\eta} \tilde v)_t = B(\tilde A, D\tilde v)$. By the fundamental theorem of calculus,

$$\operatorname{curl}_{\tilde\eta} \tilde v(t) = \operatorname{curl} u_0 + \int_0^t B(\tilde A, D\tilde v). \tag{4.6}$$

Applying the gradient operator D to (4.6) and a second application of the fundamental theorem of calculus, we find that

$$\begin{aligned} D \operatorname{curl} \tilde\eta(t) = {} & tD \operatorname{curl} u_0 - \varepsilon_{\cdot ji} D\tilde\eta^i,_s \int_0^t \tilde A_t{}^s_j \\ & + \varepsilon_{\cdot ji} \int_0^t [\tilde A_t{}^s_j D\tilde\eta^i,_s - D\tilde A^s_j \tilde v^i,_s] + \int_0^t \int_0^{t'} DB(\tilde A, D\tilde v), \end{aligned} \tag{4.7}$$

where we have used the identity $\operatorname{curl}_{\tilde\eta} D\tilde\eta = \operatorname{curl} D\tilde\eta + \varepsilon_{\cdot ji} D\tilde\eta^i,_s \int_0^t \tilde A_t{}^s_j$.

The differentiation formula (2.7) equally holds for when the gradient D replaces ∂_t; hence, the first three terms on the right-hand side of (4.7) are, with respect to the action of D^3, each bounded by $\int_0^T \mathcal{R}$.

We next analyze the highest-order term created in $\int_0^t \int_0^{t'} D^4 B(\tilde A, D\tilde v)$. With (2.7),

$$DB(\tilde A, D\tilde v) = -\varepsilon_{\cdot ji} [\tilde A^q_j D\tilde v^r,_q \tilde A^s_r \tilde v^i,_s + \tilde A^q_j \tilde v^r,_q \tilde A^s_r D\tilde v^i,_s + \tilde v^r,_q \tilde v^i,_s D(\tilde A^q_j \tilde A^s_r)],$$

from which it follows that the highest-order term of $D^4 B(\tilde A, D\tilde v)$ is

$$-\varepsilon_{\cdot ji} \int_0^t \int_0^{t'} [\tilde A^q_j D^4 \tilde v^r,_q \tilde A^s_r \tilde v^i,_s + \tilde A^q_j \tilde v^r,_q \tilde A^s_r D^4 \tilde v^i,_s].$$

With a relaxation of the precise structure of the summands in the integrands of the highest-order terms of $D^4 B(\tilde A, D\tilde v)$, we highlight the derivative count that results from integration by parts in time by writing

$$\int_0^t \int_0^{t'} D^4 B(\tilde A, D\tilde v) = - \int_0^t \int_0^{t'} D^5 \tilde\eta (D\tilde v \, \tilde A \, \tilde A)_t + \int_0^t D^5 \tilde\eta D\tilde v \, \tilde A \, \tilde A.$$

With such a temporal-integration-by-parts computation, the action of D^3 in (4.7) yields

$$\sup_{t\in[0,T]} \| \operatorname{curl} \tilde\eta(t) \|_4^2 \le \int_0^T \mathcal{R}, \tag{4.8}$$

and by the same arguments, the action of D^3 and κD^4 in (4.6) yield

$$\sup_{t\in[0,T]} \| \operatorname{curl} \tilde v(t) \|_3^2 + \sup_{t\in[0,T]} \| \kappa \operatorname{curl} \tilde v(t) \|_4^2 \le \int_0^T \mathcal{R}. \tag{4.9}$$

By returning to the Lagrangian vorticity equation $\operatorname{curl}_{\tilde\eta} \tilde v_t = 0$, we find that

$$\operatorname{curl} \tilde v_t = \varepsilon_{j \cdot i} \tilde v^i_t,_s \int_0^t \tilde A_t{}^s_j, \tag{4.10}$$

and by considering the action of D^2 and κD^3 in the identity (4.10) we infer that

$$\sup_{t\in[0,T]} \| \operatorname{curl} \tilde v_t(t) \|_2^2 + \sup_{t\in[0,T]} \| \kappa \operatorname{curl} \tilde v_t(t) \|_3^2 \le \int_0^T \mathcal{R}. \tag{4.11}$$

By considering the action of $D\partial_t$, ∂_t^2, $\kappa D^2\partial_t$, $\kappa D\partial_t^2$, and $\sqrt{\kappa}\partial_t^3$ in (4.10), and using the fundamental-theorem-of-calculus identity $\partial_t^a \tilde{v}_t = \partial_t^a \tilde{v}_t|_{t=0} + \int_0^t \partial_t^{a+1}\tilde{v}_t$ in the lower-order terms, we establish that

$$\sup_{t\in[0,T]} \sum_{a=0}^{1} \| \operatorname{curl} \partial_t^a \tilde{v}_{tt}(t)\|^2_{1-a} + \sum_{a=0}^{1} \|\kappa \operatorname{curl} \partial_t^a \tilde{v}_{tt}(t)\|^2_{2-a} + \int_0^T \|\kappa \operatorname{curl} \tilde{v}_{tttt}(t)\|_0^2 \le \int_0^T \mathcal{R}. \tag{4.12}$$

The sum of the inequalities (4.8)–(4.12) completes the proof. $\square$

Step 2: The κ-independent estimates for $\tilde{v}_{tttt}$ and $\tilde{v}_{ttt}$. We equivalently write the momentum equations (3.1a) and boundary condition (3.1b) of the κ-problem as

$$\rho_0 \tilde{v}_t^i + \tilde{a}_i^k(\rho_0^2 \tilde{J}^{-2}),_k - \kappa \rho_0 \tilde{J}^{-1} \tilde{a}_i^k (\rho_0 \tilde{J}_t),_k = 0 \qquad \text{in } \Omega \times (0,T], \tag{4.13a}$$

$$\rho_0^2 \tilde{J}^{-2} - \kappa \rho_0^2 \tilde{J}^{-1} \tilde{J}_t = \tilde{H}_\kappa \qquad \text{on } \Gamma \times (0,T], \tag{4.13b}$$

where $\tilde{H}_\kappa$ is given by (3.2). From (3.1a), we have the identity

$$\tilde{v}_t \cdot \tilde{\eta}_\gamma = \bar{\partial}_\gamma[-2\rho_0 \tilde{J}^{-1} + \kappa \rho_0 \tilde{J}_t].$$

Multiplying this identity by $\rho_0 \tilde{J}^{-1}$ yields

$$\rho_0 \tilde{J}^{-1} \tilde{v}_t \cdot \tilde{\eta}_\gamma = \bar{\partial}_\gamma[-\rho_0^2 \tilde{J}^{-2} + \kappa \rho_0^2 \tilde{J}^{-1} \tilde{J}_t] - \kappa(\rho_0 \tilde{J}^{-1}),_\gamma \rho_0 \tilde{J}_t.$$

We use the boundary condition (4.13b) to obtain the following tangential identity for v_t:

$$\rho_0 \tilde{J}^{-1} \tilde{v}_t \cdot \tilde{\eta},_\gamma = \bar{\partial}_\gamma \Big[\sigma \tilde{g}^{\mu\nu} \tilde{\eta},_{\mu\nu} \cdot \tilde{n} + \kappa \tilde{g}^{\mu\nu} \tilde{v},_{\mu\nu} \cdot \tilde{n} - \beta(t)\Big] - \kappa(\rho_0 \tilde{J}^{-1}),_\gamma \rho_0 \tilde{J}_t. \tag{4.14}$$

Proposition 4.1 (Energy estimates for the fourth time-differentiated problem)**.**

$$\sup_{t\in[0,T]} \|\tilde{v}_{tttt}(t)\|_0^2 + \sup_{t\in[0,T]} \|\partial_t^4 \tilde{J}(t)\|_0^2 + \sup_{t\in[0,T]} |\tilde{v}_{ttt} \cdot \tilde{n}(t)|_1^2$$
$$+ \int_0^T |\sqrt{\kappa}\partial_t^4 \tilde{v} \cdot \tilde{n}(t)|_1^2 + \int_0^T \|\sqrt{\kappa}\partial_t^5 \tilde{J}\|_0^2 \le \int_0^T \mathcal{R}.$$

Proof. Testing four time-derivatives of (4.13a) against $\partial_t^4 \tilde{v}$ in the $L^2(\Omega)$-inner product, and integrating by parts with respect to ∂_k in the interior integrals $\int_\Omega \tilde{a}_i^k \partial_t^4(\rho_0^2 \tilde{J}^{-2}),_k \partial_t^4 \tilde{v}^i$ and $-\kappa \int_\Omega \tilde{a}_i^k \partial_t^4[\rho_0 \tilde{J}^{-1}(\rho_0 \tilde{J}_t),_k] \partial_t^4 \tilde{v}^i$, we find that

$$\int_\Omega \partial_t^4[\rho_0 \tilde{v}_t^i] \partial_t^4 \tilde{v}^i - \underbrace{\int_\Omega \partial_t^4[\rho_0^2 \tilde{J}^{-2}] \tilde{a}_i^k \partial_t^4 \tilde{v}^i,_k}_{\mathcal{I}_1} + \kappa \underbrace{\int_\Omega \partial_t^4[\rho_0^2 \tilde{J}^{-1} \tilde{J}_t] \tilde{a}_i^k \partial_t^4 \tilde{v}^i,_k}_{\mathcal{I}_2}$$
$$+ \underbrace{\int_\Gamma \partial_t^4[\rho_0^2 \tilde{J}^{-2} - \kappa \rho_0^2 \tilde{J}^{-1} \tilde{J}_t] \tilde{a}_i^k \partial_t^4 \tilde{v}^i N^k}_{\mathfrak{i}} = \mathcal{R}.$$

It is convenient to rewrite this equation as

$$\frac{1}{2}\frac{d}{dt}\int_\Omega \rho_0 |\partial_t^4 \tilde{v}|^2 + \frac{d}{dt}\int_\Omega \rho_0^2 \tilde{J}^{-3} |\partial_t^4 \tilde{J}|^2 + \kappa \int_\Omega \rho_0^2 \tilde{J}^{-1} |\partial_t^5 \tilde{J}|^2 + \mathfrak{i} = \mathcal{R}, \tag{4.15}$$

where the identity $\tilde{J}_t = \tilde{a}_r^s \tilde{v}^r,_s$ implies that the error created in order to write $\tilde{a}_i^k \partial_t^4 \tilde{v}^i,_k$ as $\partial_t^5 \tilde{J}$ in $\mathcal{I}_1$ and $\mathcal{I}_2$ is of lower-order and so is absorbed in $\mathcal{R}$.

Rewriting the boundary integral in (4.15). Using the outward normal identity (2.9) and the boundary condition (4.13b) with $\sigma = 1$, we find that

$$\begin{aligned}
\mathfrak{i} = {} & \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\, \tilde v_{ttt},_\alpha \cdot \tilde n \, \tilde v_{ttt},_\beta \cdot \tilde n + \int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\sqrt{\kappa}\tilde v_{tttt},_\alpha \cdot \tilde n \sqrt{\kappa}\tilde v_{tttt},_\beta \cdot \tilde n \\
& \underbrace{- \int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{ttt},_\alpha \cdot \partial_t \tilde n \, \tilde v_{ttt},_\beta \cdot \tilde n}_{\mathfrak{r}_1} - \frac{1}{2}\underbrace{\int_\Gamma \partial_t[\sqrt{\tilde g}\tilde g^{\alpha\beta}]\, \tilde v_{ttt},_\alpha \cdot \tilde n \, \tilde v_{ttt},_\beta \cdot \tilde n}_{\mathcal{R}} \\
& \underbrace{+ \int_\Gamma \tilde v_{ttt},_\beta \cdot \tilde n \sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{tttt} \cdot \tilde n,_\alpha + \int_\Gamma \tilde v_{ttt},_\beta \cdot \tilde n,_\alpha \sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{tttt} \cdot \tilde n}_{\mathfrak{r}_2} \\
& \underbrace{+ \int_\Gamma \tilde v_{ttt},_\beta \cdot \tilde n (\sqrt{\tilde g}\tilde g^{\alpha\beta}),_\alpha \tilde v_{tttt} \cdot \tilde n - \sum_{l=0}^{3} c_l \int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\partial_t^l \tilde\eta^j,_{\alpha\beta}\, \partial_t^{4-l}\tilde n^j \, \tilde v_{tttt} \cdot \tilde n}_{\mathfrak{r}_3} \\
& \underbrace{- \sum_{l=1}^{4} c_l \int_\Gamma \partial_t^l \tilde g^{\alpha\beta}\partial_t^{4-l}[\tilde\eta,_{\alpha\beta} \cdot \tilde n]\sqrt{\tilde g}\tilde v_{tttt} \cdot \tilde n}_{\mathfrak{r}_4} \\
& \underbrace{+ \kappa \int_\Gamma \tilde v^i_{tttt},_\beta \left(\tilde n^i \tilde g^{\alpha\beta}\tilde n^j\right),_\alpha \sqrt{\tilde g}\tilde v^j_{tttt} - \kappa \sum_{l=1}^{4} c_l \int_\Gamma \partial_t^l(\tilde g^{\alpha\beta}\tilde n^i)\partial_t^{4-l}\tilde\eta,_{\alpha\beta} \sqrt{\tilde g}\tilde v_{tttt} \cdot \tilde n}_{\mathfrak{r}_5} \,.
\end{aligned} \tag{4.16}$$

In our analysis of $\int_0^T \mathfrak{r}_i$, $i = 1, 2, 3$, we adopt the convention of letting

$\tilde\ell$ denote a function of $L^\infty(\Gamma)$-class and $\tilde l$ a function of $H^{0.5}(\Gamma)$-class.

We recycle the symbols $\mathfrak{j}$, $\mathfrak{j}_A$, $\mathfrak{j}_B$, etc. in terming boundary integrals that require explanation.

Analysis of $\int_0^T \mathfrak{r}_1$ in the time-integral of (4.16)**.** The action of $\bar\partial_\alpha \partial_t^2$ in the tangential identity (4.14) provides that

$$\tilde v_{ttt},_\alpha \cdot \tilde\eta,_\gamma = \rho_0^{-1}\tilde J[\bar\partial_{\alpha\gamma}\left(\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu} \cdot \tilde n + \kappa \tilde g^{\mu\nu}\tilde v,_{\mu\nu} \cdot \tilde n - \beta(t)\right)_{tt} - \tilde l_{\gamma\alpha}], \tag{4.17}$$

where the lower-order $\tilde l_{\gamma\alpha} \in H^{0.5}(\Gamma)$ is given by

$$\tilde l_{\gamma\alpha} = \kappa\bar\partial_\alpha\left[(\rho_0 \tilde J^{-1}),_\gamma \rho_0 \tilde J_t\right]_{tt} + \tilde v_{ttt} \cdot (\tilde\eta,_\gamma \rho_0 \tilde J^{-1}),_\alpha + \sum_{a=1}^{2} c_a [\partial_t^a \tilde v \cdot \partial_t^{3-a}(\tilde\eta,_\gamma \rho_0 \tilde J^{-1})],_\alpha \,.$$

Setting $\tilde\ell^{\alpha\beta}_\gamma = \rho_0^{-1}\tilde J\sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde g^{\gamma\delta}\tilde v,_\delta \cdot \tilde n$ we use the tangential identity (4.17), together with the outward normal differentiation formula (2.11c), to find that

$$\mathfrak{r}_1 = \underbrace{\int_\Gamma \tilde\ell^{\alpha\beta}_\gamma \bar\partial_{\gamma\alpha}[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu} \cdot \tilde n]_{tt}\, \tilde v_{ttt},_\beta \cdot \tilde n}_{\mathfrak{r}_1{}'} + \underbrace{\kappa \int_\Gamma \tilde\ell^{\alpha\beta}_\gamma \bar\partial_{\gamma\alpha}[\tilde g^{\mu\nu}\tilde v,_{\mu\nu} \cdot \tilde n]_{tt}\, \tilde v_{ttt},_\beta \cdot \tilde n}_{\mathfrak{j}} + \mathcal{R}. \tag{4.18}$$

Concerning $\mathfrak{r}_1{}'$, we integrate by parts with respect to a time-derivative to write

$$\int_0^T \mathfrak{r}_1{}' = -\underbrace{\int_\Gamma \bar\partial_\gamma[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n]_{tt}\,\bar\partial_\alpha[\tilde\ell_\gamma^{\alpha\beta}\tilde v_{tt},_\beta\cdot\tilde n]}_{\mathfrak{j}_A}\Big|_0^T + \int_0^T \underbrace{\int_\Gamma \bar\partial_\gamma[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n]_{tt}\,\bar\partial_\alpha[\tilde\ell_\gamma^{\alpha\beta}\tilde v_{tt},_\beta\cdot\tilde n_t]}_{\mathfrak{j}_B}$$
$$+\int_0^T \underbrace{\int_\Gamma \bar\partial_\gamma[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n]_{tt}\,\bar\partial_\alpha[\partial_t\tilde\ell_\gamma^{\alpha\beta}\tilde v_{tt},_\beta\cdot\tilde n]}_{\mathfrak{j}_C} + \int_0^T \underbrace{\int_\Gamma \bar\partial_\gamma[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n]_{ttt}\,\bar\partial_\alpha[\tilde\ell_\gamma^{\alpha\beta}\tilde v_{tt},_\beta\cdot\tilde n]}_{\mathfrak{j}_D},$$

where we have further integrated by parts with respect to $\bar\partial_\alpha$ on the right-hand side.

Interpolation and two applications of Young's inequality provide that

$$|\tilde v_{tt},_\beta\cdot\tilde n(T)|_1^2 \le C|\tilde v_{tt}\cdot\tilde n(T)|_{1.5}|\bar\partial\tilde v_{tt}\cdot\tilde n(T)|_{1.5} \le C|\tilde v_{tt}\cdot\tilde n(T)|_{0.5}^2 + 2\delta|\bar\partial\tilde v_{tt}\cdot\tilde n(T)|_{1.5}^2.$$

Hence,

$$\mathfrak{j}_A(t)\big|_0^T \le \int_0^T \mathcal{R}.$$

We record that via Lemma 2.3,

$$|\tilde v_{tt},_{\alpha\beta}\cdot\tilde\eta,_\sigma|^2_{H^{-0.5}(\Gamma)} \le C\left[\|\bar\partial^2\tilde v_{tt}\|_0^2 + \|\operatorname{curl}\bar\partial\tilde v_{tt}\|_0^2\right] \le C\|\tilde v_{tt}\|_2^2.$$

Since $\tilde n_t = -\tilde\eta,_\sigma\,\tilde g^{\sigma\rho}\tilde v,_\rho\cdot\tilde n$ and $\bar\partial^2\tilde v_t\cdot\tilde n$ is in $H^{1.5}(\Gamma)$, we use an $H^{-0.5}(\Gamma)$-duality pairing in the highest-order term to write

$$\int_0^T \mathfrak{j}_B \le \int_0^T \mathcal{R} + C\int_0^T |\bar\partial^2\tilde v_t\cdot\tilde n(t)|_{1.5}|\tilde v_{tt},_{\alpha\beta}\cdot\tilde\eta,_\sigma|_{H^{-0.5}(\Gamma)} \le \int_0^T \mathcal{R}.$$

We use the Cauchy-Schwarz inequality for the estimate

$$\int_0^T \mathfrak{j}_C \le \int_0^T \mathcal{R}.$$

Regarding $\int_0^T \mathfrak{j}_D$, for the integral $\int_0^T\int_\Gamma \bar\partial_\gamma[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n]_{ttt}\,\bar\partial_\alpha\tilde\ell_\gamma^{\alpha\beta}\tilde v_{tt},_\beta\cdot\tilde n$, we integrate by parts with respect to $\bar\partial_\gamma$ and estimate using the Cauchy-Schwarz inequality, and for the integral $\int_0^T\int_\Gamma \bar\partial_\gamma[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n]_{ttt}\,\tilde\ell_\gamma^{\alpha\beta}\tilde v_{tt},_\beta\cdot\tilde n,_\alpha$, we also integrate by parts with respect to $\bar\partial_\gamma$ and then estimate using an $H^{-0.5}(\Gamma)$-duality pairing. These methods work equally well in all terms of the spacetime-integral $\int_0^T\int_\Gamma \bar\partial_\gamma[\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n]_{ttt}\,\tilde\ell_\gamma^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n$, as well as in $\int_0^T \mathfrak{j}$, where $\mathfrak{j}$ is defined in (4.18). Thus, we have established that

$$\int_0^T \mathfrak{r}_1 \le \int_0^T \mathcal{R}. \tag{4.19}$$

Analysis of $\int_0^T \mathfrak{r}_2$ in the time-integral of (4.16)**.** We have that

$$\mathfrak{r}_2 = \underbrace{\int_\Gamma \tilde v_{ttt},_\beta\cdot\tilde n\sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{tttt}\cdot\tilde n,_\alpha}_{\mathfrak{r}_{2a}} + \underbrace{\int_\Gamma \tilde v_{ttt},_\beta\cdot\tilde n,_\alpha\sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{tttt}\cdot\tilde n}_{\mathfrak{r}_{2b}}. \tag{4.20}$$

We write the action of ∂_t^3 in the tangential identity (4.14) as

$$\tilde v_{tttt}\cdot\tilde\eta,_\gamma = \rho_0^{-1}\tilde J\left[\bar\partial_\gamma\left(\tilde g^{\mu\nu}\tilde\eta,_{\mu\nu}\cdot\tilde n + \kappa\tilde g^{\mu\nu}\tilde v,_{\mu\nu}\cdot\tilde n - \beta(t)\right)_{ttt} - \tilde l_\gamma\right], \tag{4.21a}$$

where the lower-order $\tilde{l}_\gamma \in H^{0.5}(\Gamma)$ is given by

$$\tilde{l}_\gamma = \kappa\big[(\rho_0 \tilde{J}^{-1}),_\gamma \, \rho_0 \tilde{J}_t\big]_{ttt} + \sum_{l=1}^{3} c_a \partial_t^a \tilde{v} \cdot \partial_t^{4-l} (\tilde{\eta},_\gamma \, \rho_0 \tilde{J}^{-1}). \tag{4.21b}$$

Letting $\tilde{\ell}^\beta_\gamma = -\rho_0^{-1} \tilde{J} \sqrt{\tilde{g}} \tilde{g}^{\alpha\beta} \tilde{g}^{\gamma\delta} \tilde{\eta},_{\delta\alpha} \cdot \tilde{n}$, we use (4.21) to write

$$\mathfrak{r}_{2a} = \underbrace{-\int_\Gamma \tilde{\ell}^\beta_\gamma \, [\tilde{g}^{\mu\nu} \tilde{\eta},_{\mu\nu} \cdot \tilde{n}]_{ttt} \tilde{v}_{ttt},_{\beta\gamma} \cdot \tilde{n}}_{\mathfrak{r}_{2a}'} + \kappa \underbrace{\int_\Gamma \tilde{\ell}^\beta_\gamma \, \bar{\partial}_\gamma [\tilde{g}^{\mu\nu} \tilde{v}_{ttt},_{\mu\nu} \cdot \tilde{n}] \, \tilde{v}_{ttt},_\beta \cdot \tilde{n}}_{\mathfrak{r}_{2a}''} + \mathcal{R},$$

where we have used integration by parts with respect to $\bar{\partial}_\gamma$ to determine $\mathfrak{r}_{2a}'$. We integrate by parts with respect to time in order to write

$$\begin{aligned} -\int_0^T \mathfrak{r}_{2a}' = & -\int_\Gamma \tilde{\ell}^\beta_\gamma \, \tilde{v}_{tt},_{\beta\gamma} \cdot \tilde{n} \, [\tilde{g}^{\mu\nu} \tilde{\eta},_{\mu\nu} \cdot \tilde{n}]_{ttt} \Big|_0^T + \int_0^T \int_\Gamma \tilde{\ell}^\beta_\gamma \, \tilde{v}_{tt},_{\beta\gamma} \cdot \tilde{n}_t \, [\tilde{g}^{\mu\nu} \tilde{\eta},_{\mu\nu} \cdot \tilde{n}]_{ttt} \\ & + \int_0^T \int_\Gamma \partial_t \tilde{\ell}^\beta_\gamma \, \tilde{v}_{tt},_{\beta\gamma} \cdot \tilde{n} \, [\tilde{g}^{\mu\nu} \tilde{\eta},_{\mu\nu} \cdot \tilde{n}]_{ttt} + \int_0^T \int_\Gamma \tilde{\ell}^\beta_\gamma \, \tilde{v}_{tt},_{\beta\gamma} \cdot \tilde{n} \, [\tilde{g}^{\mu\nu} \tilde{\eta},_{\mu\nu} \cdot \tilde{n}]_{tttt} \\ = & \int_0^T \underbrace{\int_\Gamma \tilde{\ell}^\beta_\gamma \, \tilde{v}_{tt},_{\beta\gamma} \cdot \tilde{n} \, \tilde{g}^{\mu\nu} \tilde{v}_{ttt},_{\mu\nu} \cdot \tilde{n}}_{\mathfrak{j}} + \int_0^T \mathcal{R}. \end{aligned}$$

Letting $\tilde{\ell} = \rho_0^{-1} \tilde{J} \sqrt{\tilde{g}} \, \tilde{g}^{\gamma\delta} \tilde{\eta},_{\gamma\delta} \cdot \tilde{n}$ we utilize the symmetry of $\tilde{\ell}^\beta_\gamma$ to exchange $\bar{\partial}_\alpha$ and $\bar{\partial}_\gamma$ via integration by parts for

$$\begin{aligned} \int_0^T \mathfrak{j} = & \frac{1}{2} \int_\Gamma \tilde{\ell} \, |\tilde{g}^{\mu\nu} \tilde{v}_{tt},_{\mu\nu} \cdot \tilde{n}|^2 \Big|_0^T \\ & - \int_0^T \int_\Gamma \tilde{\ell} \, \tilde{v}_{tt},_{\alpha\beta} \cdot \tilde{n} \, \tilde{v}_{tt},_{\mu\nu} \cdot (n \tilde{g}^{\mu\nu})_t - \frac{1}{2} \int_0^T \int_\Gamma \tilde{\ell}_t \, |\tilde{g}^{\mu\nu} \tilde{v}_{tt},_{\mu\nu} \cdot \tilde{n}|^2 = \int_0^T \mathcal{R}. \end{aligned} \tag{4.22}$$

Next, integrating by parts in time, we find that

$$\begin{aligned} \kappa \int_0^T \mathfrak{r}_{2a}'' = & \int_\Gamma \tilde{\ell}^\beta_\gamma \, \bar{\partial}_\gamma [\tilde{g}^{\mu\nu} \tilde{v}_{tt},_{\mu\nu} \cdot \tilde{n}] \, \kappa(\tilde{v}_{ttt},_\beta \cdot \tilde{n}) \Big|_0^T \\ & + \int_0^T \int_\Gamma [\tilde{g}^{\mu\nu} \tilde{v}_{tt},_{\mu\nu} \cdot \partial_t \tilde{n}] \bar{\partial}_\gamma [\tilde{\ell}^\beta_\gamma \; \kappa(\tilde{v}_{ttt},_\beta \cdot \tilde{n})] - \int_0^T \int_\Gamma \tilde{\ell}^\beta_\gamma \, \bar{\partial}_\gamma [\tilde{g}^{\mu\nu} \kappa(\tilde{v}_{tt},_{\mu\nu} \cdot \tilde{n})] \, \tilde{v}_{ttt},_\beta \cdot \partial_t \tilde{n} \\ & - \int_0^T \underbrace{\int_\Gamma \tilde{\ell}^\beta_\gamma \, \bar{\partial}_\gamma [\tilde{g}^{\mu\nu} \sqrt{\kappa} \tilde{v}_{tt},_{\mu\nu} \cdot \tilde{n}] \, \sqrt{\kappa} \tilde{v}_{tttt},_\beta \cdot \tilde{n}}_{\mathfrak{j}} = \int_0^T \mathcal{R} + \int_0^T \mathfrak{j}. \end{aligned}$$

Interpolation and Young's inequality provide for the estimate

$$\begin{aligned} \int_0^T \mathfrak{j} & \le \delta \int_0^T |\sqrt{\kappa} \bar{\partial} \tilde{v}_{tttt} \cdot \tilde{n}|_0^2 + C_\delta \int_0^T |\sqrt{\kappa} \bar{\partial} \tilde{v}_{tt} \cdot \tilde{n}|_2^2 \\ & \le \delta \int_0^T |\sqrt{\kappa} \bar{\partial} \tilde{v}_{tttt} \cdot \tilde{n}|_0^2 + C_\delta \, T \, |\bar{\partial} \tilde{v}_{tt} \cdot \tilde{n}|_{1.5} |\kappa \bar{\partial} \tilde{v}_{tt} \cdot \tilde{n}|_{2.5} \le \int_0^T \mathcal{R}. \end{aligned}$$

We have thus established that

$$\int_0^T \mathfrak{r}_{2a} = \int_0^T \mathcal{R}.$$

The analysis of $\int_0^T \mathfrak{r}_{2b}$ is similar. We set $\tilde{\ell}^{\gamma\beta} = \sqrt{\tilde{g}}\tilde{g}^{\alpha\beta}\tilde{g}^{\gamma\delta}\tilde{\eta},_{\delta\alpha} \cdot\tilde{n}$ and write

$$\int_0^T \mathfrak{r}_{2b} = \int_\Gamma \tilde{v}_{ttt},_\beta \cdot\tilde{n},_\alpha \sqrt{\tilde{g}}\tilde{g}^{\alpha\beta}\,\tilde{v}_{ttt}\cdot\tilde{n}\Big|_0^T - \int_0^T\int_\Gamma \tilde{v}^j_{ttt},_\beta\,[\tilde{n}^j,_\alpha\sqrt{\tilde{g}}\tilde{g}^{\alpha\beta}\tilde{n}^i]_t\,\tilde{v}^i_{ttt}$$
$$+\int_0^T \underbrace{\int_\Gamma \tilde{\ell}^{\gamma\beta}\,\tilde{v}_{tttt},_\beta \cdot\tilde{\eta},_\gamma\;\tilde{v}_{ttt}\cdot\tilde{n}}_{\mathfrak{r}_{2b}{}'} = \int_0^T \mathcal{R} + \int_0^T \mathfrak{r}_{2b}{}'.$$

Regarding $\int_0^T \mathfrak{r}_{2b}{}'$, we use the identity

$$\tilde{v}_{tttt},_\beta \cdot\tilde{\eta},_\gamma = \rho_0^{-1}\tilde{J}[\bar{\partial}_\beta\bar{\partial}_\gamma\big(\tilde{g}^{\mu\nu}\tilde{\eta},_{\mu\nu}\cdot\tilde{n} + \kappa\tilde{g}^{\mu\nu}\tilde{v},_{\mu\nu}\cdot\tilde{n} - \beta(t)\big)_{ttt} - \tilde{l}'_{\beta\gamma}], \tag{4.23}$$

where $\tilde{l}'_{\beta\gamma} = \tilde{v}_{tttt}\cdot\tilde{\eta},_{\gamma\beta}\,\rho_0\tilde{J}^{-1} + \tilde{v}_{tttt}\cdot\tilde{\eta},_\gamma\,(\rho_0\tilde{J}^{-1}),_\beta + \bar{\partial}_\beta\tilde{l}_\gamma$, with $\tilde{l}_\gamma$ given by (4.21b). We integrate by parts with respect to both $\bar{\partial}_\beta$ and $\bar{\partial}_\gamma$ in $\int_0^T\int_\Gamma \rho_0^{-1}\tilde{J}\bar{\partial}_{\beta\gamma}[\tilde{g}^{\mu\nu}\tilde{\eta},_{\mu\nu}\cdot\tilde{n}]_{ttt}\,\tilde{\ell}^{\gamma\beta}\,\tilde{v}_{ttt}\cdot\tilde{n}$, where the highest-order term produced by $\bar{\partial}_{\beta\gamma}$-integration by parts is (4.22). To estimate the integral where $\bar{\partial}_\beta\tilde{l}_\gamma$ appears, we have the choice of integration by parts with respect to $\bar{\partial}_\beta$ or an $H^{-0.5}(\Gamma)$-duality pairing. In the integral where $\tilde{v}_{tttt}\cdot\tilde{\eta},_\gamma\,(\rho_0\tilde{J}^{-1}),_\beta$ appears, we use the identity (4.21a). Hence, we conclude that

$$\int_0^T \mathfrak{r}_{2b}{}' = \int_0^T \mathcal{R} - \int_0^T \underbrace{\int_\Gamma \tilde{\ell}^{\gamma\beta}\,\bar{\partial}_\beta\tilde{l}_\gamma\,\tilde{v}_{ttt}\cdot\tilde{n}}_{\mathcal{R}}\,.$$

Thus, we have established that

$$\int_0^T \mathfrak{r}_2 \le \int_0^T \mathcal{R}. \tag{4.24}$$

Analysis of $\int_0^T \mathfrak{r}_3$ in the time-integral of (4.16)**.** We have that

$$\mathfrak{r}_3 = \underbrace{\int_\Gamma \tilde{v}_{ttt},_\beta\cdot\tilde{n}(\sqrt{\tilde{g}}\tilde{g}^{\alpha\beta}),_\alpha\,\tilde{v}_{tttt}\cdot\tilde{n}}_{\mathfrak{r}_{3a}} - \sum_{l=0}^{3} c_l \underbrace{\int_\Gamma \sqrt{\tilde{g}}\tilde{g}^{\alpha\beta}\partial_t^l\tilde{\eta}^j,_{\alpha\beta}\,\partial_t^{4-l}\tilde{n}^j\,\tilde{v}_{tttt}\cdot\tilde{n}}_{\mathfrak{r}_{3b,l}}\,.$$

We will first establish that $\int_0^T \mathfrak{r}_{3a}$ cancels with a term arising in $\int_0^T \mathfrak{r}_{3b,0}$. Recalling that $\tilde{n}_t = -\tilde{g}^{\gamma\delta}[\tilde{v},_\delta\cdot\tilde{n}]\tilde{\eta},_\gamma$, the highest-order term of $\tilde{n}_{tttt}$ is $-\tilde{g}^{\gamma\delta}[\tilde{v}_{ttt},_\delta\cdot\tilde{n}]\tilde{\eta},_\gamma$. It follows that

$$-\int_0^T \mathfrak{r}_{3b,0} = \int_0^T \underbrace{\int_\Gamma \sqrt{\tilde{g}}\,\tilde{g}^{\alpha\beta}\tilde{g}^{\gamma\delta}\tilde{\eta},_{\alpha\beta}\cdot\tilde{\eta},_\gamma\,\tilde{v}_{ttt},_\delta\cdot\tilde{n}\,\tilde{v}_{tttt}\cdot\tilde{n}}_{\mathfrak{j}} + \int_0^T \mathcal{R}, \tag{4.25}$$

where the estimation of the lower-order terms of $\int_0^T \mathfrak{r}_{3b,0}$ requires integration by parts with respect to a time-derivative of $\tilde{v}_{tttt}$. Using the differentiation formulas (2.10a) and (2.10b), we write

$$\mathfrak{j} = -\underbrace{\int_\Gamma \tilde{v}_{ttt},_\beta\cdot\tilde{n}(\sqrt{\tilde{g}}\tilde{g}^{\alpha\beta}),_\alpha\,\tilde{v}_{tttt}\cdot\tilde{n}}_{\mathfrak{r}_{3a}}\,.$$

Thus,

$$\int_0^T \mathfrak{r}_{3a} - \int_0^T \mathfrak{r}_{3b,0} = \int_0^T \mathcal{R}.$$

The terms $\mathfrak{r}_{3b,1}$ and $\mathfrak{r}_{3b,2}$ are analyzed by integrating by parts with respect to a time-derivative of $\tilde v_{tttt}$, followed by elementary estimates. Thus,

$$\int_0^T \mathfrak{r}_{3b,1} + \int_0^T \mathfrak{r}_{3b,2} = \int_0^T \mathcal{R}.$$

We next examine $\mathfrak{r}_{3b,3} = \int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n_t\,\tilde v_{tttt}\cdot\tilde n$. After integration by parts with respect to a time-derivative of $\tilde v_{tttt}$, we find that

$$\mathfrak{r}_{3b,3} = -\int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{ttt},_{\alpha\beta}\cdot\tilde n_t\,\tilde v_{ttt}\cdot\tilde n + \mathcal{R} = \underbrace{\int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde v_{ttt},_\beta\cdot\tilde n_t\,\tilde v_{ttt},_\alpha\cdot\tilde n}_{\mathfrak{r}_{3b,3}{}'} + \mathcal{R}.$$

Letting $\tilde\ell^\gamma_{\alpha\beta} = \sqrt{\tilde g}\tilde g^{\alpha\beta}\tilde g^{\gamma\delta}\tilde v,_\delta\cdot\tilde n$, we have that

$$\begin{aligned}\int_0^T \mathfrak{r}_{3b,3}{}' = \int_\Gamma \tilde\ell^\gamma_{\alpha\beta}\,\tilde v_{ttt},_\beta\cdot\tilde\eta,_\gamma\,\tilde v_{tt},_\alpha\cdot\tilde n\Big|_0^T &- \int_0^T\int_\Gamma \tilde v_{ttt},^j_\beta\,\tilde v^i_{tt},_\alpha\cdot(\tilde n^i\,\tilde\eta^j,_\beta\,\tilde\ell^\gamma_{\alpha\beta})_t \\ &- \int_0^T \underbrace{\int_\Gamma \tilde\ell^\gamma_{\alpha\beta}\,\tilde v_{tttt},_\beta\cdot\tilde\eta,_\gamma\,\tilde v_{tt},_\alpha\cdot\tilde n}_{\mathfrak{j}} = -\int_0^T \mathfrak{j} + \int_0^T \mathcal{R}.\end{aligned}$$

By using the identity (4.23) in $\int_0^T \mathfrak{j}$, we note that the term corresponding with $\tilde l_\gamma$ has an elementary estimate after integration by parts with respect to $\bar\partial_\beta$. The remaining terms are similarly analyzed, except for $\int_0^T\int_\Gamma \tilde\ell^\gamma_{\alpha\beta}\rho_0^{-1}\tilde J\bar\partial_{\beta\gamma}[\tilde g^{\mu\nu}\tilde v_{tt},_{\mu\nu}\cdot\tilde n]\,\tilde v_{tt},_\alpha\cdot\tilde n$ in which we integrate by parts with respect to both $\bar\partial_\beta$ and $\bar\partial_\gamma$. For a certain highest-order term created by $\bar\partial_{\beta\gamma}$-integration by parts, we form an exact derivative: setting $\tilde\ell_\gamma = \rho_0^{-1}\tilde J\sqrt{\tilde g}\tilde g^{\gamma\delta}\tilde v,_\delta\cdot\tilde n$,

$$\int_0^T\int_\Gamma \tilde\ell_\gamma\,[\tilde g^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n],_\gamma\,\tilde g^{\mu\nu}\tilde v_{tt},_{\mu\nu}\cdot\tilde n = -\frac{1}{2}\int_0^T\int_\Gamma \bar\partial_\gamma\tilde\ell_\gamma|\tilde g^{\mu\nu}\tilde v_{tt},_{\mu\nu}\cdot\tilde n|^2.$$

This establishes that $\int_0^T \mathfrak{j} = \int_0^T \mathcal{R}$. We thus conclude that

$$\int_0^T \mathfrak{r}_{3b,3} = \int_0^T \mathcal{R}.$$

Hence,

$$\int_0^T \mathfrak{r}_3 \le \int_0^T \mathcal{R}. \tag{4.26}$$

Analysis of $\int_0^T \mathfrak{r}_4$ in the time-integral of (4.16)**.** We integrate by parts with respect to a time derivative of $\tilde v_{ttt}$ in $\int_0^T \mathfrak{r}_4$ and, if need be, spatially integrate by parts. For example, letting $\mathfrak{r}_4 = \sum_{l=1}^4 \mathfrak{r}_{4,l}$, we find that after integration by parts with respect to time,

$$\begin{aligned}\int_0^T \mathfrak{r}_{4,1} &= \int_0^T\int_\Gamma (\sqrt{\tilde g}\tilde g^{\alpha\beta})_t(\tilde\eta,_{\alpha\beta}\cdot\tilde n)_{tttt}\,\tilde v_{ttt}\cdot\tilde n + \int_0^T \mathcal{R} \\ &= -\int_0^T\int_\Gamma \tilde v_{ttt},_\alpha\cdot\bar\partial_\beta[\tilde n\,(\sqrt{\tilde g}\tilde g^{\alpha\beta})_t\tilde v_{ttt}\cdot\tilde n] + \int_0^T \mathcal{R} = \int_0^T \mathcal{R}.\end{aligned}$$

Similarly, integration by parts with respect to time provides for the expression

$$\int_0^T \mathfrak{r}_{4,2} + \int_0^T \mathfrak{r}_{4,3} = \int_0^T \mathcal{R}.$$

Finally, using the differentiation formulas (2.10a) and (2.10b),

$$\int_0^T \mathfrak{r}_{4,4} = \int_0^T \int_\Gamma \sqrt{\tilde g}(2\tilde g^{\alpha\mu}\tilde g^{\nu\beta} - \tilde g^{\alpha\beta}\tilde g^{\mu\nu})\, \tilde v_{ttt},_\mu \cdot \tilde\eta,_\nu \ \tilde\eta,_{\alpha\beta} \cdot \tilde n\, \tilde v_{tttt} \cdot \tilde n + \int_0^T \mathcal{R} = \int_0^T \mathcal{R},$$

where the second equality follows from our above analysis of $\int_0^T \mathfrak{r}_{2b}$.

Hence,

$$\int_0^T \mathfrak{r}_4 \le \int_0^T \mathcal{R}. \tag{4.27}$$

Analysis of $\int_0^T \mathfrak{r}_5$ in the time-integral of (4.16)**.** In the first term defining $\mathfrak{r}_5$, we integrate by parts with respect to $\bar\partial_\alpha$ when $\bar\partial_\beta$ acts on $\tilde n^j$. This term is bounded by $\int_0^T \mathcal{R}$, since thanks to Lemma 2.1,

$$\int_0^T |\sqrt{\kappa}\tilde v_{tttt}|_0^2 \le C \int_0^T |\sqrt{\kappa}\tilde v_{tttt}|_{0.25}^2 \le \delta \int_0^T \|\sqrt{\kappa}\tilde v_{tttt}\|_1^2 + C_\delta\, T \sup_{t\in[0,T]} \|\tilde v_{tttt}(t)\|_0^2.$$

The remaining terms are similarly estimated. Hence,

$$\int_0^T \mathfrak{r}_5 \le \int_0^T \mathcal{R}. \tag{4.28}$$

Rewriting the equation (4.15). Summing the inequalities (4.19), (4.24), (4.26), (4.27), (4.28) yields

$$\sum_{i=1}^5 \int_0^T \mathfrak{r}_i \le \int_0^T \mathcal{R}. \tag{4.29}$$

Thanks to the inequality (4.29) and the identity (4.16), we equivalently write (4.15) as

$$\begin{aligned} &\frac{1}{2}\frac{d}{dt}\int_\Omega \rho_0 |\partial_t^4 \tilde v|^2 + \frac{d}{dt}\int_\Omega \rho_0^2 \tilde J^{-3} |\partial_t^4 \tilde J|^2 + \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\tilde g}\tilde g^{\alpha\beta}\, \tilde v_{ttt},_\alpha \cdot \tilde n\, \tilde v_{ttt},_\beta \cdot \tilde n \\ &\quad + \int_\Gamma \sqrt{\kappa}\tilde v_{tttt},_\beta \cdot \tilde n \sqrt{\tilde g}\tilde g^{\alpha\beta} \sqrt{\kappa}\tilde v_{tttt},_\alpha \cdot \tilde n + \int_\Omega \rho_0^2 \tilde J^{-1} |\sqrt{\kappa}\partial_t^5 \tilde J|^2 = \mathcal{R}. \end{aligned} \tag{4.30}$$

The time-integral of (4.30) completes the proof. ☐

Via Proposition 2.1, the estimates of Lemma 4.2 and Proposition 4.1 imply the following

Proposition 4.2 (The κ-independent estimates for $\tilde v_{ttt}$ and $\sqrt{\kappa}\tilde v_{tttt}$)**.**

$$\sup_{t\in[0,T]} \|\tilde v_{ttt}(t)\|_1^2 + \int_0^T \|\sqrt{\kappa}\tilde v_{tttt}\|_1^2 \le \int_0^T \mathcal{R}.$$

Proof. The fundamental theorem of calculus provides for

$$\operatorname{div} \partial_t^a \tilde\eta = \tilde a_r^s \partial_t^a \tilde\eta^r,_s - \partial_t^a \tilde\eta^r,_s \int_0^t \tilde a_{t\,r}^{\,s}, \quad a = 0, 1, 2, 3, 4. \tag{4.31}$$

The identity $\tilde J_t = \tilde a_r^s \tilde v^r,_s$ implies that for $a = 1, 2, 3$, $\partial_t^a \tilde J_t$ is equal to $\tilde a_r^s \partial_t^a \tilde v^r,_s + \tilde{\jmath}$, where $\tilde{\jmath}$ scales like $D\partial_t^{a-1}\tilde v$. The estimate for $\partial_t^4 \tilde J$ stated in Proposition 4.1 therefore implies that

$$\sup_{t\in[0,T]} \|\operatorname{div} \tilde v_{ttt}(t)\|_0^2 \le \int_0^T \mathcal{R}. \tag{4.32}$$

Using the identity $N = \tilde{n} - \int_0^t \partial_t \tilde{n}$, and the estimate

$$|\bar{\partial}\partial_t^a \tilde{v} \cdot \int_0^t \partial_t \tilde{n}|^2_{2.5-a} \leq T\, C \|\partial_t^a \tilde{v}\|^2_{4-a} \quad \text{for } a = 0, 1, 2, 3,$$

we infer from the trace-estimate stated in Proposition 4.1 that

$$\sup_{t\in[0,T]} |\bar{\partial}\tilde{v}_{ttt}(t) \cdot N|^2_{-0.5} \leq \int_0^T \mathcal{R}.$$

Thanks to the curl-estimates of Lemma 4.2, Proposition 2.1 therefore establishes that

$$\sup_{t\in[0,T]} \|\tilde{v}_{ttt}(t)\|^2_1 \leq \int_0^T \mathcal{R}.$$

A similar analysis establishes the $L^2(0,T; H^1(\Omega))$-estimate for $\sqrt{\kappa}\tilde{v}_{tttt}$. □

We now improve the normal trace-estimates of Proposition 4.1.

Proposition 4.3 (An improved κ-independent normal trace-estimate for $\tilde{v}_{ttt}$)**.**

$$\sup_{t\in[0,T]} |\tilde{v}_{ttt} \cdot \tilde{n}(t)|^2_1 \leq \int_0^T \mathcal{R}.$$

Proof. The estimate of Proposition 4.2 together with the trace theorem implies that

$$\sup_{t\in[0,T]} |\tilde{v}_{ttt}|^2_{0.5} \leq \int_0^T \mathcal{R}.$$

Combining this estimate with the normal trace-estimate of Proposition 4.1 completes the proof. □

Step 3: The κ-independent divergence- and normal trace-estimates. We equivalently write the approximate surface tension problem (4.13a) as

$$2\rho_0 \tilde{J}^{-2} \tilde{A}^k_i \tilde{J},_k + \kappa\rho_0 \tilde{A}^k_i \tilde{J}_t,_k = \tilde{v}^i_t + [2\tilde{J}^{-1} - \kappa\tilde{J}_t]\tilde{A}^k_i \rho_0,_k\,,$$

or equivalently, setting $\kappa' = \kappa/2$,

$$\tilde{J}^{-2} \tilde{A}^k_{\cdot} \tilde{J},_k + \kappa' \tilde{A}^k_{\cdot} \tilde{J}_t,_k = \underbrace{\tfrac{1}{2\rho_0}[\tilde{v}_t + [2\tilde{J}^{-1} - \kappa\tilde{J}_t]\tilde{A}^k_{\cdot} \rho_0,_k\,]}_{\tilde{\mathcal{V}}_t}\,. \tag{4.33}$$

The fundamental theorem of calculus provides that (4.33) is equivalently written as

$$D\tilde{J} + \kappa' D\tilde{J}_t = \tilde{\mathcal{V}}_t - \underbrace{[\tilde{J},_k \int_0^t \partial_t(\tilde{J}^{-2}\tilde{A}^k_{\cdot}) + \kappa\tilde{J}_t,_k \int_0^t \partial_t \tilde{A}^k_{\cdot}]}_{\tilde{\jmath}}\,. \tag{4.34}$$

Lemma 4.3 (Estimates for $\tilde{J}_{ttt}$ and $\kappa\, \tilde{J}_{tttt}$ via Lemma 2.4)**.**

$$\sup_{t\in[0,T]} \|\tilde{J}_{ttt}(t)\|^2_1 + \sup_{t\in[0,T]} \|\kappa\, \tilde{J}_{tttt}(t)\|^2_1 \leq \int_0^T \mathcal{R}.$$

Proof. Taking three time-derivatives of (4.34) produces an equation of the form $f + \kappa f_t = g$:

$$D\tilde{J}_{ttt} + \kappa' \partial_t(D\tilde{J}_{ttt}) = \partial_t^3 \tilde{\mathcal{V}}_t + \tilde{\jmath}_{ttt}. \tag{4.35}$$

By (4.33) we have that $\partial_t^3\tilde{\mathcal{V}}_t$ scales like $\tilde{v}_{tttt} + T\, D\tilde{v}_{ttt} + \kappa\partial_t^5 \tilde{J}$. Proposition 4.2 therefore implies that $\|\partial_t^3 \tilde{\mathcal{V}}_t(t)\|^2_0 \leq \int_0^T \mathcal{R}$. According to (4.34), we have that $\tilde{\jmath}_{ttt}$ scales like $T\, D\tilde{J}_{ttt} +$

$T\,\kappa D\tilde{J}_{tttt} + T\,D\tilde{v}_{ttt}$. So, $\|\tilde{j}_{ttt}(t)\|_0^2 \leq \int_0^T \mathcal{R}$. The fundamental theorem of calculus provides a good estimate for $\tilde{J}_{ttt}$ in $L^\infty(0,T;L^2(\Omega))$. Thus, Lemma 2.4 implies that

$$\sup_{t\in[0,T]} \|\tilde{J}_{ttt}(t)\|_1^2 \leq \int_0^T \mathcal{R}.$$

We similarly infer from equation (4.35) and the fundamental theorem of calculus that

$$\sup_{t\in[0,T]} \|\kappa\tilde{J}_{tttt}(t)\|_1^2 \leq \int_0^T \mathcal{R}.$$

This completes the proof. ☐

Lemma 4.4 (Normal trace-estimates for $\tilde{v}_{tt}$ and $\kappa\tilde{v}_{ttt}$)**.**

$$\sup_{t\in[0,T]} |\bar{\partial}\tilde{v}_{tt}\cdot\tilde{n}(t)|_{0.5}^2 + \sup_{t\in[0,T]} |\kappa\bar{\partial}\tilde{v}_{ttt}\cdot\tilde{n}(t)|_{0.5}^2 \leq \int_0^T \mathcal{R}.$$

Proof. The fundamental theorem of calculus implies the desired estimates. For example,

$$\begin{aligned} |\bar{\partial}\tilde{v}_{tt}\cdot\tilde{n}(t)|_{0.5}^2 &\leq C|\tilde{v}_{tt}\cdot\tilde{n}(t)|_{0.5}|\bar{\partial}\tilde{v}_{tt}\cdot\tilde{n}(t)|_{1.5} \\ &\leq \mathcal{N}_0 + C_\delta\int_0^T \|\tilde{v}_{ttt}\|_1^2 + \delta\sup_{t\in[0,T]}|\bar{\partial}\tilde{v}_{tt}\cdot\tilde{n}(t)|_{1.5}^2 \leq \int_0^T \mathcal{R}. \end{aligned}$$

☐

Step 4: The κ-independent higher-order estimates via Proposition 2.1. The divergence- and normal trace-estimates obtained in Step 3, together with the curl-estimates of Lemma 4.2, imply a good estimate for $\tilde{v}_{tt}$. Hence, successively repeating Step 3 yields

$$\sup_{t\in[0,T]}\sum_{a=0}^{2}\|\partial_t^a\tilde{v}(t)\|_{4-a}^2 + \sup_{t\in[0,T]}\sum_{a=0}^{2}\|\kappa\partial_t^a\tilde{v}_t(t)\|_{4-a}^2 \leq \int_0^T \mathcal{R}. \tag{4.36}$$

We recall the boundary condition (3.1b) is

$$\tilde{g}^{\alpha\beta}\tilde{\eta},_{\alpha\beta}\cdot\tilde{n} + \kappa\tilde{g}^{\alpha\beta}\tilde{v},_{\alpha\beta}\cdot\tilde{n} = \underbrace{\beta(t) - \rho_0^2\tilde{J}^{-2} + \kappa\rho_0^2\tilde{J}^{-1}\tilde{J}_t}_{\tilde{j}}. \tag{4.37}$$

Writing the boundary condition (4.37) as an equation of the form $f + \kappa f_t = g$ yields

$$\tilde{g}^{\alpha\beta}\tilde{\eta},_{\alpha\beta}\cdot\tilde{n} + \kappa\partial_t(\tilde{g}^{\alpha\beta}\tilde{\eta},_{\alpha\beta}\cdot\tilde{n}) = \tilde{j} + \kappa\tilde{\eta},_{\alpha\beta}\cdot(\tilde{n}\tilde{g}^{\alpha\beta})_t. \tag{4.38}$$

As $\kappa\tilde{\eta}$ is of the same regularity as $\tilde{\eta}$, the flow map for the κ-problem does not witness an improved boundary-regularity. The fundamental theorem of calculus, however, does imply a good estimate for the right-hand side of (4.38). Hence, we infer from Lemma 2.4 that the normal trace of $\tilde{\eta}$ and $\kappa\tilde{v}$ each has a good estimate in $L^\infty(0,T;H^{4.5}(\Gamma))$:

$$\sup_{t\in[0,T]}\|\tilde{\eta}(t)\|_5^2 + \sup_{t\in[0,T]}\|\kappa\tilde{v}(t)\|_5^2 \leq \int_0^T \mathcal{R}. \tag{4.39}$$

We collect the estimates (4.36) and (4.39) in the following

Proposition 4.4 (Higher-order κ-independent estimates)**.**

$$\sup_{t\in[0,T]}\sum_{a=0}^{3}\|\partial_t^a\tilde{\eta}(t)\|_{5-a}^2 + \sup_{t\in[0,T]}\sum_{a=0}^{3}\|\kappa\partial_t^a\tilde{v}_t(t)\|_{5-a}^2 \leq \int_0^T \mathcal{R}.$$

Step 5: The κ-independent improved boundary-regularity estimates. With the estimates provided by Proposition 4.4 for $\|\partial_t^a\tilde\eta\|^2_{5-a}$ and $\|\kappa\partial_t^a\tilde v\|^2_{5-a}$, $a=0,1,2,3$, we are in position to establish the improved boundary-regularity estimates.

Proposition 4.5 (The κ-independent improved boundary-regularity via Lemma 2.4)**.**

$$\sup_{t\in[0,T]}\sum_{a=0}^{2}|\bar\partial^2\partial_t^a\tilde v\cdot\tilde n(t)|^2_{2.5-a}+\sup_{t\in[0,T]}\sum_{a=0}^{2}|\bar\partial^2\partial_t^a\tilde v_t\cdot\tilde n(t)|^2_{2.5-a}\le\int_0^T\mathcal{R}.$$

Proof. Taking three time-derivatives of (4.37), we obtain

$$\tilde g^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n+\kappa\partial_t(\tilde g^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n)=\partial_t^3\tilde\jmath-\underbrace{[\partial_t^3(\tilde g^{\alpha\beta}\tilde\eta,_{\alpha\beta}\cdot\tilde n)-\tilde g^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n]}_{\tilde\jmath_A}$$
$$-\underbrace{[\kappa\partial_t^3(\tilde g^{\alpha\beta}\tilde v,_{\alpha\beta}\cdot\tilde n)-\kappa\partial_t(\tilde g^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n)]}_{\tilde\jmath_B}. \qquad (4.40)$$

By Proposition 4.4, the right-hand side of (4.40) has a good estimate in $L^\infty(0,T;H^{0.5}(\Gamma))$. Lemma 2.4 therefore provides that

$$\sup_{t\in[0,T]}|\bar\partial^2\tilde v_{tt}\cdot\tilde n(t)|^2_{0.5}\le\int_0^T\mathcal{R}.$$

Since $\kappa\partial_t(\tilde g^{\alpha\beta}\tilde v_{tt},_{\alpha\beta}\cdot\tilde n)=\kappa\tilde g^{\alpha\beta}\tilde v_{ttt},_{\alpha\beta}\cdot\tilde n+\kappa\tilde v_{tt},_{\alpha\beta}\cdot(\tilde n\tilde g^{\alpha\beta})_t$, it follows from (4.40) that

$$\sup_{t\in[0,T]}|\kappa\bar\partial^2\tilde v_{ttt}\cdot\tilde n(t)|^2_{0.5}\le\int_0^T\mathcal{R}.$$

The higher-order estimates are similarly established. □

Step 6: Concluding the proof of Lemma 4.1. The sum of the estimates given in Propositions 4.1–4.5 competes the proof of Lemma 4.1. Taking δ sufficiently small in the inequality (4.5) yields a polynomial-type inequality of the form (2.19). Hence, for sufficiently small $T>0$ and independent of $\kappa>0$,

$$\sup_{t\in[0,T]}E^\kappa(t)\le 2\mathcal{N}_0, \qquad (4.41)$$

where the higher-order energy function $E^\kappa(t)$ is defined in (4.1).

5. CONSTRUCTION OF SOLUTIONS TO THE κ-PROBLEM (3.1)

In this section, we prove the following

Theorem 5.1 (Solutions to the κ-problem)**.** *For C^∞-class initial data (ρ_0,u_0,Ω) satisfying the conditions* (1.13) *and* (1.14)*, and for some $T=T_\kappa>0$, there exists a unique solution $\tilde v$ to the κ-problem* (3.1) *verifying $(\tilde v,\tilde v_t,\dots,\tilde v_{tttt})|_{t=0}=(u_0,\mathrm{v}_1,\dots,\mathrm{v}_4)$ and*

$$\sup_{t\in[0,T]}\|\tilde v_{tttt}(t)\|_0^2+\int_0^T|\tilde v_{tttt}\cdot\tilde n|_1^2+\sum_{a=0}^{4}\int_0^T\|\partial_t^a\tilde v\|^2_{9-2a}<\infty. \qquad (5.1)$$

Remark 12. We recall that the initial data v_a, $a=1,2,3,4$, is defined in Section 3.3.

We establish Theorem 5.1 via a succession of two asymptotic estimates that correspond with two intermediate approximate problems defined below. Each of the two intermediate problems involves the use of a convolution operator that smooths in the direction tangent to the moving boundary. We use $\epsilon>0$ and $\mu>0$ as the smoothing parameters.

Our first intermediate problem, which we call the $\kappa\epsilon$-problem, is defined by smoothing the moving domain of the κ-problem. The overall structure of the $\kappa\epsilon$-problem matches that of the κ-problem. It naturally follows that the ϵ-independent a priori estimates closely resemble the κ-independent a priori estimates of Section 4. The $\epsilon = 0$ formal limit of the $\kappa\epsilon$-problem is the κ-problem.

Our second intermediate problem, which we call the μ-problem, is defined by smoothing the κ-artificial viscosity term $\kappa g_\epsilon^{\alpha\beta} v,_{\alpha\beta} \cdot n_\epsilon$ appearing in the boundary condition of the $\kappa\epsilon$-problem. The μ-problem is a nonlinear heat-type problem with Neumann-type boundary conditions. The $\mu = 0$ formal limit of the heat-type μ-problem is equivalent to the $\kappa\epsilon$-problem. The key to obtaining the μ-independent a priori estimates is that the diffusive term of the heat-type μ-problem yields a trace-estimate on the boundary Γ.

5.1. **Horizontal convolution-by-layers.** For $\epsilon > 0$, we let $0 \leq \bar\varphi_\epsilon \in C_0^\infty(\mathbb{R}^2)$ with $\operatorname{spt}(\bar\varphi_\epsilon) \subset \overline{B(0,\epsilon)}$ denote the family of standard mollifiers on $\mathbb{R}^2$. With $x_h = (x_1, x_2)$, we define the operation of *horizontal convolution-by-layers* as follows:

$$\Lambda_\epsilon f(x_h, x_3) = \int_{\mathbb{R}^2} \bar\varphi_\epsilon(x_h - y_h) f(y_h, x_3) dy_h \quad \text{for } f \in L^1_{loc}(\mathbb{R}^2).$$

By standard properties of convolution, there exists a constant C which is independent of ϵ, such that for $s \geq 0$,

$$|\Lambda_\epsilon f|_s \leq C|f|_s \quad \forall f \in H^s(\Gamma).$$

Furthermore,

$$\epsilon |\Lambda_\epsilon f|_1 \leq C|f|_0 \quad \forall f \in L^2(\Gamma). \tag{5.2}$$

We recall the local coordinates near Γ are defined in Section 2.1.5. We set

$$\epsilon_0 = \min_{l=1}^{K} \operatorname{dist}(\operatorname{spt} \xi_l \circ \theta_l, \partial\mathcal{B}_l^+ \setminus \mathcal{D}_l). \tag{5.3}$$

We define the horizontally-convolved vector field v_ϵ on Γ of a given vector v by

$$v_\epsilon = \sum_{l=1}^{K} \Lambda_\epsilon[(\xi_l v) \circ \theta_l] \circ \theta_l^{-1} \quad \text{on } \Gamma. \tag{5.4}$$

Given a sufficiently smooth vector field $\check v$, we set $\check\eta = e + \int_0^t \check v$ in Ω and $\check\eta_\epsilon = e + \int_0^t \check v_\epsilon$ on Γ. We define $\check\zeta_\epsilon$ to be the solution of the following time-dependent elliptic Dirichlet problem:

$$\Delta\check\zeta_\epsilon = \Delta\check\eta \quad \text{in } \Omega, \tag{5.5a}$$

$$\check\zeta_\epsilon = \check\eta_\epsilon \quad \text{on } \Gamma. \tag{5.5b}$$

We define the following ϵ-approximate Lagrangian variables:

$$\check A_\epsilon = [D\check\zeta_\epsilon]^{-1}, \quad \check J_\epsilon = \det D\check\zeta_\epsilon, \quad \check a_\epsilon = \check J_\epsilon \check A_\epsilon, \quad [\check g_\epsilon]_{\alpha\beta} = \check\zeta_\epsilon,_\alpha \cdot \check\zeta_\epsilon,_\beta, \text{ and } \sqrt{\check g_\epsilon}\,\check n_\epsilon = [\check a_\epsilon]^T N.$$

5.2. **The $\kappa\epsilon$-problem and its a priori estimates.** We define an intermediate approximate problem, which we will refer to as the $\kappa\epsilon$-problem, that is asymptotically consistent with the κ-problem (3.1). To indicate the dependence on the smoothing parameter ϵ of all the variables in the $\kappa\epsilon$-problem, we place the symbol ˇ over each of the variables. In the $\kappa\epsilon$-problem, we smooth the moving boundary and use the corresponding twice-mean-curvature function

$$\sigma H(\check\zeta_\epsilon) = -\sigma \check g_\epsilon^{\alpha\beta} \check\zeta_\epsilon,_{\alpha\beta} \cdot \check n_\epsilon.$$

Definition 5.1 (The $\kappa\epsilon$-problem)**.** *For $\kappa > 0$ and $\epsilon > 0$, we define $\check{v}$ as the solution of*

$$\check{v}_t^i + 2[\check{A}_\epsilon]_i^k(\rho_0 \check{J}_\epsilon^{-1}),_k - \kappa[\check{A}_\epsilon]_i^k(\rho_0 \check{\mathcal{J}}_t),_k = 0 \qquad \text{in } \Omega \times (0, T_\kappa(\epsilon)], \tag{5.6a}$$

$$\rho_0^2 \check{J}_\epsilon^{-2} - \kappa\rho_0^2 \check{J}_\epsilon^{-1}\check{\mathcal{J}}_t = \check{H}_\epsilon \qquad \text{on } \Gamma \times (0, T_\kappa(\epsilon)], \tag{5.6b}$$

$$(\check{\zeta}_\epsilon, \check{v})|_{t=0} = (e, u_0) \quad \text{on } \Omega. \tag{5.6c}$$

The function $\check{\mathcal{J}}_t$ appearing in the left-hand side of equations (5.6a) *and* (5.6b) *is defined as*

$$\check{\mathcal{J}}_t = \check{J}_\epsilon \operatorname{div}_{\check{\zeta}_\epsilon} \check{v}. \tag{5.7}$$

The function $\check{H}_\epsilon$ appearing in the right-hand side of (5.6b) *is defined as*

$$\check{H}_\epsilon = \beta_\epsilon(t) - \sigma\check{g}_\epsilon^{\alpha\beta}\check{\zeta}_\epsilon,_{\alpha\beta} \cdot \check{n}_\epsilon - \kappa\check{g}_\epsilon^{\alpha\beta}\check{v},_{\alpha\beta} \cdot \check{n}_\epsilon. \tag{5.8}$$

The function $\beta_\epsilon(t)$ appearing in the right-hand side of (5.8) *is defined as*

$$\beta_\epsilon(t) = \beta + \sum_{a=0}^{3} \frac{t^a}{a!}\partial_t^a[\rho_0^2 \check{J}_\epsilon^{-2} - \beta + \sigma\check{g}_\epsilon^{\alpha\beta}\check{\zeta}_\epsilon,_{\alpha\beta} \cdot \check{n}_\epsilon]$$
$$+ \kappa\sum_{a=0}^{3} \frac{t^a}{a!}\partial_t^a[-\rho_0^2 \check{J}_\epsilon^{-1}\check{\mathcal{J}}_t + \check{g}_\epsilon^{\alpha\beta}\check{v},_{\alpha\beta} \cdot \check{n}_\epsilon]|_{t=0}. \tag{5.9}$$

Remark 13. Elliptic regularity provides that The ϵ-approximate Lagrangian flow map $\check{\zeta}_\epsilon$ defined by (5.5) satisfies $\|\check{\zeta}_\epsilon\|_s \le C\|\check{\eta}\|_s$ for a positive constant C independent of ϵ. Standard properties of convolution provide that setting $\epsilon = 0$ in (5.5) yields $\check{\zeta}_\epsilon = \check{\eta}$. It follows that the $\kappa\epsilon$-problem (5.6) is asymptotically consistent with the κ-problem (3.1).

5.2.1. *The constant-in-time vectors* $\mathbf{v}_a$ *for* $a = 1, 2, 3, 4$. Letting $\check{v}$ solve (5.6), the vector field $\partial_t^a \check{v}|_{t=0}$ for all $a \in \mathbb{N}$ is computed as follows:

$$\partial_t^a \check{v}|_{t=0} = \frac{\partial^{a-1}}{\partial t^{a-1}}\left(\kappa[\check{A}_\epsilon]_\cdot^k(\rho_0\check{\mathcal{J}}_t),_k - 2[\check{A}_\epsilon]_\cdot^k(\rho_0\check{J}_\epsilon^{-1}),_k\right)\Big|_{t=0} \quad \text{on } \Omega.$$

This formula makes it clear that each $\partial_t^a \check{v}|_{t=0}$ is a function of space-derivatives of the initial data u_0, $\Lambda_\epsilon u_0$ and ρ_0. We define the constant-in-time vectors $\mathbf{v}_a$ as

$$\mathbf{v}_a = \frac{\partial^{a-1}}{\partial t^{a-1}}\left(\kappa[\check{A}_\epsilon]_\cdot^k(\rho_0\check{\mathcal{J}}_t),_k - 2[\check{A}_\epsilon]_\cdot^k(\rho_0\check{J}_\epsilon^{-1}),_k\right)\Big|_{t=0}, \quad \text{for } a = 1, 2, 3, 4. \tag{5.10}$$

We have that $\mathbf{v}_a \to \mathrm{v}_a$ as $\epsilon \to 0$ where v_a are defined in Section 3.3. We use (5.6b) and the definition (5.9) of $\beta_\epsilon(t)$ to compute the following identities: for $a = 0, 1, 2, 3$,

$$\partial_t^a[\rho_0^2\check{J}_\epsilon^{-2} - \kappa\rho_0^2\check{J}_\epsilon^{-1}\check{\mathcal{J}}_t]\big|_{t=0} = \partial_t^a[\beta_\epsilon(t)]\big|_{t=0} - \partial_t^a[\sigma\check{g}_\epsilon^{\alpha\beta}\check{\zeta}_\epsilon,_{\alpha\beta} \cdot \check{n}_\epsilon + \kappa\check{g}_\epsilon^{\alpha\beta}\check{v},_{\alpha\beta} \cdot \check{n}_\epsilon]\big|_{t=0}. \tag{5.11}$$

5.2.2. *A priori estimates for the $\kappa\epsilon$-problem* (5.6). For $\epsilon > 0$, we define the following higher-order energy function:

$$E^\epsilon(t) = 1 + \|\check{v}_{tt}(t)\|_0^2 + \int_0^t |\check{v}_{tt} \cdot \check{n}_\epsilon|_1^2 + \sum_{a=0}^{2}\int_0^t \|\partial_t^a \check{v}\|_{5-2a}^2. \tag{5.12}$$

Definition 5.2 (Notational convention for constants depending on $1/\delta\kappa > 0$)**.** *We let $\dot{\mathcal{P}}$ denote a generic polynomial with constant and coefficients depending on $1/\delta\kappa > 0$.*

We define the constant $\dot{\mathcal{N}}_0 > 0$ by

$$\dot{\mathcal{N}}_0 = \dot{\mathcal{P}}(\|u_0\|_{100}, \|\rho_0\|_{100}).$$

We let $\dot{\mathcal{R}}$ denote generic lower-order terms satisfying

$$\int_0^T \dot{\mathcal{R}} \le \dot{\mathcal{N}}_0 + \delta \sup_{t\in[0,T]} E^\epsilon(t) + T\,\dot{\mathcal{P}}(\sup_{t\in[0,T]} E^\epsilon(t)).$$

We assume $T>0$ is taken sufficiently small to ensure that

$$\frac{1}{2} \le \check{J} \le \frac{3}{2} \ \text{ and } \ \frac{1}{2} \le \check{J}_\epsilon \le \frac{3}{2} \ \text{ for all } t\in[0,T] \text{ and } x\in\Omega.$$

Lemma 5.1 (A priori estimates for the $\kappa\epsilon$-problem)**.** *We let $\check{v}$ solve the $\kappa\epsilon$-problem* (5.6) *on a time-interval* $[0,T]$ *for some $T=T_\kappa(\epsilon)>0$. Then independent of ϵ,*

$$\sup_{t\in[0,T]} E^\epsilon(t) \le \int_0^T \dot{\mathcal{R}}. \tag{5.13}$$

We will establish Lemma 5.1 in the following four steps:

Step 1: The ϵ-independent curl-estimates. Taking the ϵ-approximate Lagrangian curl of the equations (5.6a) yields

$$\operatorname{curl}_{\check{\zeta}_\epsilon} \check{v}_t = 0. \tag{5.14}$$

Integrating the identity (5.14) in time from 0 to $t\in(0,T]$ provides that

$$\operatorname{curl}_{\check{\zeta}_\epsilon} \check{v} = \operatorname{curl} u_0 + \varepsilon_{\cdot ji} \int_0^t \partial_t [\check{A}_\epsilon]^s_j \check{v}^i,_s\,. \tag{5.15}$$

We may therefore infer the curl estimates from Lemma 4.2. We record this fact as

Lemma 5.2 (The ϵ-independent curl-estimates for $\partial^a\check{v}$, for $a=0,1,2$)**.**

$$\sum_{a=0}^{2} \int_0^T \|\operatorname{curl}\partial_t^a \check{v}\|^2_{4-2a} \le \int_0^T \dot{\mathcal{R}}.$$

Step 2: The ϵ-independent estimate for $\check{v}_{tt}$. We equivalently write the momentum equations (5.6a) as

$$\rho_0 \check{v}^i_t + [\check{a}_\epsilon]^k_i (\rho_0^2 \check{J}_\epsilon^{-2}),_k - \kappa\rho_0 \check{J}_\epsilon^{-1} [\check{a}_\epsilon]^k_i (\rho_0 \check{\mathcal{J}}_t),_k = 0 \qquad \text{in } \Omega\times(0,T_\kappa(\epsilon)]. \tag{5.16}$$

Lemma 5.3 (Energy estimates for the action of ∂_t^2 in (5.16))**.**

$$\sup_{t\in[0,T]} \|\check{v}_{tt}(t)\|^2_0 + \int_0^T \|\check{\mathcal{J}}_{ttt}\|^2_0 + \int_0^T |\bar{\partial}\check{v}_{tt}\cdot \check{n}_\epsilon|^2_0 \le \int_0^T \dot{\mathcal{R}}.$$

Proof. Testing the action of ∂_t^2 in (5.16) against $\check{v}_{tt}$ in the $L^2(\Omega)$-inner product and integrating by parts with respect to ∂_k in the interior integrals $\int_\Omega [\check{a}_\epsilon]^k_i \partial_t^2 (\rho_0^2 \check{J}_\epsilon^{-2}),_k \check{v}^i_{tt}$ and $-\kappa \int_\Omega [\check{a}_\epsilon]^k_i \partial_t^2 \big[\rho_0 \check{J}_\epsilon^{-1} (\rho_0 \check{\mathcal{J}}_t),_k\big] \check{v}^i_{tt}$ yields

$$\frac{1}{2}\frac{d}{dt}\int_\Omega \rho_0 |\check{v}_{tt}|^2 - \underbrace{\int_\Omega \partial_t^2[\rho_0^2 \check{J}_\epsilon^{-2}][\check{a}_\epsilon]^k_i \check{v}^i_{tt},_k}_{\dot{\mathcal{R}}} + \kappa \underbrace{\int_\Omega \partial_t^2[\rho_0^2 \check{J}_\epsilon^{-1}\check{\mathcal{J}}_t][\check{a}_\epsilon]^k_i \check{v}^i_{tt},_k}_{\mathcal{I}}$$
$$+ \underbrace{\int_\Gamma \partial_t^2[\rho_0^2 \check{J}_\epsilon^{-2} - \kappa\rho_0^2 \check{J}_\epsilon^{-1} \check{\mathcal{J}}_t][\check{a}_\epsilon]^k_i \check{v}^i_{tt} N^k}_{\mathfrak{i}} = \dot{\mathcal{R}}. \tag{5.17}$$

It is convenient to write the equation (5.17) as

$$\frac{1}{2}\frac{d}{dt}\int_\Omega \rho_0|\check{v}_{tt}|^2 + \kappa\int_\Omega \rho_0^2 \check{J}_\epsilon^{-1}|\check{\mathcal{J}}_{ttt}|^2 + \mathfrak{i} = \dot{\mathcal{R}}, \tag{5.18}$$

where the error created in order to write $[\check{a}_\epsilon]_i^k \partial_t^2 \check{v}^i,_k$ as $\check{\mathcal{J}}_{ttt}$ in $\mathcal{I}$ is of lower-order and so is absorbed in $\dot{\mathcal{R}}$.

We write the boundary integral $\mathfrak{i}$ appearing in the left-hand side of (5.18) as

$$\mathfrak{i} = \int_\Gamma \check{H}_{\epsilon tt}\sqrt{\check{g}_\epsilon}\check{v}_{tt}\cdot \check{n}_\epsilon, \tag{5.19}$$

where we have used the boundary condition (5.6b) and the formula (2.9). The definition (5.8) provides that $\check{H}_\epsilon = \beta_\epsilon(t) - \sigma \check{g}_\epsilon^{\alpha\beta}\check{\zeta}_\epsilon,_{\alpha\beta}\cdot\check{n}_\epsilon - \kappa\check{g}_\epsilon^{\alpha\beta}\check{v},_{\alpha\beta}\cdot\check{n}_\epsilon$. Thus,

$$\mathfrak{i} = \kappa\int_\Gamma \sqrt{\check{g}_\epsilon}\check{g}_\epsilon^{\alpha\beta}\check{v}_{tt},_\beta\cdot\check{n}_\epsilon\check{v}_{tt},_\alpha\cdot\check{n}_\epsilon + \kappa\underbrace{\int_\Gamma \check{v}^i_{tt},_\beta\left(\check{n}^i_\epsilon\sqrt{\check{g}_\epsilon}\check{g}_\epsilon^{\alpha\beta}\check{n}^j_\epsilon\right),_\alpha\check{v}^j_{tt}}_{\mathfrak{j}} + \dot{\mathcal{R}}. \tag{5.20}$$

Thanks to Lemma 2.1,

$$\int_0^T |\check{v}_{tt}|_0^2 \le C\int_0^T |\check{v}_{tt}|_{0.25}^2 \le \int_0^T \dot{\mathcal{R}}. \tag{5.21}$$

Employing the Cauchy-Schwarz inequality and Young's inequality with $\delta > 0$ thus yields

$$\int_0^T \mathfrak{j} - \underbrace{\int_0^T \check{v}_{tt},_\beta\cdot\check{n}_\epsilon,_\alpha\sqrt{\check{g}_\epsilon}\check{g}_\epsilon^{\alpha\beta}\check{v}_{tt}\cdot\check{n}_\epsilon}_{\mathfrak{j}'} \le \int_0^T \dot{\mathcal{R}}.$$

Integrating by parts with respect to $\bar{\partial}_\beta$, we similarly have that $\int_0^T \mathfrak{j}' \le \int_0^T \dot{\mathcal{R}}$. We have therefore established that the identity (5.20) is equivalently written as

$$\mathfrak{i} = \kappa\int_\Gamma \sqrt{\check{g}_\epsilon}\check{g}_\epsilon^{\alpha\beta}\check{v}_{tt},_\beta\cdot\check{n}_\epsilon\check{v}_{tt},_\alpha\cdot\check{n}_\epsilon + \dot{\mathcal{R}}.$$

Using this identity for $\mathfrak{i}$ in the time-integral of (5.18) completes the proof. ☐

Proposition 5.1 (The ϵ-independent estimates for $\check{v}_{tt}$)**.**

$$\sup_{t\in[0,T]}\|\check{v}_{tt}(t)\|_0^2 + \int_0^T |\check{v}_{tt}\cdot\check{n}_\epsilon|_1^2 + \int_0^T \|\check{v}_{tt}\|_1^2 \le \dot{\mathcal{R}}.$$

Proof. The desired $L^\infty(0,T;L^2(\Omega))$-estimate is provided by Lemma 5.3. The inequality (5.21) and the trace-estimate stated in Lemma 5.3 establish the desired $L^2(0,T;H^1(\Gamma))$-estimate. We infer from the arguments proving Proposition 4.2 that Lemma 5.3 implies a divergence- and normal trace-estimate for $\check{v}_{tt}$. Thanks to the curl-estimate for $\check{v}_{tt}$ stated in Lemma 5.2, we have by Proposition 2.1 that the proof is complete. ☐

Step 3: The ϵ-independent estimate for $\check{v}_t$.

Proposition 5.2 (The ϵ-independent estimate for $\check{v}_t$)**.**

$$\int_0^T \|\check{v}_t\|_3^2 \le \int_0^T \dot{\mathcal{R}}.$$

Proof. We write the boundary condition (5.6b) as

$$\kappa \check{g}_\epsilon^{\alpha\beta}\check{v},_{\alpha\beta}\cdot\check{n}_\epsilon = \underbrace{\kappa\rho_0^2\check{J}_\epsilon^{-1}\check{\mathcal{J}}_t - \rho_0^2\check{J}_\epsilon^{-2} + \beta_\epsilon(t) - \sigma\check{g}_\epsilon^{\alpha\beta}\check{\zeta}_\epsilon,_{\alpha\beta}\cdot\check{n}_\epsilon}_{\check{\jmath}}.$$

Taking a time-derivative of this equation yields

$$\kappa\check{g}_\epsilon^{\alpha\beta}\check{v}_t,_{\alpha\beta}\cdot\check{n}_\epsilon = \check{\jmath}_t + \kappa\check{v},_{\alpha\beta}\cdot\partial_t(\check{n}_\epsilon\check{g}_\epsilon^{\alpha\beta}). \tag{5.22}$$

The right-hand side of (5.22) scales like $T\,\bar{\partial}^2\check{v}_t$. We infer that

$$\int_0^T|\bar{\partial}^2\check{v}_t\cdot\check{n}_\epsilon|_{0.5}^2 \le \int_0^T\dot{\mathcal{R}}. \tag{5.23}$$

We equivalently write the momentum equations (5.16) as

$$[\check{A}_\epsilon]_i^k\check{\mathcal{J}}_t,_k = \underbrace{(\kappa\rho_0)^{-1}\big[\check{v}_t^i + 2[\check{A}_\epsilon]_i^k(\rho_0\check{J}_\epsilon^{-1}),_k - \kappa\check{\mathcal{J}}_t[\check{A}_\epsilon]_i^k\rho_0,_k\big]}_{\check{\mathcal{V}}_t}.$$

Using the fundamental theorem of calculus, we have that

$$D\check{\mathcal{J}}_t = \check{\mathcal{V}}_t + \check{\mathcal{J}}_t,_k\int_0^t\partial_t[\check{A}_\epsilon]_i^k. \tag{5.24}$$

Since the right-hand side of (5.24) scales like $\check{v}_t + D\check{v}$, a time-derivative of (5.24) yields

$$\int_0^T\|D\check{\mathcal{J}}_{tt}\|_1^2 \le \int_0^T\dot{\mathcal{R}}.$$

Lemma 5.3 provides a good estimate for $\check{\mathcal{J}}_{tt}$ in $L^2(0,T;L^2(\Omega))$. Thus,

$$\int_0^T\|\check{\mathcal{J}}_{tt}\|_2^2 \le \int_0^T\dot{\mathcal{R}}. \tag{5.25}$$

Via Proposition 2.1, the curl-, normal trace- and divergence-estimates for $\check{v}_t$, respectively given in Lemma 5.2 and inequalities (5.23) and (5.25), complete the proof. ☐

Step 4: Concluding the proof of Lemma 5.1. Repeating Step 3 establishes

Proposition 5.3 (The ϵ-independent estimate for $\check{v}$)**.**

$$\int_0^T\|\check{v}\|_5^2 \le \int_0^T\dot{\mathcal{R}}.$$

The sum of the ϵ-independent estimates stated in Propositions 5.1, 5.2 and 5.3 completes the proof of Lemma 5.1.

As an intermediate step in proving Theorem 5.1, we will establish the following

Theorem 5.2 (Solutions to the $\kappa\epsilon$-problem)**.** *For C^∞-class initial data (ρ_0, u_0, Ω) satisfying the conditions* (1.13) *and* (1.14)*, and for some $T = T_\kappa(\epsilon) > 0$, there exists a unique solution $\check{v}$ to the $\kappa\epsilon$-problem* (5.6) *verifying $(\check{v}, \check{v}_t, \ldots, \check{v}_{tttt})|_{t=0} = (u_0, \mathbf{v}_1, \ldots, \mathbf{v}_4)$ and*

$$\sup_{t\in[0,T]}\|\check{v}_{tttt}(t)\|_0^2 + \sum_{a=0}^4\int_0^T\|\partial_t^a\check{v}\|_{9-2a}^2 + \sum_{a=0}^4\int_0^T|\partial_t^a\check{v}\cdot\check{n}_\epsilon|_{9-2a}^2 < \infty. \tag{5.26}$$

Remark 14. We recall that the initial data $\mathbf{v}_a$, $a = 1, 2, 3, 4$, is defined in Section 5.2.1.

5.3. **Deriving a heat-type problem with Neumann-type boundary conditions.** We will derive a nonlinear heat-type problem with Neumann-type boundary conditions which is equivalent to the $\kappa\epsilon$-problem (5.6).

5.3.1. *Rewriting the momentum equations* (5.6a) *via* (5.15). Setting

$$\check{\varrho} = \kappa \rho_0 \check{J}_\epsilon, \tag{5.27}$$

and using the definition (5.7), the momentum equations (5.6a) are equivalently written as

$$\check{v}_t - \check{\varrho}[\check{A}_\epsilon]^k_\cdot (\operatorname{div}_{\check{\zeta}_\epsilon} \check{v}),_k = \operatorname{div}_{\check{\zeta}_\epsilon} \check{v} [\check{A}_\epsilon]^k_\cdot \check{\varrho},_k - 2[\check{A}_\epsilon]^k_\cdot (\rho_0 \check{J}_\epsilon^{-1}),_k . \tag{5.28}$$

Given a sufficiently smooth vector $\check{v}$, the identity $-\Delta\check{v} = \operatorname{curl}\operatorname{curl}\check{v} - D\operatorname{div}\check{v}$ in the ϵ-approximate Lagrangian variables is the identity

$$-[\check{A}_\epsilon]^j_r [[\check{A}_\epsilon]^k_r \check{v},_k],_j = \operatorname{curl}_{\check{\zeta}_\epsilon} \operatorname{curl}_{\check{\zeta}_\epsilon} \check{v} - [\check{A}_\epsilon]^s_\cdot (\operatorname{div}_{\check{\zeta}_\epsilon} \check{v}),_s . \tag{5.29}$$

Using (5.29), the equations (5.28) are equivalently written as

$$\check{v}_t - \check{\varrho}[\check{A}_\epsilon]^j_r \big([\check{A}_\epsilon]^k_r \check{v},_k\big),_j = \check{\varrho} \operatorname{curl}_{\check{\zeta}_\epsilon} (\operatorname{curl}_{\check{\zeta}_\epsilon} \check{v}) + \operatorname{div}_{\check{\zeta}_\epsilon} \check{v} [\check{A}_\epsilon]^k_\cdot \check{\varrho},_k - 2[\check{A}_\epsilon]^k_\cdot (\rho_0 \check{J}_\epsilon^{-1}),_k . \tag{5.30}$$

Thanks to the vorticity equation (5.15), we further have that (5.6a) is equivalent to

$$\check{v}_t - \check{\varrho}[\check{A}_\epsilon]^j_r \big([\check{A}_\epsilon]^k_r \check{v},_k\big),_j = \check{\mathcal{K}}, \tag{5.31}$$

where the vector field $\check{\mathcal{K}}$ appearing in the right-hand side of (5.31) is defined as

$$\check{\mathcal{K}} = \check{\varrho} \operatorname{curl}_{\check{\zeta}_\epsilon} \Big(\operatorname{curl} u_0 + \varepsilon_{\cdot ji} \int_0^t \partial_t [\check{A}_\epsilon]^s_j \check{v}^i,_s\Big) + \operatorname{div}_{\check{\zeta}_\epsilon} \check{v} [\check{A}_\epsilon]^k_\cdot \check{\varrho},_k - 2[\check{A}_\epsilon]^k_\cdot (\rho_0 \check{J}_\epsilon^{-1}),_k . \tag{5.32}$$

5.3.2. *Deriving a Neumann-type boundary condition for* $\check{v}$. We decompose any vector field $\xi \in \mathbb{R}^3$ evaluated on Γ into tangential components and a normal component as

$$\xi = \xi^\alpha \check{g}_\epsilon^{\alpha\beta} \check{\zeta}_\epsilon,_\beta + \xi^3 \check{n}_\epsilon, \tag{5.33}$$

where $\xi^\alpha = \xi \cdot \check{\zeta}_\epsilon,_\alpha$ and $\xi^3 = \xi \cdot \check{n}_\epsilon$. In the special case of a flat boundary, we may use the standard orthogonal unit vectors e_k to write (5.33) as

$$\xi = \xi^\alpha e_\alpha + \xi^3 e_3. \tag{5.34}$$

For the special case where (5.33) takes the form (5.34), we have that

$$\frac{\partial \check{v}^i}{\partial N} = [(\operatorname{curl}\check{v}) \times N]^i + \check{v}^3,_\alpha e^i_\alpha + [\operatorname{div}\check{v} - \check{v}^\alpha,_\alpha] N^i. \tag{5.35}$$

We define the vector $\check{b}$ as the sum

$$\check{b} = \check{b}_{\text{curl}} + \check{b}_{\text{div}}, \tag{5.36}$$

where $\check{b}_{\text{curl}}$ is a tangent vector and $\check{b}_{\text{div}}$ is a normal vector, respectively defined as

$$\check{b}_{\text{curl}} = \frac{\sqrt{\check{g}_\epsilon}}{\check{J}_\epsilon} \big[\check{g}_\epsilon^{\alpha\beta} \check{v}^3,_\beta + \check{v}^\gamma \check{g}_\epsilon^{\gamma\delta} \check{g}_\epsilon^{\alpha\beta} \check{\zeta}_\epsilon,_{\delta\beta} \cdot \check{n}_\epsilon\big] \check{\zeta}_\epsilon,_\alpha , \tag{5.37a}$$

$$\check{b}_{\text{div}} = -\frac{\sqrt{\check{g}_\epsilon}}{\check{J}_\epsilon} \big[\check{g}_\epsilon^{\alpha\beta} \check{v}^\alpha,_\beta + \check{v}^\alpha (\sqrt{\check{g}_\epsilon} \check{g}_\epsilon^{\alpha\beta}),_\beta + \check{v}^3 H(\check{\zeta}_\epsilon)\big] \check{n}_\epsilon. \tag{5.37b}$$

For the moving boundary $\Gamma_\epsilon(t) = \check{\zeta}_\epsilon(\Gamma)$, the identity (5.35) is written as

$$\frac{\check{J}_\epsilon}{\sqrt{\check{g}_\epsilon}} N^j [\check{A}_\epsilon]^j_r [\check{A}_\epsilon]^k_r \check{v}^i,_k = \big((\operatorname{curl}_{\check{\zeta}_\epsilon} \check{v}) \times \check{n}_\epsilon\big)^i + \check{n}^i_\epsilon \operatorname{div}_{\check{\zeta}_\epsilon} \check{v} + \frac{\check{J}_\epsilon}{\sqrt{\check{g}_\epsilon}} \check{b}^i, \tag{5.38}$$

or equivalently,

$$N^j[\check{A}_\epsilon]^j_r[\check{A}_\epsilon]^k_r\check{v},_k = \sqrt{\check{g}_\epsilon}\check{J}_\epsilon^{-1}(\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v})\times\check{n}_\epsilon + \sqrt{\check{g}_\epsilon}\check{J}_\epsilon^{-1}(\operatorname{div}_{\check{\zeta}_\epsilon}\check{v})\check{n}_\epsilon + \check{b}. \tag{5.39}$$

Thanks to (5.15), we have that (5.39) multiplied by $\check{\varrho} = \kappa\rho_0\check{J}_\epsilon$ is equivalent to

$$\check{\varrho}N^j[\check{A}_\epsilon]^j_r[\check{A}_\epsilon]^k_r\check{v},_k = \check{h}_{\rm curl} + \frac{\sqrt{\check{g}_\epsilon}}{\rho_0}\kappa\rho_0^2(\operatorname{div}_{\check{\zeta}_\epsilon}\check{v})\check{n}_\epsilon + \check{\varrho}\check{b}, \tag{5.40}$$

where the tangential vector field $\check{h}_{\rm curl}$ appearing in the right-hand side of (5.40) is defined via the ϵ-approximate Lagrangian vorticity equation (5.15) as

$$\check{h}_{\rm curl} = \check{\varrho}\sqrt{\check{g}_\epsilon}\check{J}_\epsilon^{-1}\big(\operatorname{curl} u_0 + \varepsilon_{\cdot ji}\int_0^t \partial_t[\check{A}_\epsilon]^s_j\check{v}^i,_s\big)\times\check{n}_\epsilon. \tag{5.41}$$

We write the boundary condition (5.6b) as

$$\kappa\rho_0^2\operatorname{div}_{\check{\zeta}_\epsilon}\check{v} = \rho_0^2\check{J}_\epsilon^{-2} - \beta_\epsilon(t) + \sigma\check{g}_\epsilon^{\alpha\beta}\check{\zeta}_\epsilon,_{\alpha\beta}\cdot\check{n}_\epsilon + \kappa\check{g}_\epsilon^{\alpha\beta}\check{v},_{\alpha\beta}\cdot\check{n}_\epsilon. \tag{5.42}$$

Setting

$$\check{\mathfrak{g}}_\epsilon^{\alpha\beta} = \kappa\frac{\sqrt{\check{g}_\epsilon}}{\rho_0}\check{g}_\epsilon^{\alpha\beta}, \tag{5.43}$$

$$\check{h}_{\rm div} = \frac{\sqrt{\check{g}_\epsilon}}{\rho_0}\Big[\rho_0^2\check{J}_\epsilon^{-2} - \beta_\epsilon(t) + \sigma\check{g}_\epsilon^{\alpha\beta}\check{\zeta}_\epsilon,_{\alpha\beta}\cdot\check{n}_\epsilon\Big]\check{n}_\epsilon, \tag{5.44}$$

we find by using the identity (5.42) for $\kappa\rho_0^2\operatorname{div}_{\check{\zeta}_\epsilon}\check{v}$ in the equation (5.40) that

$$\check{\varrho}N^j[\check{A}_\epsilon]^j_r[\check{A}_\epsilon]^k_r\check{v},_k = \check{h}_{\rm curl} + \check{h}_{\rm div} + \check{\mathfrak{g}}_\epsilon^{\alpha\beta}[\check{v},_{\alpha\beta}\cdot\check{n}_\epsilon]\check{n}_\epsilon + \check{\varrho}\check{b}. \tag{5.45}$$

We define the vector field $\check{h}$ as the sum

$$\check{h} = \check{h}_{\rm curl} + \check{h}_{\rm div} + \check{\mathfrak{g}}_\epsilon^{\alpha\beta}[\check{v},_{\alpha\beta}\cdot\check{n}_\epsilon]\check{n}_\epsilon + \check{\varrho}[\check{b}_{\rm curl} + \check{b}_{\rm div}], \tag{5.46}$$

with the vectors $\check{h}_{\rm curl}$, $\check{h}_{\rm div}$, $\check{b}_{\rm curl}$ and $\check{b}_{\rm div}$ respectively given in (5.41), (5.44), (5.37a) and (5.37b), and the function $\check{\varrho}$ defined in (5.27). We equivalently express the identity (5.45) as

$$\check{\varrho}N^j[\check{A}_\epsilon]^j_r[\check{A}_\epsilon]^k_r\check{v},_k = \check{h}. \tag{5.47}$$

Remark 15. We record that the identities (5.42) and (5.44) establish the following identity:

$$\check{\varrho}\sqrt{\check{g}_\epsilon}\check{J}_\epsilon^{-1}\operatorname{div}_{\check{\zeta}_\epsilon}\check{v} = \check{h}_{\rm div}\cdot\check{n}_\epsilon + \check{\mathfrak{g}}_\epsilon^{\alpha\beta}\check{v},_{\alpha\beta}\cdot\check{n}_\epsilon. \tag{5.48}$$

5.3.3. *The heat-type $\kappa\epsilon$-problem with Neumann-type boundary conditions.*

Definition 5.3 (The heat-type $\kappa\epsilon$-problem)**.** *For $\kappa > 0$ and $\epsilon > 0$ given, we define $\check{v}$ as the solution of the nonlinear heat-type system*

$$\check{v}_t - \check{\varrho}[\check{A}_\epsilon]^j_r\big([\check{A}_\epsilon]^k_r\check{v},_k\big),_j = \check{\mathcal{K}} \qquad \text{in } \Omega\times(0,T_\kappa(\epsilon)], \tag{5.49a}$$

$$\check{\varrho}N^j[\check{A}_\epsilon]^j_r[\check{A}_\epsilon]^s_r\check{v},_s = \check{h} + c(t) \qquad \text{on } \Gamma\times(0,T_\kappa(\epsilon)], \tag{5.49b}$$

$$(\check{\zeta}_\epsilon,\check{v})|_{t=0} = (e,u_0) \qquad \text{on } \Omega. \tag{5.49c}$$

The bounded, nonnegative function $\check{\varrho}$ is defined in (5.27)*. The vector fields $\check{\mathcal{K}}$ and $\check{h}$ are respectively defined in* (5.32) *and* (5.46)*. The vector field $c(t)$ appearing in the right-hand side of* (5.49b) *is defined as*

$$c(t) = \sum_{a=0}^{2}\frac{t^a}{a!}\partial_t^a\big[\check{\varrho}N^j[\check{A}_\epsilon]^j_r[\check{A}_\epsilon]^s_r\check{v},_s - \check{h}\big]|_{t=0}. \tag{5.50}$$

Remark 16. The vector fields $\mathbf{v}_a$, for $a = 1, 2, 3, 4$, defined in Section 5.2.1, are equivalently defined using the $(a-1)$th time-derivative of the momentum equations (5.49a) evaluated at time $t = 0$.

Remark 17. According to Appendix C, the heat-type $\kappa\epsilon$-problem (5.49) is equivalent to the $\kappa\epsilon$-problem (5.6).

Proposition 5.4 (Solutions to the heat-type $\kappa\epsilon$-problem (5.49))**.** *For C^∞-class initial data (ρ_0, u_0, Ω) satisfying the conditions* (1.13) *and* (1.14)*, and for some $T = T_\kappa(\epsilon) > 0$, there exists a unique $\check{v} \in L^2(0, T; H^9(\Omega))$ with $\partial_t^a \check{v} \in L^2(0, T; H^{9-2a}(\Omega))$, for $a = 1, 2, 3, 4$, and $\check{v}_{tttt} \in L^\infty(0, T; L^2(\Omega))$ that solves the nonlinear heat-type $\kappa\epsilon$-problem* (5.49) *on a time-interval $[0, T]$ and verifies $(\check{v}, \check{v}_t, \ldots, \check{v}_{tttt})|_{t=0} = (u_0, \mathbf{v}_1, \ldots, \mathbf{v}_4)$.*

5.4. **The μ-problem and its a priori estimates.** We now define our second intermediate problem, which we term the μ-problem. The μ-problem is a system of nonlinear heat-type equations that is asymptotically consistent with the heat-type $\kappa\epsilon$-problem (5.49). To indicate the dependence on the smoothing paramater μ of all the variables in the following problem, we place the symbol $^\circ$ over each of the variables.

Given a sufficiently smooth vector field $\mathring{v}$, we set $\mathring{\eta} = e + \int_0^t \mathring{v}$ in Ω and $\mathring{\eta}_\epsilon = e + \int_0^t \mathring{v}_\epsilon$ on Γ. We define $\mathring{\zeta}_\epsilon$ to be the solution of the following time-dependent elliptic Dirichlet problem:

$$\Delta\mathring{\zeta}_\epsilon = \Delta\mathring{\eta} \quad \text{in } \Omega, \tag{5.51a}$$

$$\mathring{\zeta}_\epsilon = \mathring{\eta}_\epsilon \quad \text{on } \Gamma. \tag{5.51b}$$

We define the following ϵ-approximate Lagrangian variables:

$$\mathring{A}_\epsilon = [D\mathring{\zeta}_\epsilon]^{-1}, \quad \mathring{J}_\epsilon = \det D\mathring{\zeta}_\epsilon, \quad \mathring{a}_\epsilon = \mathring{J}_\epsilon \mathring{A}_\epsilon, \ [\mathring{g}_\epsilon]_{\alpha\beta} = \mathring{\zeta}_\epsilon,_\alpha \cdot \mathring{\zeta}_\epsilon,_\beta \,, \text{ and } \sqrt{\mathring{g}_\epsilon}\mathring{n}_\epsilon = [\mathring{a}_\epsilon]^T N.$$

We recall that the convolution operator Λ_μ for $\mu > 0$ is defined via Section 5.1.

Definition 5.4 (The μ-problem)**.** *Given $\kappa > 0$ and $\epsilon > 0$, for $\mu > 0$ we define $\mathring{v}$ as the solution of the nonlinear heat-type system*

$$\mathring{v}_t - \mathring{\varrho}[\mathring{A}_\epsilon]^j_r \big([\mathring{A}_\epsilon]^k_r \mathring{v},_k\big),_j = \mathring{\mathcal{K}} \qquad \textit{in } \Omega \times (0, T_\kappa(\epsilon\mu)], \tag{5.52a}$$

$$\mathring{\varrho} N^j [\mathring{A}_\epsilon]^j_r [\mathring{A}_\epsilon]^s_r \mathring{v},_s = \mathring{h}^\mu + c^\mu(t) \qquad \textit{on } \Gamma \times (0, T_\kappa(\epsilon\mu)], \tag{5.52b}$$

$$(\mathring{\zeta}_\epsilon, \mathring{v})|_{t=0} = (e, u_0) \qquad \textit{on } \Omega. \tag{5.52c}$$

The bounded, nonnegative function $\mathring{\varrho}$ is defined as

$$\mathring{\varrho} = \kappa\rho_0 \mathring{J}_\epsilon. \tag{5.53}$$

The vector field $\mathring{\mathcal{K}}$ appearing in the right-hand side of (5.52a) *is defined as*

$$\mathring{\mathcal{K}} = \mathring{\varrho}\, \operatorname{curl}_{\mathring{\zeta}_\epsilon} \Big(\operatorname{curl} u_0 + \varepsilon_{\cdot ji} \int_0^t \partial_t [\mathring{A}_\epsilon]^s_j \mathring{v}^i,_s\Big) + \operatorname{div}_{\mathring{\zeta}_\epsilon} \mathring{v} [\mathring{A}_\epsilon]^k_\cdot \mathring{\varrho},_k - 2[\mathring{A}_\epsilon]^k_\cdot (\rho_0 \mathring{J}_\epsilon^{-1}),_k \,. \tag{5.54}$$

The vector field $\mathring{h}^\mu$ appearing in the right-hand side of (5.52b) *is given by*

$$\mathring{h}^\mu = \mathring{h}_{\text{curl}} + \mathring{h}_{\text{div}} + \sum_{l=1}^K \sqrt{\xi_l}\Big(\Lambda_\mu\Big[\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l} \Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v},_{\alpha\beta} \cdot \mathring{n}_\epsilon) \circ \theta_l\big]\Big]\Big) \circ \theta_l^{-1} \mathring{n}_\epsilon + \mathring{\varrho}\Big[\mathring{b}^\mu_{\text{curl}} + \mathring{b}^\mu_{\text{div}}\Big], \tag{5.55}$$

where $\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l}$ appearing in the right-hand side of (5.55) *is defined as*

$$\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l} = \kappa \frac{\sqrt{\mathring{g}_\epsilon}}{\rho_0} \mathring{g}^{\alpha\beta}_\epsilon \circ \theta_l, \tag{5.56}$$

and $\mathring{h}_{\text{curl}}$, $\mathring{h}_{\text{div}}$, $\mathring{b}^{\mu}_{\text{curl}}$, $\mathring{b}^{\mu}_{\text{div}}$ appearing in the right-hand side of (5.55) *are defined as*

$$\mathring{h}_{\text{curl}} = \mathring{\varrho}\sqrt{\mathring{g}_\epsilon}\mathring{J}_\epsilon^{-1}\big(\operatorname{curl} u_0 + \varepsilon_{\cdot ji}\int_0^t \partial_t[\mathring{A}_\epsilon]^s_j \mathring{v}^i,_s\big) \times \mathring{n}_\epsilon, \tag{5.57a}$$

$$\mathring{h}_{\text{div}} = \frac{\sqrt{\mathring{g}_\epsilon}}{\rho_0}\Big[\rho_0^2\mathring{J}_\epsilon^{-2} - \beta_\epsilon(t) + \sigma\mathring{g}_\epsilon^{\alpha\beta}\mathring{\zeta}_\epsilon,_{\alpha\beta}\cdot\mathring{n}_\epsilon\Big]\mathring{n}_\epsilon, \tag{5.57b}$$

$$\mathring{b}^{\mu}_{\text{curl}} = \frac{\sqrt{\mathring{g}_\epsilon}}{\mathring{J}_\epsilon}\Big[\sum_{l=1}^K \mathring{g}_\epsilon^{\alpha\beta}\sqrt{\xi_l}\big(\Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}\cdot\mathring{n}_\epsilon),_\beta \circ\theta_l\big]\big)\circ\theta_l^{-1} + \mathring{v}^\gamma\mathring{g}_\epsilon^{\gamma\delta}\mathring{g}_\epsilon^{\alpha\beta}\mathring{\zeta}_\epsilon,_{\delta\beta}\cdot\mathring{n}_\epsilon\Big]\mathring{\zeta}_\epsilon,_\alpha\,, \tag{5.57c}$$

$$\mathring{b}^{\mu}_{\text{div}} = -\sum_{l=1}^K\sqrt{\xi_l}\Lambda_\mu\Big(\sqrt{\xi_l}\frac{\sqrt{\mathring{g}_\epsilon}}{\mathring{J}_\epsilon}\big[\mathring{g}_\epsilon^{\alpha\beta}\mathring{v}^\alpha,_\beta + \mathring{v}^\alpha(\sqrt{\mathring{g}_\epsilon}\mathring{g}_\epsilon^{\alpha\beta}),_\beta + \mathring{v}^3 H(\mathring{\zeta}_\epsilon)\big]\circ\theta_l\Big)\circ\theta_l^{-1}\mathring{n}_\epsilon. \tag{5.57d}$$

The vector field $c^\mu(t)$ appearing in the right-hand side of (5.52b) *is defined as*

$$c^\mu(t) = \sum_{a=0}^2\frac{t^a}{a!}\partial_t^a\big[\mathring{\varrho}N^j[\mathring{A}_\epsilon]^j_r[\mathring{A}_\epsilon]^s_r\mathring{v},_s - \mathring{h}^\mu\big]|_{t=0}. \tag{5.58}$$

Remark 18. The fixed-point solution to the μ-problem (5.52) is established in Appendix B.

Remark 19. For any h defined on Γ, we have that $h = \sum_{l=1}^K \xi_l h = \sum_{l=1}^K \sqrt{\xi_l}\big([\sqrt{\xi_l}h\circ\theta_l]\circ\theta_l^{-1}\big)$.

For $\mu > 0$, we define the following higher-order energy function:

$$E^\mu(t) = 1 + \|\mathring{v}_{tt}(t)\|_0^2 + \sum_{a=0}^2\int_0^t\|\partial_t^a\mathring{v}\|^2_{5-2a} + \sum_{a=0}^2\sum_{l=1}^K\int_0^t|\Lambda_\mu[(\sqrt{\xi_l}\partial_t^a\mathring{v}\cdot\mathring{n}_\epsilon)\circ\theta_l]|^2_{5-2a,\mathcal{D}_l}. \tag{5.59}$$

Definition 5.5 (Notational convention for constants depending on $1/\delta\kappa\epsilon > 0$)**.** *We let $\ddot{\mathcal{P}}$ denote a generic polynomial with constant and coefficients depending on $1/\delta\kappa\epsilon > 0$.*

We define the constant $\ddot{\mathcal{N}}_0 > 0$ by

$$\ddot{\mathcal{N}}_0 = \ddot{\mathcal{P}}(\|u_0\|_{100}, \|\rho_0\|_{100}). \tag{5.60}$$

We let $\ddot{\mathcal{R}}$ denote generic lower-order terms satisfying

$$\int_0^T\ddot{\mathcal{R}} \le \ddot{\mathcal{N}}_0 + \delta\sup_{t\in[0,T]}E^\mu(t) + T\,\ddot{\mathcal{P}}(\sup_{t\in[0,T]}E^\mu(t)).$$

Lemma 5.4 (A priori estimates for the μ-problem)**.** *We let $\mathring{v}$ solve the μ-problem* (5.52) *on a time-interval $[0,T]$ for some $T = T_\kappa(\epsilon\mu) > 0$. Then independent of μ,*

$$\sup_{t\in[0,T]}E^\mu(t) \le \int_0^T\ddot{\mathcal{R}}. \tag{5.61}$$

We will establish Lemma 5.4 in the following four steps:

Step 1: The μ-independent estimates for $\mathring{v}_{tt}$. Testing two time-derivatives of (5.52a) against $\mathring{v}_{tt}$ in the $L^2(\Omega)$-inner product and integrating by parts in the interior integral

$-\int_\Omega \partial_t^2 \big(\mathring{\varrho}[\mathring{A}_\epsilon]^j_r \big([\mathring{A}_\epsilon]^k_r \mathring{v},_k\big),_j\big)\mathring{v}_{tt}$ yields

$$\frac{1}{2}\frac{d}{dt}\int_\Omega |\mathring{v}_{tt}|^2 + \int_\Omega \mathring{\varrho}|[\mathring{A}_\epsilon]^k_\cdot \mathring{v}_{tt},_k|^2 - \underbrace{\int_\Gamma [\mathring{h}^\mu + c^\mu(t)]_{tt}\mathring{v}_{tt}}_{\mathfrak{i}}$$

$$= \underbrace{\int_\Omega \partial_t^2\big(\mathring{\mathcal{K}} - (\mathring{\varrho}[\mathring{A}_\epsilon]^j_r),_j\, [\mathring{A}_\epsilon]^k_r \mathring{v},_k\big)\mathring{v}_{tt}}_{\ddot{\mathcal{R}}} + \ddot{\mathcal{R}}. \tag{5.62}$$

The definition (5.55) of the vector field $\mathring{h}^\mu$ provides that

$$\mathfrak{i} = \underbrace{-\sum_{l=1}^K \int_{\mathcal{D}_l} \Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt},_{\alpha\beta} \cdot \mathring{n}_\epsilon)\circ\theta_l\big]\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l}\Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l\big]}_{\mathfrak{i}'} - \underbrace{\int_\Gamma (\mathring{\varrho}[\mathring{b}^\mu_{\text{curl}} + \mathring{b}^\mu_{\text{div}}])_{tt}\mathring{v}_{tt}}_{\mathfrak{j}} + \ddot{\mathcal{R}}.$$

We integrate by parts with respect to $\bar{\partial}_\alpha$ in $\mathfrak{i}'$ to find that

$$\begin{aligned}
\mathfrak{i}' = & \sum_{l=1}^K \int_{\mathcal{D}_l} \Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt},_\beta \cdot\mathring{n}_\epsilon)\circ\theta_l\big]\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l}\Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt},_\alpha \cdot\mathring{n}_\epsilon)\circ\theta_l\big] \\
& + \underbrace{\sum_{l=1}^K \int_{\mathcal{D}_l} \Lambda_\mu\big[(\mathring{v}_{tt},_\beta \cdot(\mathring{n}_\epsilon\sqrt{\xi_l}),_\alpha)\circ\theta_l\big]\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l}\Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l\big]}_{\mathfrak{i}'_A} \\
& + \underbrace{\sum_{l=1}^K \int_{\mathcal{D}_l} \Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt},_\beta \cdot\mathring{n}_\epsilon)\circ\theta_l\big]\bar{\partial}_\alpha\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l}\Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l\big]}_{\mathfrak{i}'_B} \\
& + \underbrace{\sum_{l=1}^K \int_{\mathcal{D}_l} \Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt},_\beta \cdot\mathring{n}_\epsilon)\circ\theta_l\big]\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l}\Lambda_\mu\big[(\mathring{v}_{tt}\cdot(\mathring{n}_\epsilon\sqrt{\xi_l}),_\alpha)\circ\theta_l\big]}_{\mathfrak{i}'_C}.
\end{aligned}$$

We employ an $H^{-0.5}(\mathcal{D}_l)$-duality pairing in the integrals $\mathfrak{i}'_A$, $\mathfrak{i}'_B$ and $\mathfrak{i}'_C$. For example,

$$\begin{aligned}
\mathfrak{i}'_A &\le C|\bar{\partial}\mathring{v}_{tt}\circ\theta_l|_{-0.5,\mathcal{D}_l}|\Lambda_\mu[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l]|_{0.5,\mathcal{D}_l} \\
&\le \delta\|\mathring{v}_{tt}\|_1^2 + C_\delta|\Lambda_\mu[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l]|^2_{0.5,\mathcal{D}_l}.
\end{aligned}$$

Thanks to Lemma 2.1, $|\Lambda_\mu[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l]|^2_{0.5,\mathcal{D}_l} \le C|\mathring{v}_{tt}|_0|\Lambda_\mu[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l]|_{1,\mathcal{D}_l} \le \ddot{\mathcal{R}} + \delta|\Lambda_\mu\big(\mathring{v}_{tt}\cdot\mathring{n}_\epsilon\big)|^2_{1,\mathcal{D}_l}$. Thus, we infer that

$$\mathfrak{i} = \sum_{l=1}^K \int_{\mathcal{D}_l} \Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon),_\beta\circ\theta_l\big]\mathring{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l}\Lambda_\mu\big[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon),_\alpha\circ\theta_l\big] + \ddot{\mathcal{R}},$$

where we have used the definitions (5.57c) and (5.57d) in analyzing $\mathfrak{j}$. We infer from the time integral of (5.62) that

$$\sup_{t\in[0,T]}\|\mathring{v}_{tt}(t)\|_0^2 + \int_0^T \|\mathring{v}_{tt}\|_1^2 + \sum_{l=1}^K \int_0^t |\Lambda_\mu[(\sqrt{\xi_l}\mathring{v}_{tt}\cdot\mathring{n}_\epsilon)\circ\theta_l]|^2_{1,\mathcal{D}_l} \le \int_0^T \ddot{\mathcal{R}}. \tag{5.63}$$

Step 2: The μ-independent estimates for $\mathring{v}_t$. Similar to Step 1, testing the action of $\bar{\partial}^2\partial_t$ in the equations (5.52a) against $\bar{\partial}^2\mathring{v}_t$ in the $L^2(\Omega)$-inner product yields

$$\sup_{t\in[0,T]}\|\bar{\partial}^2\mathring{v}_t(t)\|_0^2+\int_0^T\|\bar{\partial}^2\mathring{v}_t\|_1^2+\sum_{l=1}^K\int_0^t|\Lambda_\mu[(\sqrt{\xi_l}\mathring{v}_t\cdot\mathring{n}_\epsilon)\circ\theta_l]|_{3,\mathcal{D}_l}^2\le\int_0^T\ddot{\mathcal{R}}. \tag{5.64}$$

The inequality (5.64) provides that $\int_0^T|\mathring{v}_t|_{2.5}^2\le\int_0^T\ddot{\mathcal{R}}$. We infer from Step 2 of the proof of Lemma B.2 that by viewing a time-derivative of the equations (5.52a) as an elliptic Dirichlet problem for $\mathring{v}_t$ that

$$\int_0^T\|\mathring{v}_t\|_3^2\le\int_0^T\ddot{\mathcal{R}}. \tag{5.65}$$

Step 3: The μ-independent estimates for $\mathring{v}$. Repeating Step 1, we test four tangential-derivatives of (5.52a) against $\bar{\partial}^4\mathring{v}$ in the $L^2(\Omega)$-inner product and integrate by parts in the integral $-\int_\Omega\bar{\partial}^4\big(\mathring{\varrho}[\mathring{A}_\epsilon]_r^j\big([\mathring{A}_\epsilon]_r^k\mathring{v},_k\big),_j\big)\bar{\partial}^4\mathring{v}$ to find that

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\int_\Omega|\bar{\partial}^4\mathring{v}|^2&+\sum_{l=1}^K\int_{\mathcal{D}_l}\Lambda_\mu\big[\bar{\partial}^4(\sqrt{\xi_l}\mathring{v}\cdot\mathring{n}_\epsilon),_\beta\circ\theta_l\big]\mathring{\mathfrak{g}}_{\epsilon,l}^{\alpha\beta}\Lambda_\mu\big[\bar{\partial}^4(\sqrt{\xi_l}\mathring{v}\cdot\mathring{n}_\epsilon),_\alpha\circ\theta_l\big]\\&+\int_\Omega\mathring{\varrho}|[\mathring{A}_\epsilon]_\cdot^k\bar{\partial}^4\mathring{v},_k|^2=\underbrace{\int_\Gamma[\mathring{\varrho}\sqrt{\mathring{g}_\epsilon}\mathring{J}_\epsilon^{-1}\big(\varepsilon_{\cdot ji}\int_0^t\partial_t[\mathring{A}_\epsilon]_j^s\bar{\partial}^4\mathring{v}^i\big),_s\times\mathring{n}_\epsilon]\cdot\bar{\partial}^4\mathring{v}}_{\ddot{\mathcal{R}}}\\&\qquad+\underbrace{\int_\Omega\bar{\partial}^4\Big[\mathring{\mathcal{K}}-(\mathring{\varrho}[\mathring{A}_\epsilon]_r^j),_j[\mathring{A}_\epsilon]_r^k\mathring{v},_k\Big]\bar{\partial}^4\mathring{v}}_{\mathcal{I}}+\ddot{\mathcal{R}}.\end{aligned} \tag{5.66}$$

We use an $H^{-0.5}(\mathcal{D}_l)$-duality pairing in analyzing the first term appearing in the right-hand side of (5.66) and conclude a good estimate thanks to the time-integral. Tangentially integrating by parts in $\mathcal{I}$, we find that

$$\int_0^T\mathcal{I}\le\int_0^T\ddot{\mathcal{R}}.$$

Hence, the time-integral of (5.66) yields

$$\sup_{t\in[0,T]}\|\bar{\partial}^4\mathring{v}(t)\|_0^2+\int_0^T\|\bar{\partial}^4\mathring{v}\|_1^2+\sum_{l=1}^K\int_0^t|\Lambda_\mu[(\sqrt{\xi_l}\mathring{v}\cdot\mathring{n}_\epsilon)\circ\theta_l]|_{5,\mathcal{D}_l}^2\le\int_0^T\ddot{\mathcal{R}}. \tag{5.67}$$

The inequality (5.67) yields

$$\int_0^T|\mathring{v}|_{4.5}^2\le\int_0^T\ddot{\mathcal{R}}. \tag{5.68}$$

Viewing the equations (5.52a) as an elliptic Dirichlet problem for $\mathring{v}$, we infer from elliptic regularity and the inequalities (5.65) and (5.68) that

$$\int_0^T\|\mathring{v}\|_5^2\le\int_0^T\ddot{\mathcal{R}}. \tag{5.69}$$

Step 4: Concluding the proof of Lemma 5.4. The sum of the inequalities (5.63), (5.65), (5.67) and (5.69) completes the proof of Lemma 5.4. Taking δ sufficiently small in the inequality (5.61) yields a polynomial-type inequality of the form (2.19). Hence, for sufficiently small $T = T_\kappa(\epsilon) > 0$ and independently of $\mu > 0$,

$$\sup_{t\in[0,T]} E^\mu(t) \le 2\ddot{\mathcal{N}}_0, \tag{5.70}$$

where the higher-order energy function $E^\mu(t)$ is defined in (5.59).

5.5. **The proof of Proposition 5.4.** Proposition B.1 establishes the existence and uniqueness of a solution $\mathring{v}$ to the μ-problem (5.52). Given the μ-independent estimate (5.70), standard compactness arguments provide for the existence of the strong convergence, as μ tends to zero,

$$\begin{aligned} \mathring{\check{\zeta}}_\epsilon &\to \check{\zeta}_\epsilon \text{ in } L^2(0,T;H^4(\Omega)), & \mathring{\check{\mathcal{K}}} &\to \check{\mathcal{K}} \quad \text{in } L^2(0,T;H^2(\Omega)),\\ \mathring{v} &\to \check{v} \text{ in } L^2(0,T;H^4(\Omega)), & \mathring{h}^\mu &\to \check{h} \quad \text{in } L^2(0,T;H^{1.5}(\Gamma)),\\ \mathring{v}_t &\to \check{v}_t \text{ in } L^2(0,T;H^2(\Omega)), & c^\mu(t) &\to c(t) \text{ in } L^2(0,T;H^{1.5}(\Gamma)). \end{aligned}$$

The definitions (5.54), (5.55), (5.58) respectively define $\mathring{\check{\mathcal{K}}}$, $\mathring{h}^\mu$, $c^\mu(t)$. The limiting vectors $\check{\mathcal{K}}$, $\check{h}$ and $c(t)$ are respectively defined by (5.32), (5.46) and (5.50). Letting $\phi \in L^2(0,T;H^1(\Omega))$, we have that the variational form of the μ-problem (5.52) is

$$\int_0^T\int_\Omega \mathring{v}_t\cdot\phi + \int_0^T\int_\Omega [\mathring{A}_\epsilon]^k_r \mathring{v},_k \left(\mathring{\varrho}[\mathring{A}_\epsilon]^j_r\phi\right),_j = \int_0^T\int_\Omega \mathring{\check{\mathcal{K}}}\cdot\phi + \int_0^T\int_\Gamma [\mathring{h}^\mu + c^\mu(t)]\cdot\phi.$$

We infer from the strong convergence of the sequences $(\mathring{\check{\zeta}}_\epsilon, \mathring{v}, \mathring{v}_t, \mathring{\check{\mathcal{K}}}, \mathring{h}^\mu, c^\mu(t))$ that the limit $(\check{\zeta}_\epsilon, \check{v}, \check{v}_t, \check{\mathcal{K}}, \check{h}, c(t))$ satisfies

$$\int_0^T\int_\Omega \check{v}_t\cdot\phi + \int_0^T\int_\Omega [\check{A}_\epsilon]^k_r \check{v},_k \left(\varrho[\check{A}_\epsilon]^j_r\phi\right),_j = \int_0^T\int_\Omega \check{\mathcal{K}}\cdot\phi + \int_0^T\int_\Gamma [\check{h} + c(t)]\cdot\phi.$$

Thus, $\check{v}$ is a solution of the nonlinear heat-type $\kappa\epsilon$-problem (5.49) on a time-interval $[0,T]$ for some $T = T_\kappa(\epsilon) > 0$. Standard arguments provide that $\check{\zeta}_\epsilon(0) = e$ and $(\check{v}, \check{v}_t, \dots, \check{v}_{tttt})|_{t=0} = (u_0, \mathbf{v}_1, \dots, \mathbf{v}_4)$. Furthermore, according to the inequality (5.70),

$$\sup_{t\in[0,T]} \|\check{v}_{tt}(t)\|_0^2 + \sum_{a=0}^2 \int_0^T \|\partial_t^a \check{v}\|^2_{5-2a} + \sum_{a=0}^2 \int_0^T |\partial_t^a \check{v}\cdot\check{n}_\epsilon|^2_{5-2a} \le 2\ddot{\mathcal{N}}_0.$$

By the higher-order regularity stated in Proposition B.1, we infer that

$$\sup_{t\in[0,T]} \|\check{v}_{tttt}(t)\|_0^2 + \sum_{a=0}^4 \int_0^T \|\partial_t^a \check{v}\|^2_{9-2a} + \sum_{a=0}^4 \int_0^T |\partial_t^a \check{v}\cdot\check{n}_\epsilon|^2_{9-2a} < \infty. \tag{5.71}$$

5.6. **The proof of Theorem 5.2.** By Lemma C.1, the heat-type $\kappa\epsilon$-problem (5.49) is equivalent to the $\kappa\epsilon$-problem (5.6). Hence, Proposition 5.4 establishes the existence and uniqueness of a solution to the $\kappa\epsilon$-problem on a time-interval $[0,T]$ for some $T = T_\kappa(\epsilon) > 0$ verifying $(\check{v}, \check{v}_t, \dots, \check{v}_{tttt})|_{t=0} = (u_0, \mathbf{v}_1, \dots, \mathbf{v}_4)$. The inequality (5.71) establishes the inequality (5.26).

5.7. **The proof of Theorem 5.1.**

A unique solution to the $\kappa\epsilon$-problem (5.6) exists by Theorem 5.2.

Taking δ sufficiently small in the inequality (5.13) yields a polynomial-type inequality of the form (2.19). Hence, for sufficiently small $T = T_\kappa > 0$ and independently of $\epsilon > 0$,

$$\sup_{t\in[0,T]} E^\epsilon(t) \le 2\dot{\mathcal{N}}_0, \tag{5.72}$$

where the higher-order energy function $E^\epsilon(t)$ is defined in (5.12).

For $\phi \in L^2(0,T;H^1(\Omega))$ such that $\phi\cdot\check{n}_\epsilon \in L^2(0,T;H^1(\Gamma))$, the variational equation for the $\kappa\epsilon$-problem (5.6) is

$$\int_0^T\int_\Omega \rho_0\check{v}_t\cdot\phi - \int_0^T\int_\Omega \rho_0^2\check{J}_\epsilon^{-2}[\check{a}_\epsilon]_i^k\phi^i,_k + \kappa\int_0^T\int_\Omega \rho_0\check{\mathcal{J}}_t[\check{a}_\epsilon]_i^k(\rho_0\check{J}_\epsilon^{-1}\phi^i),_k$$
$$+\int_0^T\int_\Gamma \beta_\epsilon(t)\sqrt{\check{g}_\epsilon}\phi\cdot\check{n}_\epsilon + \int_0^T\int_\Gamma [\sigma\check{\zeta}_\epsilon,_\beta + \kappa\check{v},_\beta]^i(\check{n}_\epsilon^i\sqrt{\check{g}_\epsilon}\check{g}_\epsilon^{\alpha\beta}\phi\cdot\check{n}_\epsilon),_\alpha = 0. \tag{5.73}$$

We infer from Section 5.5 and the pointwise convergence $\beta_\epsilon(t)\to\beta(t)$ that the variational equation (5.73) converges to the variational equation (3.4) as ϵ tends to zero. Hence, the $\epsilon = 0$ limit $\tilde{v}$ of the solutions $\check{v}$ to the $\kappa\epsilon$-problem (5.6) solves the κ-problem (3.1). By Section 5.2.1, the solution $\tilde{v}$ of the κ-problem verifies $(\tilde{v},\tilde{v}_t,\dots,\tilde{v}_{tttt})|_{t=0} = (u_0,\mathrm{v}_1,\dots,\mathrm{v}_4)$. According to (5.72),

$$\sup_{t\in[0,T]}\|\tilde{v}_{tt}(t)\|_0^2 + \int_0^T|\tilde{v}_{tt}\cdot\tilde{n}|_1^2 + \sum_{a=0}^2\int_0^T\|\partial_t^a\tilde{v}\|_{5-2a}^2 \le 2\dot{\mathcal{N}}_0.$$

It follows from the proof of Lemma 5.1 that by including two more time-derivatives in the definition (5.12) of the energy function $E^\epsilon(t)$, the solution $\tilde{v}$ of the κ-problem (3.1) satisfies

$$\sup_{t\in[0,T]}\|\tilde{v}_{tttt}(t)\|_0^2 + \int_0^T|\tilde{v}_{tttt}\cdot\tilde{n}|_1^2 + \sum_{a=0}^4\int_0^T\|\partial_t^a\tilde{v}\|_{9-2a}^2 < \infty.$$

This establishes the inequality (5.1).

6. WELL-POSEDNESS OF THE SURFACE TENSION PROBLEM (1.7)

In this section, we prove Theorem 1.1 via the κ-independent a priori estimates of Section 4.

6.1. **Existence.**

We obtain a solution v to surface tension problem (1.7) in the limit of $\tilde{v}$ as the parabolic parameter κ tends to zero. According to Remark 9, $\beta(t) = \beta$ in the $\kappa = 0$ limit. Letting $\kappa = 0$ in Section 4, we therefore conclude that the right-hand side of the inequality (4.41) depends only on $M_0 = P(E(0))$. That is, for sufficiently small $T > 0$ the energy function $E(t)$ defined in (1.9) satisfies

$$\sup_{t\in[0,T]} E(t) \le 2M_0. \tag{6.1}$$

The assumption (4.3) on J remains valid by taking $T > 0$ even smaller if necessary. Hence,

$$f(t) = \rho_0 J^{-1}(t) \ge \tfrac{4}{3}\lambda.$$

Taking $T > 0$ even smaller if necessary, we ensure that $\rho(t) = f\circ\eta^{-1}(t)$ satisfies

$$\rho(t) \ge \lambda \ \text{ in } \overline{\Omega}(t).$$

Since $p(t) = \rho^2(t) - \beta > -\beta$ on $\Gamma(t)$, the boundary condition (1.1c) establishes that

$$\sigma H(t) > -\beta \ \text{ on } \Gamma(t).$$

6.2. **Optimal regularity for the initial data.** In order to obtain the $H^5(\Omega)$-regularity of our existence theory, we assumed that the given initial data is of C^∞-class in Section 3.1. In fact, by virtue of the estimate (6.1), it suffices for the regularity of the initial data to be such that $E(0) < \infty$.

6.3. **Uniqueness.** We define

$$\mathbf{E}(v,t) = 1 + \sum_{a=0}^{6} \|\partial_t^a \eta(t)\|^2_{6-a} + |\partial_t^4 v \cdot n(t)|_1^2 + \sum_{a=0}^{4} |\bar\partial^2 \partial_t^a \eta \cdot n(t)|^2_{4.5-a}.$$

We suppose that (η_1, v_1, f_1) and (η_2, v_2, f_2) are two solutions of the compressible surface tension problem (1.7) with $\mathbf{E}(v,0)$, for $v = v_1, v_2$, bounded by some $\mathbf{M}_0 > 0$.

Then by setting

$$\zeta = \eta_1 - \eta_2, \quad w = v_1 - v_2, \quad \varrho = [f_1]^2 - [f_2]^2,$$

we have that (ζ, w, ϱ) satisfies

$$\zeta = \int_0^t w \qquad \text{in } \Omega \times (0,T], \tag{6.2a}$$

$$\rho_0 w_t^i + [a_1]_i^k \varrho,_k = [a_2 - a_1]_i^k ([f_2]^2),_k \qquad \text{in } \Omega \times (0,T], \tag{6.2b}$$

$$\begin{aligned} \varrho = -\sigma [g_1]^{\alpha\beta} \zeta,_{\alpha\beta} \cdot n_1 - \sigma [g_1]^{\alpha\beta} \eta_2,_{\alpha\beta} \cdot n_1 & \\ + \sigma [g_2]^{\alpha\beta} \eta_2,_{\alpha\beta} \cdot n_2 & \qquad \text{on } \Gamma \times (0,T], \end{aligned} \tag{6.2c}$$

$$(\zeta, w, \varrho)|_{t=0} = (0,0,0) \qquad \text{on } \Omega. \tag{6.2d}$$

We will establish that $w = 0$. Setting

$$E^\zeta(t) = \sum_{a=0}^{5} \|\partial_t^a \zeta(t)\|^2_{5-a}, \tag{6.3}$$

we follow Section 4 with κ set to zero.

The Lagrangian curl operator $\operatorname{curl}_{\eta_1}$ applied to

$$w_t^i + 2[A_1]_i^k f_1,_k = 2[A_2]_i^k f_2,_k$$

provides the following vorticity equation for the difference $w = v_1 - v_2$:

$$\operatorname{curl}_{\eta_1} w_t = 2\varepsilon_{\cdot ji} [A_1]_j^s ([A_2]_i^k f_2,_k\,),_s\,. \tag{6.4}$$

The time integral of the surface tension problem satisfied by v_2 yields

$$v_2 - u_0 = -2 \int_0^t [A_2]_\cdot^k f_2,_k\,,$$

by which we infer that

$$\int_0^T \|D^2 \int_0^t [A_2]_\cdot^k f_2,_k\,\|_3^2 \leq T\,P(\sup_{t\in[0,T]} \|v_2(t)\|_5^2) \leq C\,T\,\mathbf{M}_0.$$

The curl-estimates for w therefore follow from the analysis proving Lemma 4.2 with the vorticity equation (6.4) replacing the homogeneous vorticity equation (2.13).

Repeating the energy estimate for the fourth time-differentiated problem in Proposition 4.1, the highest-order term of the interior forcing term $\int_0^t \int_\Omega \partial_t^4 [[a_2 - a_1]_i^k ([f_2]^2),_k\,] w^i_{tttt}$

obeys

$$\int_0^t \int_\Omega [a_2 - a_1]^k_i ([f_2]^2)_{tttt},_k \, w^i_{tttt} \le C \int_0^T \|Dv_{2ttt}\|_1 \|w_{tttt}\|_0 \le T\, P(\sup_{t\in[0,T]} E^\zeta(t)).$$

The boundary integral corresponding with

$$-\sigma [g_1]^{\alpha\beta} \eta_2,_{\alpha\beta} \cdot n_1 + \sigma [g_2]^{\alpha\beta} \eta_2,_{\alpha\beta} \cdot n_2 = -\sigma [g_1]^{\alpha\beta} \eta_2,_{\alpha\beta} \cdot [n_1 - n_2] - \sigma [g_1 - g_2]^{\alpha\beta} \eta_2,_{\alpha\beta} \cdot n_2$$

of (6.2c) is similarly bounded.

We notice that with

$$\int_\Omega \rho_0^2 (J_1)^{-3} [a_1]^q_p \, {v_1}_{ttt}{}^p,_q \, [a_1]^s_r w^r_{tttt},_s = \int_\Omega \rho_0^2 (J_1)^{-3} [a_1]^q_p \, w^p_{ttt},_q \, [a_1]^s_r w^r_{tttt},_s + \int_\Omega \rho_0^2 (J_1)^{-3} [a_1]^q_p \, {v_2}_{ttt}{}^p,_q \, [a_1]^s_r w^r_{tttt},_s \,,$$

we preserve an energy estimate for $\|[a_1]^s_r w^r_{ttt},_s\|^2_0$ in Proposition 4.1.

Following the arguments in Step 3 of Section 4 provides control of the divergence and normal trace of the functions of $E^\zeta(t)$. Proposition 2.1 and the initial condition (6.2d) imply that

$$\sup_{t\in[0,T]} E^\zeta(t) \le T\, P(\sup_{t\in[0,T]} E^\zeta(t)),$$

for the higher-order energy function $E^\zeta(t)$ defined in (6.3). Using the polynomial-type inequality (2.19), we infer that $w = 0$ as desired.

7. The asymptotic limit as surface tension tends to zero

In this section, we establish an existence theory for the zero surface tension limit of (1.7) via a priori estimates that are independent of the surface tension parameter $\sigma > 0$. This asymptotic limit holds whenever the initial data satisfies the Taylor sign condition (1.15), and provides the following:

$$0 < \nu \int_\Gamma |\bar\partial^4 \eta \cdot n|^2 \le - \int_\Gamma \frac{1}{\sqrt{g}} N^j a^j_\ell a^k_\ell (\rho_0^2 J^{-2}),_k \, |\bar\partial^4 \eta \cdot n|^2.$$

We recall that according to Theorem 1.1, a solution to (1.7) satisfies

$$\sup_{t\in[0,T]} \sum_{a=0}^{5} \|\partial_t^a \eta(t)\|^2_{5-a} + |v_{ttt} \cdot n(t)|^2_1 + \sum_0^2 |\bar\partial^2 \partial_t^a v \cdot n(t)|^2_{2.5-a} \le C_\sigma, \tag{7.1}$$

for a finite bound $C_\sigma > 0$ depending on $1/\sigma$. The Taylor-sign-condition assumption of Theorem 1.2 provides for σ-independent a priori estimates under the higher-order energy function $E^\sigma(t)$ defined below in (7.5).

7.1. Assuming C^∞-class initial data.

In our construction of solutions to the zero surface tension limit of (1.7), we will assume that the initial data (ρ_0, u_0, Ω) is of C^∞-class and satisfy the conditions (1.13), (1.15) and (1.16), as in Appendix A. Later, in Section 7.4.2, we will recover the optimal regularity of the initial data stated in Theorem 1.2.

7.2. **The σ-problem.** For $\sigma > 0$ taken sufficiently small, solutions to the following problem are provided by Theorem 1.1. To indicate the dependence on the surface tension paramater σ of all the variables in the following problem, we place the symbol $\smile$ over each of the variables.

Definition 7.1 (The σ-problem)**.** *For $\sigma > 0$, we define $\breve{v}$ as the solution of*

$$\rho_0 \breve{v}_t^i + \breve{a}_i^k (\rho_0^2 \breve{J}^{-2}),_k = 0 \qquad \text{in } \Omega \times (0, T_\sigma], \tag{7.2a}$$

$$\rho_0^2 \breve{J}^{-2} = \beta_\sigma(t) - \sigma \breve{g}^{\alpha\beta} \breve{\eta},_{\alpha\beta} \cdot \breve{n} \qquad \text{on } \Gamma \times (0, T_\sigma], \tag{7.2b}$$

$$(\breve{\eta}, \breve{v})|_{t=0} = (e, u_0) \qquad \text{on } \Omega. \tag{7.2c}$$

The function $\beta_\sigma(t)$ appearing in the right-hand side of (7.2b) *is defined as*

$$\beta_\sigma(t) = \beta + \sum_{a=0}^{6} \frac{t^a}{a!} \partial_t^a [\rho_0^2 \breve{J}^{-2} - \beta + \sigma \breve{g}^{\alpha\beta} \breve{\eta},_{\alpha\beta} \cdot \breve{n}]\big|_{t=0}.$$

Remark 20. The initial data satisfy the compatibility conditions (1.16). Thus,

$$\beta_\sigma(t) = \beta + \sigma \sum_{a=0}^{6} \frac{t^a}{a!} \partial_t^a [\breve{g}^{\alpha\beta} \breve{\eta},_{\alpha\beta} \cdot \breve{n}]\big|_{t=0}. \tag{7.3}$$

The $\sigma = 0$ formal limit of $\beta_\sigma(t)$ is β. It follows that the σ-problem (7.2) is asymptotically consistent with the zero surface tension limit of (1.7).

Remark 21. We use (7.2b) to compute the following identities: for $a = 0, \dots, 6$,

$$\partial_t^a [\rho_0^2 \breve{J}^{-2}]|_{t=0} = \partial_t^a \beta_\sigma(t)|_{t=0} - \sigma \partial_t^a [\breve{g}^{\alpha\beta} \breve{\eta},_{\alpha\beta} \cdot \breve{n}]|_{t=0}. \tag{7.4}$$

7.3. **The a priori estimates for the σ-problem.** For $\sigma > 0$, we define

$$\begin{aligned} E^\sigma(t) = 1 &+ \sum_{a=0}^{7} \|\partial_t^a \breve{\eta}(t)\|_{4.5-\frac{1}{2}a}^2 + \sum_{a=0}^{5} \|\partial_t^a \breve{J}(t)\|_{4.5-\frac{1}{2}a}^2 + \|\partial_t^6 \breve{J}(t)\|_1^2 \\ &+ \sum_{a=0}^{5} \|\sqrt{\sigma} \partial_t^a \breve{\eta}(t)\|_{5.5-\frac{1}{2}a}^2 + \|\sqrt{\sigma} \partial_t^5 \breve{v}(t)\|_2^2 + \sum_{a=0}^{1} |\sqrt{\sigma} \partial_t^{5+a} \breve{v} \cdot \breve{n}(t)|_{2-a}^2 \\ &+ \sum_{a=0}^{3} \|\sigma \partial_t^a \breve{\eta}(t)\|_{6.5-\frac{1}{2}a}^2 + \|\sigma \breve{v}_{ttt}(t)\|_4^2 + \sum_{a=0}^{1} |\sigma \partial_t^{3+a} \breve{v} \cdot \breve{n}(t)|_{4-\frac{1}{2}a}^2 + |\sigma \partial_t^5 \breve{v} \cdot \breve{n}(t)|_{2.5}^2. \end{aligned} \tag{7.5}$$

We will allow constants to depend on $1/\delta > 0$:

Definition 7.2 (Notational convention for constants depending on $1/\delta > 0$)**.** *We let $\dddot{\mathcal{P}}$ denote a generic polynomial with constant and coefficients depending on $1/\delta > 0$.*

We define the constant $\dddot{\mathcal{N}}_0 > 0$ by

$$\dddot{\mathcal{N}}_0 = \dddot{\mathcal{P}}(\|u_0\|_{100}, \|\rho_0\|_{100}). \tag{7.6}$$

We let $\dddot{\mathcal{R}}$ denote generic lower-order terms satisfying

$$\int_0^T \dddot{\mathcal{R}} \le \dddot{\mathcal{N}}_0 + \delta \sup_{t \in [0,T]} E^\sigma(t) + T\, \dddot{\mathcal{P}}(\sup_{t \in [0,T]} E^\sigma(t)).$$

We infer from the estimates (4.4) and (7.1) that for $T > 0$ taken sufficiently small,

$$\frac{1}{2} \le \breve{J} \le \frac{3}{2} \quad \text{for all } t \in [0, T] \text{ and } x \in \Omega. \tag{7.7a}$$

Since the initial data satisfy the Taylor sign condition (1.15), we also assume that

$$0 < \nu \le -\frac{1}{\sqrt{\breve{g}}} N^j \breve{a}^j_\ell \breve{a}^k_\ell (\rho_0^2 \breve{J}^{-2}),_k \quad \text{for all } t \in [0,T] \text{ and } x \in \Gamma. \tag{7.7b}$$

Lemma 7.1 (A priori estimates for the σ-problem)**.** *We let* $\breve{v}$ *solve the* σ*-problem* (7.2) *on a time-interval* $[0,T]$*, for some* $T = T_\sigma > 0$*. Then independent of* $1 >> \sigma > 0$*,*

$$\sup_{t\in[0,T]} E^\sigma(t) \le \int_0^T \dddot{\mathcal{R}}. \tag{7.8}$$

We will establish Lemma 7.1 in the following nine steps:

Step 1: The σ-independent curl-estimates. We infer the following lemma from the arguments proving Lemma 4.2.

Lemma 7.2 (The σ-independent curl-estimates)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{7} \| \operatorname{curl} \partial_t^a \breve{\eta}(t)\|^2_{3.5-\frac{1}{2}a} + \sup_{t\in[0,T]} \sum_{a=0}^{1} \| \sqrt{\sigma} \operatorname{curl} \partial_t^{3+a} \breve{v}(t)\|^2_{2.5-\frac{1}{2}a} + \sup_{t\in[0,T]} \| \sqrt{\sigma} \operatorname{curl} \partial_t^5 \breve{v}(t)\|^2_1$$
$$+ \sup_{t\in[0,T]} \sum_{a=0}^{3} \| \sigma \operatorname{curl} \partial_t^a \breve{\eta}(t)\|^2_{5.5-\frac{1}{2}a} + \sup_{t\in[0,T]} \| \sigma \operatorname{curl} \partial_t^3 \breve{v}(t)\|^2_3 \le \int_0^T \dddot{\mathcal{R}}.$$

Step 2: The σ-independent estimates for $\partial_t^7 \breve{J}$, $\partial_t^6 \breve{J}$ and $\sqrt{\sigma}\partial_t^6 \breve{v} \cdot \breve{n}$. We recall that

$$\breve{f} = \rho_0 \breve{J}^{-1}$$

is the Lagrangian density. Using the identity $\breve{J}^{-1}\breve{J}_t = \operatorname{div}_{\breve{\eta}} \breve{v}$, we have that

$$\begin{aligned} \partial_t \breve{f}^2 &= -2\breve{f}^2 \operatorname{div}_{\breve{\eta}} \breve{v}, \\ \partial_t^2 \breve{f}^2 &= -2\breve{f}^2 \operatorname{div}_{\breve{\eta}} \breve{v}_t - 2(\breve{f}^2 \breve{A}^s_r)_t \breve{v}^r,_s . \end{aligned}$$

Letting the operator $-2[\breve{f}\breve{A}^j_i \partial_j]\breve{J}^{-1}$ act in the Euler equations (7.2a) yields

$$-2\breve{f}^2 \operatorname{div}_{\breve{\eta}} \breve{v}_t - 2\breve{f}\breve{A}^j_i \left[\breve{A}^k_i (\rho_0^2 \breve{J}^{-2}),_k \right],_j = \breve{v}^i_t \breve{A}^j_i \breve{f}^2,_j .$$

Using the Euler equations (7.2a) to write $\breve{v}^i_t = -\rho_0^{-1} \breve{a}^k_i \breve{f}^2,_k$, we infer that $\breve{f}^2$ satisfies

$$\partial_t^2 \breve{f}^2 - 2\breve{f}\breve{A}^j_i \left[\breve{A}^k_i \breve{f}^2,_k \right],_j = \underbrace{-\rho_0^{-1} \breve{a}^k_i \breve{f}^2,_k \breve{A}^j_i \breve{f}^2,_j - 2(\breve{f}^2 \breve{A}^j_i)_t \breve{v}^i,_j}_{\breve{F}} . \tag{7.9}$$

Since $\breve{F}$ scales like $D\breve{J} + D\breve{v}$, it follows that $\partial_t^6 \breve{F}$ is in $L^2(\Omega)$.

Similar to the tangential identity (4.14),

$$\rho_0 \breve{J}^{-1} \breve{v}_t \cdot \breve{\eta},_\gamma = \bar{\partial}_\gamma \left[\sigma \breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n} - \beta_\sigma(t) \right] \quad \text{on } \Gamma, \tag{7.10}$$

thanks to the Euler equations (7.2a) and the Laplace-Young boundary condition (7.2b).

Proposition 7.1 (Energy estimates for the action of ∂_t^6 in the wave-type equation (7.9))**.**

$$\sup_{t\in[0,T]} \|\partial_t^7 \breve{J}(t)\|^2_0 + \sup_{t\in[0,T]} \|\partial_t^6 \breve{J}(t)\|^2_1 + \sup_{t\in[0,T]} |\sqrt{\sigma}\partial_t^6 \breve{v} \cdot \breve{n}(t)|^2_1 \le \int_0^T \dddot{\mathcal{R}}.$$

Proof. Testing six time-derivatives of (7.9) against $\rho_0^{-2}\breve{J}^3\partial_t^7\breve{f}^2$ in the $L^2(\Omega)$-inner product, and integrating by parts with respect to ∂_j in the integral $-2\int_\Omega \rho_0^{-1}\breve{a}_i^j\partial_t^6\big[\breve{a}_i^k\breve{f}^2,_k\big],_j\partial_t^7\breve{f}^2$ yields

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\int_\Omega \rho_0^{-2}\breve{J}^3|\partial_t^7\breve{f}^2|^2 &+ 2\underbrace{\int_\Omega \rho_0^{-1}\partial_t^6\big[\breve{a}_i^k\breve{f}^2,_k\big]\breve{a}_i^j\partial_t^7\breve{f}^2,_j}_{\mathcal{I}} - 2\underbrace{\int_\Gamma \rho_0^{-1}\partial_t^6\big[\breve{a}_i^k\breve{f}^2,_k\big]\breve{a}_i^jN^j\partial_t^7\breve{f}^2}_{\mathfrak{i}}\\
&= \underbrace{\frac{1}{2}\int_\Omega(\rho_0^{-2}\breve{J}^3)_t|\partial_t^7\breve{f}^2|^2}_{\dddot{\mathcal{R}}} + \underbrace{2\sum_{l=1}^6 c_l\int_\Omega \breve{f}\breve{A}_i^j\big(\partial_t^l\breve{J}^{-1}\partial_t^{6-l}\big[\breve{a}_i^k\breve{f}^2,_k\big]\big),_j\rho_0^{-2}\breve{J}^3\partial_t^7\breve{f}^2}_{\dddot{\mathcal{R}}}\\
&+ \underbrace{2\sum_{l=1}^6 c_l\int_\Omega \partial_t^l(\breve{f}\breve{A}_i^j)\partial_t^{6-l}\big(\breve{A}_i^k\breve{f}^2,_k\big),_j\rho_0^{-2}\breve{J}^3\partial_t^7\breve{f}^2}_{\dddot{\mathcal{R}}} + \underbrace{\int_\Omega \partial_t^6F\rho_0^{-2}\breve{J}^3\partial_t^7\breve{f}^2}_{\dddot{\mathcal{R}}}\\
&+ \underbrace{2\int_\Omega \rho_0^{-1}\breve{a}_i^j(\breve{J}^{-1}),_j\partial_t^6[\tilde{a}_i^k\breve{f}^2,_k]\breve{J}\partial_t^7\breve{f}^2}_{\dddot{\mathcal{R}}}\,.\end{aligned} \tag{7.11}$$

We have used the Cauchy-Schwarz inequality to analyze all of the terms in the right-hand side of (7.11) except for the highest-order terms of $\int_\Omega \partial_t^4(\breve{f}\breve{A}_i^j)\partial_t^2\big(\breve{A}_i^k\breve{f}^2,_k\big),_j\rho_0^{-2}\breve{J}^3\partial_t^7\breve{f}^2$, where we have used an L^4–L^4–L^2 Hölder inequality.

Writing $\partial_t\breve{f}^2 = -2\rho_0^2\breve{J}^{-3}\breve{J}_t$, it follows that

$$\partial_t^7\breve{f}^2 = -2\rho_0^2\breve{J}^{-3}\partial_t^7\breve{J} - 2\sum_{l=1}^6 c_l\partial_t^l(\rho_0^2\breve{J}^{-3})\partial_t^{7-l}\breve{J}. \tag{7.12}$$

The identity (7.12) provides that the equation (7.11) multiplied by $\frac{1}{2}$ is equivalent to

$$\frac{d}{dt}\int_\Omega \rho_0^2\breve{J}^{-3}|\partial_t^7\breve{J}|^2 + \mathcal{I} - \mathfrak{i} = \dddot{\mathcal{R}}. \tag{7.13}$$

Analysis of $\mathcal{I}$ in (7.13). We equivalently write

$$\mathcal{I} = \underbrace{\int_\Omega \rho_0^{-1}\breve{a}_i^k\partial_t^6\breve{f}^2,_k\breve{a}_i^j\partial_t^7\breve{f}^2,_j}_{\mathcal{I}_a} + \sum_{l=0}^5 c_l\underbrace{\int_\Omega \rho_0^{-1}\partial_t^{6-l}\breve{a}_i^k\partial_t^l\breve{f}^2,_k\breve{a}_i^j\partial_t^7\breve{f}^2,_j}_{\mathcal{I}_{b,l}}\,.$$

We have that

$$\mathcal{I}_a = \frac{1}{2}\frac{d}{dt}\int_\Omega \rho_0^{-1}|\breve{a}_\cdot^k\partial_t^6\breve{f}^2,_k|^2 - \int_\Omega \rho_0^{-1}\breve{a}_i^k\partial_t^6\breve{f}^2,_k\partial_t\breve{a}_i^j\partial_t^6\breve{f}^2,_j\,.$$

Similar to (7.12), we have that $\partial_t^6\breve{f}^2$ is equal to $-2\breve{f}^2\breve{J}^{-1}\partial_t^6\breve{J}$ plus lower-order terms. Thus,

$$\mathcal{I}_a = 2\frac{d}{dt}\int_\Omega \rho_0^{-1}\breve{f}^4|\breve{A}_\cdot^k\partial_t^6\breve{J},_k|^2 + \dddot{\mathcal{R}}.$$

For $\int_0^T\mathcal{I}_{b,l}$, we integrate by parts with respect to a time-derivative of $\partial_t^7\breve{f}^2,_j$. For example,

$$\int_0^T\mathcal{I}_{b,0} = \int_\Omega \rho_0^{-1}\partial_t^6\breve{a}_i^k\breve{f}^2,_k\breve{a}_i^j\partial_t^6\breve{f}^2,_j\Big|_0^T - \int_0^T\int_\Omega \rho_0^{-1}\partial_t\big[\partial_t^6\breve{a}_i^k\breve{f}^2,_k\breve{a}_i^j\big]\partial_t^6\breve{f}^2,_j = \int_0^T\dddot{\mathcal{R}}.$$

Since $\partial_t^6 \breve{a}$ scales like $D\partial_t^5 \breve{v}$, the fundamental theorem of calculus ensures a good estimate for the term evaluated at time $t = T$. The terms $\int_0^T \mathcal{I}_{b,l}$, $l = 1, \ldots, 5$, are similarly analyzed:

$$\sum_{l=0}^{5} \mathcal{I}_{b,l} = \dddot{\mathcal{R}}.$$

This establishes that

$$\mathcal{I} = 2\frac{d}{dt}\int_\Omega \rho_0^{-1} \breve{f}^4 |\breve{A}^k_{\cdot} \partial_t^6 \breve{J},_k|^2 + \dddot{\mathcal{R}}. \tag{7.14}$$

Rewriting the boundary integral in (7.13)**.** We use the trace of the action of ∂_t^6 in the Euler equations (7.2a) to write $\mathfrak{i} = \int_\Gamma \partial_t^7 \breve{f}^2 \partial_t^7 \breve{v}^i \breve{a}_i^j N^j$, or equivalently,

$$\mathfrak{i} = \int_\Gamma \partial_t^7 \breve{f}^2 \sqrt{\breve{g}} \partial_t^7 \breve{v} \cdot \breve{n}.$$

Using the Laplace-Young boundary condition (7.2b), we find that

$$\begin{aligned}
\mathfrak{i} = {} & \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\alpha \cdot \breve{n}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\beta \cdot \breve{n} \\
& \underbrace{-\sigma \int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta} \partial_t^6 \breve{v},_\alpha \cdot \breve{n}_t\, \partial_t^6 \breve{v},_\beta \cdot \breve{n}}_{\mathfrak{j}_1} - \frac{1}{2} \underbrace{\int_\Gamma (\sqrt{\breve{g}}\breve{g}^{\alpha\beta})_t\, \sqrt{\sigma}\partial_t^6 \breve{v},_\alpha \cdot \breve{n}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\beta \cdot \breve{n}}_{\dddot{\mathcal{R}}} \\
& + \underbrace{\sigma \int_\Gamma \partial_t^6 \breve{v},_\beta \cdot \breve{n} \sqrt{\breve{g}}\breve{g}^{\alpha\beta} \partial_t^7 \breve{v} \cdot \breve{n},_\alpha + \sigma \int_\Gamma \partial_t^6 \breve{v},_\beta \cdot \breve{n},_\alpha \sqrt{\breve{g}}\breve{g}^{\alpha\beta} \partial_t^7 \breve{v} \cdot \breve{n}}_{\mathfrak{j}_2} \\
& + \underbrace{\sigma \int_\Gamma \partial_t^6 \breve{v},_\beta \cdot \breve{n} (\sqrt{\breve{g}}\breve{g}^{\alpha\beta}),_\alpha \partial_t^7 \breve{v} \cdot \breve{n} - \sigma \sum_{l=0}^{6} c_l \int_\Gamma \breve{g}^{\alpha\beta} \partial_t^l \breve{\eta}^j,_{\alpha\beta}\, \partial_t^{7-l} \breve{n}^j \sqrt{\breve{g}} \partial_t^7 \breve{v} \cdot \breve{n}}_{\mathfrak{j}_3} \\
& \underbrace{- \sigma \sum_{l=1}^{7} c_l \int_\Gamma \partial_t^l \breve{g}^{\alpha\beta} \partial_t^{7-l} [\breve{\eta},_{\alpha\beta} \cdot \breve{n}] \sqrt{\breve{g}} \partial_t^7 \breve{v} \cdot \breve{n}}_{\mathfrak{j}_4} + \underbrace{\int_\Gamma \partial_t^7 \beta_\sigma(t) \sqrt{\breve{g}} \partial_t^7 \breve{v} \cdot \breve{n}}_{0}\,.
\end{aligned} \tag{7.15}$$

Analysis of $\int_0^T \mathfrak{j}_1$ in the time-integral of (7.15)**.** The action of $\bar{\partial}_\alpha \partial_t^5$ in the tangential identity (7.10) provides that

$$\partial_t^6 \breve{v},_\alpha \cdot \breve{\eta},_\gamma = \rho_0^{-1} \breve{J}[\bar{\partial}_{\alpha\gamma}\partial_t^5 \big(\sigma \breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n} - \partial_t^5 \beta_\sigma(t)\big) - \breve{l}_{\gamma\alpha}], \tag{7.16a}$$

where $\breve{l}_{\gamma\alpha}$ is such that $\breve{l}_{\gamma\alpha} \in H^{-0.5}(\Gamma)$, $\sqrt{\sigma}\breve{l}_{\gamma\alpha} \in H^{0.5}(\Gamma)$ and is given by

$$\breve{l}_{\gamma\alpha} = \partial_t^6 \breve{v} \cdot (\breve{\eta},_\gamma \rho_0 \breve{J}^{-1}),_\alpha + \sum_{l=0}^{4} c_l [\partial_t^a \breve{v}_t \cdot \partial_t^{5-l} (\breve{\eta},_\gamma \rho_0 \breve{J}^{-1})],_\alpha\,. \tag{7.16b}$$

Setting $\breve{\ell}_\gamma^{\alpha\beta} = \rho_0^{-1} \breve{J} \sqrt{\breve{g}} \breve{g}^{\alpha\beta} \breve{g}^{\gamma\delta} \breve{v},_\delta \cdot \breve{n}$ we use the tangential identity (7.16), together with the outward normal differentiation formula (2.11c), to find that

$$\mathfrak{j}_1 = \underbrace{\sigma \int_\Gamma \breve{\ell}_\gamma^{\alpha\beta} \bar{\partial}_{\gamma\alpha} \partial_t^5 (\breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n})\, \partial_t^6 \breve{v},_\beta \cdot \breve{n}}_{\mathfrak{j}_1{}'} + \underbrace{\int_\Gamma \breve{\ell}_\gamma^{\alpha\beta} \sqrt{\sigma}\breve{l}_{\gamma\alpha}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\beta \cdot \breve{n}}_{\dddot{\mathcal{R}}} + \dddot{\mathcal{R}}. \tag{7.17}$$

We integrate by parts with respect to a time-derivative of $\partial_t^6 \breve{v},_\beta \cdot \breve{n}$ to write

$$\int_0^T \mathfrak{j}_1{}' =$$

$$- \underbrace{\int_\Gamma \bar\partial_\gamma \partial_t^5 [\breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n}] \, \sigma^2 \bar\partial_\alpha [\breve{\ell}_\gamma^{\alpha\beta} \partial_t^5 \breve{v},_\beta \cdot \breve{n}]}_{\mathfrak{j}_1{}'_A} \Big|_0^T + \int_0^T \underbrace{\sigma^2 \int_\Gamma \bar\partial_\gamma \partial_t^5 [\breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n}] \, \bar\partial_\alpha [\breve{\ell}_\gamma^{\alpha\beta} \partial_t^5 \breve{v},_\beta \cdot \breve{n}_t]}_{\mathfrak{j}_1{}'_B}$$

$$+ \int_0^T \underbrace{\sigma^2 \int_\Gamma \bar\partial_\gamma \partial_t^5 [\breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n}] \, \bar\partial_\alpha [\partial_t \breve{\ell}_\gamma^{\alpha\beta} \partial_t^5 \breve{v},_\beta \cdot \breve{n}]}_{\mathfrak{j}_1{}'_C} + \int_0^T \underbrace{\sigma^2 \int_\Gamma \bar\partial_\gamma \partial_t^6 [\breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n}] \, \bar\partial_\alpha [\breve{\ell}_\gamma^{\alpha\beta} \partial_t^5 \breve{v},_\beta \cdot \breve{n}]}_{\mathfrak{j}_1{}'_D} .$$

Since $\sigma \partial_t^5 \breve{v} \cdot \breve{n}$ is in $H^{2.5}(\Gamma)$, we may take σ sufficiently small so that

$$|\sigma \partial_t^5 \breve{v} \cdot \breve{n}(T)|_2^2 \le \sqrt{\sigma} C |\sqrt{\sigma} \partial_t^5 \breve{v}(T)|_{1.5} |\sigma \partial_t^5 \breve{v} \cdot \breve{n}(T)|_{2.5} \le \int_0^T \dddot{\mathcal{R}}.$$

Thus, the Cauchy-Schwarz inequality provides that

$$\mathfrak{j}_1{}'_A \Big|_0^T = - \int_\Gamma \breve{g}^{\mu\nu} \bar\partial_\gamma [\partial_t^4 \breve{v},_{\mu\nu} \cdot \breve{n}] \, \sigma^2 \breve{\ell}_\gamma^{\alpha\beta} \bar\partial_\alpha [\partial_t^5 \breve{v},_\beta \cdot \breve{n}] \Big|_0^T + \int_0^T \dddot{\mathcal{R}} = \int_0^T \dddot{\mathcal{R}}. \tag{7.18}$$

Since $\sigma \partial_t^5 \breve{v} \cdot \breve{n}$ is in $H^{3.5}(\Gamma)$, we find by use of an $H^{-0.5}(\Gamma)$-duality pairing that

$$\mathfrak{j}_1{}'_B = \underbrace{\sqrt{\sigma} \int_\Gamma \bar\partial_\gamma [\breve{g}^{\mu\nu} \sigma \partial_t^4 \breve{v},_{\mu\nu} \cdot \breve{n}] \, \breve{\ell}_\gamma^{\alpha\beta} \sqrt{\sigma} \partial_t^5 \breve{v},_{\alpha\beta} \cdot \breve{n}_t}_{\dddot{\mathcal{R}}} + \int_0^T \dddot{\mathcal{R}}. \tag{7.19}$$

We employ the Cauchy-Schwarz inequality to conclude that

$$\int_0^T \mathfrak{j}_1{}'_C \le \int_0^T \dddot{\mathcal{R}}. \tag{7.20}$$

We employ the Cauchy-Schwarz inequality or an $H^{-0.5}(\Gamma)$-duality pairing to conclude that

$$\int_0^T \mathfrak{j}_1{}'_D \le \int_0^T \dddot{\mathcal{R}}. \tag{7.21}$$

The inequalities (7.18), (7.19), (7.20), (7.21) establish that

$$\int_0^T \mathfrak{j}_1 \le \int_0^T \dddot{\mathcal{R}}. \tag{7.22}$$

Analysis of $\int_0^T \mathfrak{j}_2$ in the time-integral of (7.15)**.** We equivalently write $\mathfrak{j}_2$ as

$$\mathfrak{j}_2 = \underbrace{\sigma \int_\Gamma \partial_t^6 \breve{v},_\beta \cdot \breve{n} \sqrt{\breve{g}} \breve{g}^{\alpha\beta} \partial_t^7 \breve{v} \cdot \breve{n},_\alpha}_{\mathfrak{j}_{2a}} + \underbrace{\sigma \int_\Gamma \partial_t^6 \breve{v},_\beta \cdot \breve{n},_\alpha \sqrt{\breve{g}} \breve{g}^{\alpha\beta} \partial_t^7 \breve{v} \cdot \breve{n}}_{\mathfrak{j}_{2b}} . \tag{7.23}$$

The action of ∂_t^6 in the tangential identity (7.10) yields

$$\partial_t^7 \breve{v} \cdot \breve{\eta},_\gamma = \rho_0^{-1} \breve{J} [\bar\partial_\gamma \partial_t^6 \big(\sigma \breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n} - \partial_t^6 \beta_\sigma(t) \big) - \breve{l}_\gamma], \tag{7.24a}$$

where $\breve{l}_\gamma$ is such that $\sqrt{\sigma} \breve{l}_\gamma$ is in $H^{0.5}(\Gamma)$ and is given by

$$\breve{l}_\gamma = \sum_{l=0}^5 c_l \partial_t^l \breve{v}_t \cdot \partial_t^{6-l} (\breve{\eta},_\gamma \rho_0 \breve{J}^{-1}). \tag{7.24b}$$

Letting $\breve\ell_\gamma^\beta = -\rho_0^{-1}\breve J\sqrt{\breve g}\breve g^{\alpha\beta}\breve g^{\gamma\delta}\breve\eta,_{\delta\alpha}\cdot\breve n$ and using the tangential identity (7.24), we have that

$$\mathfrak{j}_{2a} = \underbrace{-\sigma^2\int_\Gamma [\partial_t^6\breve v,_\beta\cdot\breve n\breve\ell_\gamma^\beta],_\gamma\,\partial_t^6[\breve g^{\mu\nu}\breve\eta,_{\mu\nu}\cdot\breve n]}_{\mathfrak{j}_{2a}{}'} + \dddot{\mathcal{R}}.$$

We integrate by parts with respect to a time-derivative of $\partial_t^6\breve v,_\beta$ in order to write

$$\begin{aligned}
-\int_0^T \mathfrak{j}_{2a}{}' &= -\sigma^2\int_\Gamma [\breve\ell_\gamma^\beta\,\partial_t^5\breve v,_\beta\cdot\breve n],_\gamma\;\partial_t^6[\breve g^{\mu\nu}\breve\eta,_{\mu\nu}\cdot\breve n]\Big|_0^T + \int_0^T\sigma^2\int_\Gamma[\breve\ell_\gamma^\beta\,\partial_t^5\breve v,_\beta\cdot\breve n_t],_\gamma\;\partial_t^6[\breve g^{\mu\nu}\breve\eta,_{\mu\nu}\cdot\breve n]\\
&\quad+\int_0^T\sigma^2\int_\Gamma\partial_t[\breve\ell_\gamma^\beta\,\partial_t^5\breve v,_\beta\cdot\breve n],_\gamma\;\partial_t^6[\breve g^{\mu\nu}\breve\eta,_{\mu\nu}\cdot\breve n] + \int_0^T\sigma^2\int_\Gamma[\breve\ell_\gamma^\beta\,\partial_t^5\breve v,_\beta\cdot\breve n],_\gamma\;\partial_t^7[\breve g^{\mu\nu}\breve\eta,_{\mu\nu}\cdot\breve n]\\
&= \int_0^T \underbrace{\sigma^2\int_\Gamma[\breve\ell_\gamma^\beta\,\partial_t^5\breve v,_\beta\cdot\breve n],_\gamma\;\breve g^{\mu\nu}\partial_t^6\breve v,_{\mu\nu}\cdot\breve n}_{\mathfrak{j}_{2a}{}''} + \int_0^T\dddot{\mathcal{R}}.
\end{aligned}$$

We write

$$\int_0^T\mathfrak{j}_{2a}{}'' = \int_0^T\underbrace{\sigma^2\int_\Gamma\breve\ell_\gamma^\beta\,\partial_t^5\breve v,_{\beta\gamma}\cdot\breve n\,\breve g^{\mu\nu}\partial_t^6\breve v,_{\mu\nu}\cdot\breve n}_{\mathfrak{j}_{2a}{}'''} + \int_0^T\underbrace{\sigma^2\int_\Gamma\partial_t^5\breve v,_\beta\cdot[\breve n\breve\ell_\gamma^\beta],_\gamma\;\breve g^{\mu\nu}\partial_t^6\breve v,_{\mu\nu}\cdot\breve n}_{\mathfrak{j}}\,.$$

Using integration by parts with respect to a time-derivative of $\partial_t^6\breve v,_{\mu\nu}$, we conclude that $\int_0^T\mathfrak{j} = \int_0^T\dddot{\mathcal{R}}$. Letting $\breve\ell = \rho_0^{-1}\breve J\sqrt{\breve g}\,\breve g^{\gamma\delta}\breve\eta,_{\gamma\delta}\cdot\breve n$ we utilize the symmetry of $\breve\ell_\gamma^\beta$ to exchange $\bar\partial_\alpha$ and $\bar\partial_\gamma$ via integration by parts for

$$\begin{aligned}
\int_0^T\mathfrak{j}_{2a}{}''' &= \frac12\int_\Gamma\breve\ell\,|\breve g^{\mu\nu}\sigma\partial_t^5\breve v,_{\mu\nu}\cdot\breve n|^2\Big|_0^T\\
&\quad-\int_0^T\sigma^2\int_\Gamma\breve\ell\,\partial_t^5\breve v,_{\alpha\beta}\cdot\breve n\,\partial_t^5\breve v,_{\mu\nu}\cdot(n\breve g^{\mu\nu})_t - \frac12\int_0^T\int_\Gamma\breve\ell_t\,|\breve g^{\mu\nu}\sigma\partial_t^5\breve v,_{\mu\nu}\cdot\breve n|^2 = \int_0^T\dddot{\mathcal{R}}.
\end{aligned}\tag{7.25}$$

We have thus established that

$$\int_0^T\mathfrak{j}_{2a} = \int_0^T\dddot{\mathcal{R}}.$$

The analysis of $\int_0^T\mathfrak{j}_{2b}$ is similar. We set $\breve\ell^{\gamma\beta} = \sqrt{\breve g}\breve g^{\alpha\beta}\breve g^{\gamma\delta}\breve\eta,_{\delta\alpha}\cdot\breve n$ and write

$$\begin{aligned}
\int_0^T\mathfrak{j}_{2b} &= -\sigma^{\frac32}\int_\Gamma\partial_t^6\breve v\cdot\Big[\breve n,_\alpha\,\sqrt{\breve g}\breve g^{\alpha\beta}\,\sqrt{\sigma}\partial_t^6\breve v\cdot\breve n\Big],_\beta\Big|_0^T - \int_0^T\sigma^2\int_\Gamma\partial_t^6\breve v^j,_\beta\,[\breve n^j,_\alpha\,\sqrt{\breve g}\breve g^{\alpha\beta}\breve n^i]_t\,\partial_t^6\breve v^i\\
&\quad+\int_0^T\underbrace{\sigma^2\int_\Gamma\breve\ell^{\gamma\beta}\,\partial_t^7\breve v,_\beta\cdot\breve\eta,_\gamma\;\partial_t^6\breve v\cdot\breve n}_{\mathfrak{j}_{2b}{}'} = \int_0^T\dddot{\mathcal{R}} + \int_0^T\mathfrak{j}_{2b}{}'.
\end{aligned}$$

Regarding $\int_0^T\mathfrak{j}_{2b}{}'$, we use the identity

$$\partial_t^7\breve v,_\beta\cdot\breve\eta,_\gamma = \rho_0^{-1}\breve J[\bar\partial_{\beta\gamma}\partial_t^6\big(\breve g^{\mu\nu}\breve\eta,_{\mu\nu}\cdot\breve n - \beta_\sigma(t)\big) - \breve l'_{\beta\gamma}],\tag{7.26}$$

where $\breve l'_{\beta\gamma} = \partial_t^7\breve v\cdot\breve\eta,_{\gamma\beta}\,\rho_0\breve J^{-1} + \partial_t^7\breve v\cdot\breve\eta,_\gamma\,(\rho_0\breve J^{-1}),_\beta + \bar\partial_\beta\breve l_\gamma$, with $\breve l_\gamma$ given by (7.24b). We integrate by parts with respect to $\bar\partial_{\beta\gamma}$ in $\sigma^3\int_0^T\int_\Gamma\rho_0^{-1}\breve J\bar\partial_{\beta\gamma}\partial_t^6[\breve g^{\mu\nu}\breve\eta,_{\mu\nu}\cdot\breve n]\,\breve\ell^{\gamma\beta}\,\partial_t^6\breve v\cdot\breve n$, where the highest-order term produced by $\bar\partial_{\beta\gamma}$-integration by parts is (7.25). To estimate the integral where $\bar\partial_\beta\breve l_\gamma$ appears, we have the choice of integration by parts with respect to $\bar\partial_\beta$

or an $H^{-0.5}(\Gamma)$-duality pairing. In the integral where $\partial_t^7\breve{v}\cdot\breve{\eta},_\gamma(\rho_0\breve{J}^{-1}),_\beta$ appears, we use the identity (7.24). Thus, we conclude that $\int_0^T \mathfrak{j}_{2b}{}' = \int_0^T \dddot{\mathcal{R}}$ and

$$\int_0^T \mathfrak{j}_2 \le \int_0^T \dddot{\mathcal{R}}. \tag{7.27}$$

Analysis of $\int_0^T \mathfrak{j}_3$ in the time-integral of (7.15)**.** We equivalently write $\mathfrak{j}_3$ as

$$\mathfrak{j}_3 = \underbrace{\sigma\int_\Gamma \partial_t^6\breve{v},_\beta\cdot\breve{n}(\sqrt{\breve{g}}\breve{g}^{\alpha\beta}),_\alpha\,\partial_t^7\breve{v}\cdot\breve{n}}_{\mathfrak{j}_{3a}} - \sum_{l=0}^{6} c_l\,\underbrace{\sigma\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\partial_t^l\breve{\eta}^j,_{\alpha\beta}\,\partial_t^{7-l}\breve{n}^j\,\partial_t^7\breve{v}\cdot\breve{n}}_{\mathfrak{j}_{3b,l}}\,.$$

We infer from our analysis of (4.25) that

$$-\int_0^T \mathfrak{j}_{3b,0} = \int_0^T \underbrace{\sigma\int_\Gamma \sqrt{\breve{g}}\,\breve{g}^{\alpha\beta}\breve{g}^{\gamma\delta}\breve{\eta},_{\alpha\beta}\cdot\breve{\eta},_\gamma\,\partial_t^6\breve{v},_\delta\cdot\breve{n}\,\partial_t^7\breve{v}\cdot\breve{n}}_{-\mathfrak{j}_{3a}} + \int_0^T \dddot{\mathcal{R}}.$$

Hence,

$$\mathfrak{j}_3 = -\sum_{l=1}^{6} c_l\,\mathfrak{j}_{3b,l} + \dddot{\mathcal{R}}.$$

The terms $\int_0^T \mathfrak{j}_{3b,l}$ for $l = 1,\ldots,5$, are analyzed by integrating by parts with respect to a time-derivative of $\partial_t^7\breve{v}$ and then using elementary estimates. Thus,

$$\mathfrak{j}_3 = -\mathfrak{j}_{3b,6} + \dddot{\mathcal{R}}.$$

Integration by parts with respect to a time-derivative of $\partial_t^7\breve{v}$ yields

$$\begin{aligned}\int_0^T \mathfrak{j}_{3b,6} &= -\int_0^T \sigma\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\partial_t^6\breve{v},_{\alpha\beta}\cdot\breve{n}_t\,\partial_t^6\breve{v}\cdot\breve{n} + \int_0^T \dddot{\mathcal{R}}\\ &= \int_0^T \underbrace{\sigma\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\partial_t^6\breve{v},_\beta\cdot\breve{n}_t\,\partial_t^6\breve{v},_\alpha\cdot\breve{n}}_{\mathfrak{j}_{3b,6}{}'} + \int_0^T \dddot{\mathcal{R}}.\end{aligned}$$

Letting $\breve{\ell}^\gamma_{\alpha\beta} = \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\breve{g}^{\gamma\delta}\breve{v},_\delta\cdot\breve{n}$, we once again integrate by parts with respect to time:

$$\begin{aligned}&\int_0^T \mathfrak{j}_{3b,6}{}' = -\int_\Gamma \partial_t^6\breve{v},_\beta\cdot\Big[\breve{\eta},_\gamma\,\breve{\ell}^\gamma_{\alpha\beta}\sigma\partial_t^5\breve{v},_\alpha\cdot\breve{n}\Big],_\beta\Big|_0^T\\ &-\int_0^T \sigma\int_\Gamma \partial_t^6\breve{v}^j,_\beta\,\partial_t^5\breve{v}^i,_\alpha\cdot(\breve{n}^i\,\breve{\eta}^j,_\beta\,\breve{\ell}^\gamma_{\alpha\beta})_t - \int_0^T \underbrace{\sigma\int_\Gamma \breve{\ell}^\gamma_{\alpha\beta}\,\partial_t^7\breve{v},_\beta\cdot\breve{\eta},_\gamma\,\partial_t^5\breve{v},_\alpha\cdot\breve{n}}_{\mathfrak{j}} = -\int_0^T \mathfrak{j} + \int_0^T \dddot{\mathcal{R}}.\end{aligned}$$

We conclude via the tangential identity (7.26) that $\mathfrak{j} = \dddot{\mathcal{R}}$. Hence,

$$\int_0^T \mathfrak{j}_3 \le \int_0^T \dddot{\mathcal{R}}. \tag{7.28}$$

Analysis of $\int_0^T \mathfrak{j}_4$ in the time-integral of (7.15)**.** We integrate by parts with respect to a time-derivative of $\partial_t^7 \breve{v}$ in the $\int_0^T \mathfrak{j}_4$-terms and, if need be, spatially integrate by parts. For example, letting $\mathfrak{j}_4 = \sum_{l=1}^7 \mathfrak{j}_{4,l}$, we find that after integration by parts with respect to time,

$$\begin{aligned}\int_0^T \mathfrak{j}_{4,1} &= \int_0^T \sigma \int_\Gamma (\sqrt{\breve{g}}\breve{g}^{\alpha\beta})_t \partial_t^7 (\breve{\eta},_{\alpha\beta} \cdot \breve{n})\, \partial_t^6 \breve{v} \cdot \breve{n} + \int_0^T \dddot{\mathcal{R}} \\ &= -\int_0^T \sigma \int_\Gamma \partial_t^6 \breve{v},_\alpha \cdot \bar{\partial}_\beta [\breve{n}\, (\sqrt{\breve{g}}\breve{g}^{\alpha\beta})_t \partial_t^6 \breve{v} \cdot \breve{n}] + \int_0^T \dddot{\mathcal{R}} = \int_0^T \dddot{\mathcal{R}}.\end{aligned}$$

Similarly, integration by parts with respect to time provides for the expression

$$\sum_{l=2}^6 \int_0^T \mathfrak{j}_{4,l} = \int_0^T \dddot{\mathcal{R}}.$$

Finally, using the differentiation formulas (2.10a) and (2.10b),

$$\int_0^T \mathfrak{j}_{4,7} = \int_0^T \int_\Gamma \sqrt{\breve{g}}(2\breve{g}^{\alpha\mu}\breve{g}^{\nu\beta} - \breve{g}^{\alpha\beta}\breve{g}^{\mu\nu})\, \partial_t^6 \breve{v},_\mu \cdot \breve{\eta},_\nu\ \breve{\eta},_{\alpha\beta} \cdot \breve{n}\, \partial_t^7 \breve{v} \cdot \breve{n} + \int_0^T \dddot{\mathcal{R}} = \int_0^T \dddot{\mathcal{R}},$$

where the second equality follows from our above analysis of $\int_0^T \mathfrak{j}_{2b}$.

Hence,

$$\int_0^T \mathfrak{j}_4 \le \int_0^T \dddot{\mathcal{R}}. \tag{7.29}$$

Rewriting equation (7.13)**.** The inequalities (7.22), (7.27), (7.28), (7.29) provide that the boundary integral $\mathfrak{i}$ expressed as (7.15) satisfies

$$\mathfrak{i} = \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\alpha \cdot \breve{n}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\beta \cdot \breve{n} + \dddot{\mathcal{R}}. \tag{7.30}$$

Using the identities (7.14) and (7.30), we have that (7.13) is equivalently written as

$$\frac{d}{dt}\int_\Omega \rho_0^2 \breve{J}^{-3} |\partial_t^7 \breve{J}|^2 + 2\frac{d}{dt}\int_\Omega \rho_0^{-1} \breve{J}^4 |\breve{A}^k_\cdot \partial_t^6 \breve{J},_k|^2 + \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\alpha \cdot \breve{n}\, \sqrt{\sigma}\partial_t^6 \breve{v},_\beta \cdot \breve{n} = \dddot{\mathcal{R}}. \tag{7.31}$$

The time-integral of (7.31) completes the proof. □

Step 3: The energy estimates for the action of $\bar{\partial}^a \partial_t^{8-2a}$, $a = 1, 2, 3, 4$. We define the vector $\breve{a}^k_\alpha$ as $\breve{\eta},_\alpha \cdot \breve{a}^k_\cdot$ and let the vector $\breve{a}^k_{\breve{n}}$ be defined as $\breve{n} \cdot \breve{a}^k_\cdot$. Using these vector identities, we decompose the cofactor matrix $\breve{a}$ as

$$\breve{a}^k_r = \breve{a}^k_\alpha \breve{g}^{\alpha\beta} \breve{\eta}^r,_\beta + \breve{a}^k_{\breve{n}} \breve{n}^r \quad \text{on } \Gamma. \tag{7.32}$$

Since $\breve{a}^k_\cdot N^k = \sqrt{\breve{g}}\breve{n}$ by the formula (2.9), it follows that

$$\breve{a}^\cdot_{\breve{n}} = \frac{1}{\sqrt{\breve{g}}} N^j \breve{a}^j_\ell \breve{a}^\cdot_\ell. \tag{7.33}$$

According to the identity (7.33), the lower-bound (7.7b) is equivalently stated as

$$0 < \frac{\nu}{2} \le -\breve{a}^k_{\breve{n}} (\rho_0^2 \breve{J}^{-2}),_k\,. \tag{7.34}$$

Proposition 7.2 (Energy estimates for the action of $\bar{\partial}\partial_t^6$ in the Euler equations (7.2a))**.**

$$\sup_{t\in[0,T]} \|\bar{\partial}\partial_t^6 \breve{v}(t)\|_0^2 + \sup_{t\in[0,T]} |\sqrt{\sigma}\bar{\partial}\partial_t^5 \breve{v} \cdot \breve{n}(t)|_1^2 + \sup_{t\in[0,T]} |\bar{\partial}\partial_t^5 \breve{v} \cdot \breve{n}(t)|_0^2 \le \int_0^T \dddot{\mathcal{R}}.$$

Proof. Testing the action of $\bar\partial\partial_t^6$ in the Euler equations (7.2a) against $\bar\partial\partial_t^6\breve v$ in the $L^2(\Omega)$-inner product, and integrating by parts in the integral $\int_\Omega \breve a_i^k\bar\partial\partial_t^6(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^6\breve v^i$ yields

$$\int_\Omega \bar\partial\partial_t^6[\rho_0\breve v_t^i]\bar\partial\partial_t^6\breve v^i + \underbrace{\int_\Omega \bar\partial\partial_t^6\breve a_i^k[\rho_0^2\breve J^{-2}],_k\,\bar\partial\partial_t^6\breve v^i}_{\mathcal{K}} - \underbrace{\int_\Omega \bar\partial\partial_t^6[\rho_0^2\breve J^{-2}]\breve a_i^k\bar\partial\partial_t^6\breve v^i,_k}_{\mathcal{I}}$$
$$+\underbrace{\int_\Gamma \bar\partial\partial_t^6[\rho_0^2\breve J^{-2}]\breve a_i^k\bar\partial\partial_t^6\breve v^i N^k}_{\mathfrak{i}} = \dddot{\mathcal{R}}. \qquad (7.35)$$

We used the Cauchy-Schwarz inequality or an L^4–L^4–L^2 Hölder inequality to analyze the lower-order terms in (7.35).

Analysis of $\mathcal{I}$ in (7.35)**.** We have that

$$-\mathcal{I} = \underbrace{\frac{d}{dt}\int_\Omega \rho_0^2\breve J^{-3}|\bar\partial\partial_t^6\breve J|^2}_{\int_0^T\dddot{\mathcal{R}}} + \underbrace{\int_\Omega \bar\partial\partial_t^6(\rho_0^2\breve J^{-2})\bar\partial\partial_t^6\breve a_i^k\breve v^i,_k}_{\mathcal{I}_1} + \underbrace{\int_\Omega \bar\partial\partial_t^6(\rho_0^2\breve J^{-2})\partial_t\breve a_i^k\bar\partial\partial_t^5\breve v^i,_k}_{\mathcal{I}_2} + \dddot{\mathcal{R}}.$$

Integration by parts with respect to a time-derivative of $\partial_t^6(\rho_0^2\breve J^{-2})$ yields

$$\int_0^T \mathcal{I}_1 = -\underbrace{\int_\Omega \bar\partial\Big[\bar\partial\partial_t^5(\rho_0^2\breve J^{-2})\breve v^i,_k\Big]\partial_t^6\breve a_i^k\Big|_0^T}_{\mathcal{I}_{1A}} - \int_0^T\underbrace{\int_\Omega \bar\partial\partial_t^5(\rho_0^2\breve J^{-2})\partial_t\big(\bar\partial\partial_t^6\breve a_i^k\breve v^i,_k\big)}_{\mathcal{I}_{1B}}. \qquad (7.36)$$

Since $\partial_t^5\breve J$ is in $H^2(\Omega)$ and the highest-order term of $\partial_t^6\breve a_i^k$ scales like $D\partial_t^5\breve v$, the term $\mathcal{I}_{1A}$ is bounded by $\int_0^T\dddot{\mathcal{R}}$ thanks to the fundamental theorem of calculus. Since the analysis of $\mathcal{I}_{1B}$ is similar, we have established that $\mathcal{I}_1 = \dddot{\mathcal{R}}$. Similarly, $\mathcal{I}_2 = \dddot{\mathcal{R}}$. Thus,

$$\mathcal{I} = \dddot{\mathcal{R}}. \qquad (7.37)$$

Analysis of $\mathcal{K}$ in (7.35)**.** From the differentiation formula (2.6) we infer that

$$\mathcal{K} = \int_\Omega \breve J^{-1}\big(\breve a_r^s\breve a_i^k - \breve a_i^s\breve a_r^k\big)\bar\partial\partial_t^5\breve v^r,_s(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^6\breve v^i + \dddot{\mathcal{R}}.$$

Since $\breve a_r^s\bar\partial\partial_t^5\breve v^r,_s$ is equal to $\bar\partial\partial_t^6\breve J$ plus lower-order terms, we write

$$\mathcal{K} = -\int_\Omega \breve J^{-1}\breve a_r^k(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^5\breve v^r,_s\,\breve a_i^s\bar\partial\partial_t^6\breve v^i + \dddot{\mathcal{R}}$$
$$= \underbrace{\int_\Omega \breve J^{-1}\breve a_r^k(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^5\breve v^r\breve a_i^s\bar\partial\partial_t^6\breve v^i,_s}_{\mathcal{K}'} - \underbrace{\int_\Gamma \breve J^{-1}\breve a_r^k(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^5\breve v^r\breve a_i^s\bar\partial\partial_t^6\breve v^i N^s}_{\mathbf{k}} + \dddot{\mathcal{R}}.$$

Integrating by parts with respect to a time-derivative of $\bar\partial\partial_t^6\breve v^i,_s$ yields

$$\int_0^T \mathcal{K}' = \underbrace{\int_\Omega \breve J^{-1}\breve a_r^k(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^5\breve v^r\breve a_i^s\bar\partial\partial_t^5\breve v^i,_s\Big|_0^T}_{\int_0^T\dddot{\mathcal{R}}}$$
$$-\int_0^T\underbrace{\int_\Omega \breve J^{-1}\breve a_r^k(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^5\breve v^r\breve a_i^s\bar\partial\partial_t^5\breve v^i,_s}_{\dddot{\mathcal{R}}} - \int_0^T\underbrace{\int_\Omega \breve J^{-1}\breve a_r^k(\rho_0^2\breve J^{-2}),_k\,\bar\partial\partial_t^5\breve v^r(\breve a_i^s)_t\bar\partial\partial_t^5\breve v^i,_s}_{\mathcal{K}''},$$

where we have again used that $\breve{a}_i^s \bar\partial \partial_t^5 \breve{v}^i{,_s}$ is equal to $\bar\partial \partial_t^6 \breve{J}$ plus lower-order terms. Lemma 2.5 provides that

$$\int_0^T \mathcal{K}'' \leq C \int_0^T \|\bar\partial \partial_t^5 \breve{v}\|_{0.5} \|\bar\partial D \partial_t^5 \breve{v}\|_{H^{0.5}(\Omega)'} \leq C \int_0^T \|\partial_t^5 \breve{v}\|_{1.5}^2 \leq \int_0^T \dddot{\mathcal{R}}.$$

The decomposition (7.32) provides that

$$-\mathbf{k} = -\underbrace{\int_\Gamma \sqrt{\breve{g}}(\rho_0^2 \breve{J}^{-2}),_\alpha \, \bar\partial \partial_t^5 \breve{v} \cdot \breve{\eta},_\beta \, \breve{g}^{\alpha\beta} \bar\partial \partial_t^6 \breve{v} \cdot \breve{n}}_{\mathbf{k}_1} - \underbrace{\int_\Gamma \sqrt{\breve{g}} \breve{J}^{-1} \breve{a}^k_{\breve{n}} (\rho_0^2 \breve{J}^{-2}),_k \, \bar\partial \partial_t^5 \breve{v} \cdot \breve{n} \, \bar\partial \partial_t^6 \breve{v} \cdot \breve{n}}_{\mathbf{k}_2}.$$

We have used the identities $\breve{J}^{-1} \breve{a} = \breve{A}$ and $\breve{\eta},_\alpha \cdot \breve{A}^k_{\cdot} = \delta^k_\alpha$, where δ^k_α is the Kronecker delta, in writing $\mathbf{k}_1$. Thanks to the Laplace-Young boundary condition (7.2b), we have that

$$\mathbf{k}_1 = \underbrace{\int_\Gamma \sqrt{\breve{g}}(\beta_\sigma(t)),_\alpha \, \bar\partial \partial_t^5 \breve{v} \cdot \breve{\eta},_\beta \, \breve{g}^{\alpha\beta} \bar\partial \partial_t^6 \breve{v} \cdot \breve{n} - \int_\Gamma \sqrt{\breve{g}}(\breve{g}^{\mu\nu} \breve{\eta},_{\mu\nu} \cdot \breve{n}),_\alpha \sqrt{\sigma} \bar\partial \partial_t^5 \breve{v} \cdot \breve{\eta},_\beta \, \breve{g}^{\alpha\beta} \sqrt{\sigma} \bar\partial \partial_t^6 \breve{v} \cdot \breve{n}}_{\dddot{\mathcal{R}}}.$$

We have used the identity (7.3) to deduce that $\bar\partial \beta_\sigma(t)$ scales like σ in $L^\infty(\Gamma)$ in the first term on the right-hand side and we have used an L^4–L^4–L^2 Hölder inequality in the second term on the right-hand side. We also have that

$$\begin{aligned} -\mathbf{k}_2 = &\frac{1}{2} \frac{d}{dt} \int_\Gamma \sqrt{\breve{g}} \breve{J}^{-1} [-\breve{a}^k_{\breve{n}} (\rho_0^2 \breve{J}^{-2}),_k] |\bar\partial \partial_t^5 \breve{v} \cdot \breve{n}|^2 \\ &+ \underbrace{\frac{1}{2} \int_\Gamma \partial_t \big(\sqrt{\breve{g}} \breve{J}^{-1} \breve{a}^k_{\breve{n}} (\rho_0^2 \breve{J}^{-2}),_k\big) |\bar\partial \partial_t^5 \breve{v} \cdot \breve{n}|^2}_{\dddot{\mathcal{R}}} + \underbrace{\int_\Gamma \sqrt{\breve{g}} \breve{J}^{-1} \breve{a}^k_{\breve{n}} (\rho_0^2 \breve{J}^{-2}),_k \, \bar\partial \partial_t^5 \breve{v} \cdot \breve{n} \bar\partial \partial_t^5 \breve{v} \cdot \breve{n}_t}_{\dddot{\mathcal{R}}}. \end{aligned}$$

Hence,

$$\mathcal{K} = \frac{1}{2} \frac{d}{dt} \int_\Gamma \sqrt{\breve{g}} \breve{J}^{-1} [-\breve{a}^k_{\breve{n}} (\rho_0^2 \breve{J}^{-2}),_k] |\bar\partial \partial_t^5 \breve{v} \cdot \breve{n}|^2 + \dddot{\mathcal{R}}. \tag{7.38}$$

Analysis of $\mathfrak{i}$ in (7.35)**.** Using the Laplace-Young boundary condition (7.2b), we find that

$$\begin{aligned}
\mathfrak{i} = \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\breve g}\breve g^{\alpha\beta}\,\sqrt{\sigma}\bar\partial\partial_t^5\breve v,_\alpha\cdot\breve n\,\sqrt{\sigma}\bar\partial\partial_t^5\breve v,_\beta\cdot\breve n - \underbrace{\sigma\int_\Gamma \sqrt{\breve g}\partial_t^5\breve v,_{\alpha\beta}\cdot\bar\partial(\breve n\breve g^{\alpha\beta})\bar\partial\partial_t^6\breve v\cdot\breve n}_{\mathfrak{j}_1} \\
\underbrace{-\sigma\int_\Gamma \sqrt{\breve g}\breve g^{\alpha\beta}\bar\partial\partial_t^5\breve v,_\alpha\cdot\breve n_t\,\bar\partial\partial_t^5\breve v,_\beta\cdot\breve n}_{\mathfrak{j}_2} - \frac{1}{2}\underbrace{\int_\Gamma(\sqrt{\breve g}\breve g^{\alpha\beta})_t\,\sqrt{\sigma}\bar\partial\partial_t^5\breve v,_\alpha\cdot\breve n\,\sqrt{\sigma}\bar\partial\partial_t^5\breve v,_\beta\cdot\breve n}_{\dddot{\mathcal{R}}} \\
+\underbrace{\int_\Gamma \sigma\bar\partial\partial_t^5\breve v,_\beta\cdot\breve n\sqrt{\breve g}\breve g^{\alpha\beta}\bar\partial\partial_t^6\breve v\cdot\breve n,_\alpha}_{\dddot{\mathcal{R}}} + \underbrace{\sigma\int_\Gamma \bar\partial\partial_t^5\breve v,_\beta\cdot\breve n,_\alpha\,\sqrt{\breve g}\breve g^{\alpha\beta}\bar\partial\partial_t^6\breve v\cdot\breve n}_{\mathfrak{j}_3} \\
+\underbrace{\sigma\int_\Gamma \bar\partial\partial_t^5\breve v,_\beta\cdot\breve n(\sqrt{\breve g}\breve g^{\alpha\beta}),_\alpha\,\bar\partial\partial_t^6\breve v\cdot\breve n}_{\dddot{\mathcal{R}}} + \underbrace{\sum_{l=0}^{5} c_l\int_\Gamma \breve g^{\alpha\beta}\partial_t^l\breve\eta^j,_{\alpha\beta}\,\partial_t^{6-l}\breve n^j\,\bar\partial\big(\sigma\sqrt{\breve g}\bar\partial\partial_t^6\breve v\cdot\breve n\big)}_{\mathfrak{j}_4} \\
+\underbrace{\sum_{l=1}^{6} c_l\int_\Gamma \partial_t^l\breve g^{\alpha\beta}\partial_t^{6-l}[\breve\eta,_{\alpha\beta}\cdot\breve n]\bar\partial\big(\sigma\sqrt{\breve g}\bar\partial\partial_t^6\breve v\cdot\breve n\big)}_{\mathfrak{j}_5} + \underbrace{\int_\Gamma \bar\partial\partial_t^6\beta_\sigma(t)\bar\partial\partial_t^6\breve v\cdot\breve n}_{\dddot{\mathcal{R}}}\,. \qquad (7.39)
\end{aligned}$$

We integrate by parts with respect to a time-derivative of $\bar\partial\partial_t^6\breve v\cdot\breve n$ to write

$$\begin{aligned}
&\int_0^T \mathfrak{j}_1 = \\
&-\int_\Gamma \sqrt{\breve g}\partial_t^5\breve v,_\alpha\cdot\Big[\bar\partial(\breve n\breve g^{\alpha\beta})\sigma\bar\partial\partial_t^5\breve v\cdot\breve n\Big],_\beta\Big|_0^T - \int_0^T\int_\Gamma \sqrt{\breve g}\sqrt{\sigma}\partial_t^5\breve v,_{\alpha\beta}\cdot\bar\partial(\breve n\breve g^{\alpha\beta})\sqrt{\sigma}\bar\partial\partial_t^5\breve v\cdot\breve n_t \\
&+\int_0^T\int_\Gamma \partial_t^5\breve v,_\alpha\cdot\Big[[\bar\partial(\breve n\breve g^{\alpha\beta})\sqrt{\breve g}]_t\sigma\bar\partial\partial_t^5\breve v\cdot\breve n\Big],_\beta + \int_0^T\int_\Gamma \sqrt{\breve g}\partial_t^6\breve v,_\alpha\cdot\Big[\bar\partial(\breve n\breve g^{\alpha\beta})\sigma\bar\partial\partial_t^5\breve v\cdot\breve n\Big],_\beta\,.
\end{aligned}$$

Since $\sigma\partial_t^5\breve v\cdot\breve n\in H^{2.5}(\Gamma)$, it follows that

$$\int_0^T \mathfrak{j}_1 = \int_0^T \dddot{\mathcal{R}}.$$

Similarly, integration by parts with respect to a time-derivative of $\bar\partial\partial_t^5\breve v,_\beta\cdot\breve n$ yields

$$\int_0^T \mathfrak{j}_2 = \int_0^T \dddot{\mathcal{R}}.$$

Since $\sqrt{\sigma}\partial_t^5\breve v\cdot\breve n\in H^2(\Gamma)$ and $\sqrt{\sigma}\partial_t^4\breve v\in H^{2.5}(\Gamma)$, for the $l=5$ term of $\mathfrak{j}_4$, we have that

$$\begin{aligned}
&\int_0^T\int_\Gamma \breve g^{\alpha\beta}\partial_t^4\breve v^j,_{\alpha\beta}\,\partial_t\breve n^j\,\bar\partial\big(\sigma\sqrt{\breve g}\bar\partial\partial_t^6\breve v\cdot\breve n\big) = \\
&\int_\Gamma \breve g^{\alpha\beta}\sqrt{\sigma}\partial_t^4\breve v^j,_{\alpha\beta}\,\partial_t\breve n^j\,\bar\partial\big(\sqrt{\sigma}\sqrt{\breve g}\bar\partial\partial_t^5\breve v\cdot\breve n\big)\Big|_0^T - \int_0^T\int_\Gamma \breve g^{\alpha\beta}\sqrt{\sigma}\partial_t^4\breve v^j,_{\alpha\beta}\,\partial_t\breve n^j\,\bar\partial\big(\sqrt{\sigma}\sqrt{\breve g}\bar\partial\partial_t^5\breve v\cdot\breve n_t\big) \\
&-\int_0^T\int_\Gamma \partial_t^4\breve v^j,_{\alpha\beta}\,(\partial_t\breve n^j\breve g^{\alpha\beta})_t\,\bar\partial\big(\sigma\sqrt{\breve g}\bar\partial\partial_t^5\breve v\cdot\breve n\big) - \int_0^T\int_\Gamma \breve g^{\alpha\beta}\partial_t^5\breve v^j,_{\alpha\beta}\,\partial_t\breve n^j\,\bar\partial\big(\sigma\sqrt{\breve g}\bar\partial\partial_t^5\breve v\cdot\breve n\big) \\
&= \int_0^T \dddot{\mathcal{R}} - \int_0^T\int_\Gamma \breve g^{\alpha\beta}\partial_t^5\breve v^j,_{\alpha\beta}\,\partial_t\breve n^j\,\bar\partial\big(\sigma\sqrt{\breve g}\bar\partial\partial_t^5\breve v\cdot\breve n\big) = \int_0^T \dddot{\mathcal{R}}.
\end{aligned}$$

The last equality follows from the analysis of $\int_0^T \mathfrak{j}_2$. Since the analysis of $\mathfrak{j}_5$ and the $l = 0, \dots, 4$ terms of $\mathfrak{j}_4$ are similarly established, we infer that (7.39) is equally written as

$$\mathfrak{i} = \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\,\sqrt{\sigma}\bar{\partial}\partial_t^5\breve{v},_\alpha \cdot\breve{n}\,\sqrt{\sigma}\bar{\partial}\partial_t^5\breve{v},_\beta \cdot\breve{n} + \dddot{\mathcal{R}}. \tag{7.40}$$

The time-integral of (7.35)**.** Using the identities (7.37), (7.38) and (7.40) in the time-integral of the equation (7.35) completes the proof. ☐

Proposition 7.3 (Energy estimates for the action of $\bar{\partial}^2\partial_t^4$ in the Euler equations (7.2a))**.**

$$\sup_{t\in[0,T]} \|\bar{\partial}^2\partial_t^4\breve{v}(t)\|_0^2 + \sup_{t\in[0,T]} |\sqrt{\sigma}\bar{\partial}^2\breve{v}_{ttt}\cdot\breve{n}(t)|_1^2 + \sup_{t\in[0,T]} |\bar{\partial}^2\breve{v}_{ttt}\cdot\breve{n}(t)|_0^2 \le \int_0^T \dddot{\mathcal{R}}.$$

Proof. Testing the action of $\bar{\partial}^2\partial_t^4$ in the Euler equations (7.2a) against $\bar{\partial}^2\partial_t^4\breve{v}$ in the $L^2(\Omega)$-inner product, and integrating by parts in the integral $\int_\Omega \breve{a}_i^k\bar{\partial}^2\partial_t^4(\rho_0^2\breve{J}^{-2}),_k\,\bar{\partial}^2\partial_t^4\breve{v}^i$ yields

$$\int_\Omega \bar{\partial}^2\partial_t^4[\rho_0\breve{v}_t^i]\bar{\partial}^2\partial_t^4\breve{v}^i + \underbrace{\int_\Omega \bar{\partial}^2\partial_t^4\breve{a}_i^k[\rho_0^2\breve{J}^{-2}],_k\,\bar{\partial}^2\partial_t^4\breve{v}^i}_{\mathcal{K}} - \underbrace{\int_\Omega \bar{\partial}^2\partial_t^4[\rho_0^2\breve{J}^{-2}]\breve{a}_i^k\bar{\partial}^2\partial_t^4\breve{v}^i,_k}_{\mathcal{I}}$$
$$+ \underbrace{\int_\Gamma \bar{\partial}^2\partial_t^4[\rho_0^2\breve{J}^{-2}]\breve{a}_i^k\bar{\partial}^2\partial_t^4\breve{v}^i N^k}_{\mathfrak{i}} = \dddot{\mathcal{R}}.$$

The analysis of $\mathcal{K}$ and $\mathfrak{i}$ follows from the proof of Proposition 7.2. Hence,

$$\frac{1}{2}\frac{d}{dt}\int_\Omega \rho_0|\bar{\partial}^2\partial_t^4\breve{v}|^2 + \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\breve{g}}\breve{J}^{-1}[-\breve{a}_{\breve{n}}^k(\rho_0^2\breve{J}^{-2}),_k\,]|\bar{\partial}^2\breve{v}_{ttt}\cdot\breve{n}|^2$$
$$+ \frac{1}{2}\frac{d}{dt}\int_\Gamma \sqrt{\breve{g}}\breve{g}^{\alpha\beta}\,\sqrt{\sigma}\bar{\partial}^2\breve{v}_{ttt},_\alpha \cdot\breve{n}\,\sqrt{\sigma}\bar{\partial}^2\breve{v}_{ttt},_\beta \cdot\breve{n} = \dddot{\mathcal{R}} + \mathcal{I}. \tag{7.41}$$

We have that

$$-\mathcal{I} = \frac{d}{dt}\underbrace{\int_\Omega \rho_0^2\breve{J}^{-3}|\bar{\partial}^2\partial_t^4\breve{J}|^2}_{\int_0^T \dddot{\mathcal{R}}} + \int_\Omega \bar{\partial}^2\partial_t^4(\rho_0^2\breve{J}^{-2})\bar{\partial}^2\partial_t^4\breve{a}_i^k\breve{v}^i,_k + \int_\Omega \bar{\partial}^2\partial_t^4(\rho_0^2\breve{J}^{-2})\partial_t\breve{a}_i^k\bar{\partial}^2\partial_t^3\breve{v}^i,_k + \dddot{\mathcal{R}}.$$

Since $\bar{\partial}^2\partial_t^4\breve{J}$ is in $H^{0.5}(\Omega)$, we use Lemma 2.5 to conclude that

$$\mathcal{I} = \dddot{\mathcal{R}}. \tag{7.42}$$

Using (7.42) in the time-integral of (7.41) completes the proof. ☐

We infer the following two propositions from the proof of Proposition 7.3:

Proposition 7.4 (Energy estimates for the action of $\bar{\partial}^3\partial_t^2$ in the Euler equations (7.2a))**.**

$$\sup_{t\in[0,T]} \|\bar{\partial}^3\breve{v}_{tt}(t)\|_0^2 + \sup_{t\in[0,T]} |\sqrt{\sigma}\bar{\partial}^3\breve{v}_t\cdot\breve{n}(t)|_1^2 + \sup_{t\in[0,T]} |\bar{\partial}^3\breve{v}_t\cdot\breve{n}(t)|_0^2 \le \int_0^T \dddot{\mathcal{R}}.$$

Proposition 7.5 (Energy estimates for the action of $\bar{\partial}^4$ in the Euler equations (7.2a))**.**

$$\sup_{t\in[0,T]} \|\bar{\partial}^4\breve{v}(t)\|_0^2 + \sup_{t\in[0,T]} |\sqrt{\sigma}\bar{\partial}^4\breve{\eta}\cdot\breve{n}(t)|_1^2 + \sup_{t\in[0,T]} |\bar{\partial}^4\breve{\eta}\cdot\breve{n}(t)|_0^2 \le \int_0^T \dddot{\mathcal{R}}.$$

Step 4: The σ-independent higher-order estimates via Proposition 2.1.

Lemma 7.3 (The σ-independent lower-order estimates for $\partial_t^a \breve{J}$, $a = 0, \dots, 7$)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{7} \|\partial_t^a \breve{J}(t)\|^2_{3.5-\frac{1}{2}a} \le \int_0^T \ddot{\dddot{\mathcal{R}}}.$$

Proof. The $a = 6$ and $a = 7$ cases are provided by Proposition 7.1. The higher-order estimates for $a = 0, \dots, 5$, are established by interpolation and the fundamental theorem of calculus. For example, using that $\partial_t^6 \breve{J}$ is in $H^1(\Omega)$

$$\sup_{t\in[0,T]} \|\partial_t^5 \breve{J}(t)\|^2_{1.5} \le C_\delta \sup_{t\in[0,T]} \|\partial_t^5 \breve{J}(t)\|^2_{1} + \delta \sup_{t\in[0,T]} \|\partial_t^5 \breve{J}(t)\|^2_{2} \le \int_0^T \ddot{\dddot{\mathcal{R}}}.$$

☐

Lemma 7.4 (The σ-independent normal trace-estimates for $\partial_t^a \breve{\eta}$, $a = 0, \dots, 7$)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{7} |\partial_t^a \breve{\eta}(t)\cdot N|^2_{4-\frac{1}{2}a} \le \int_0^T \ddot{\dddot{\mathcal{R}}}.$$

Proof. By Lemma 2.2,

$$\sup_{t\in[0,T]} |\bar\partial \partial_t^6 \breve{v}(t)\cdot N|^2_{-0.5} \le C \sup_{t\in[0,T]} \left(\|\bar\partial \partial_t^6 \breve{v}(t)\|^2_0 + \|\operatorname{div} \partial_t^6 \breve{v}(t)\|^2_0 \right) \le \int_0^T \ddot{\dddot{\mathcal{R}}},$$

thanks to the estimates stated in Proposition 7.2 and Lemma 7.3. Using the same argument, the $L^2(\Omega)$-estimates for $\bar\partial^3 \breve{v}_{tt}$ and $\bar\partial^4 \breve{v}$ respectively given in Propositions 7.4 and 7.5, and the divergence-estimates given by Lemma 7.3 provide that

$$\sup_{t\in[0,T]} \sum_{a=0}^{3} |\partial_t^{2a} \breve{v}(t)\cdot N|^2_{3.5-a} \le \int_0^T \ddot{\dddot{\mathcal{R}}}.$$

Using the fundamental theorem of calculus, the normal trace-estimates stated in Propositions 7.2, 7.3 and 7.5 complete the proof. ☐

Via Proposition 2.1, the estimates stated in Lemmas 7.2, 7.3 and 7.4 establish

Proposition 7.6 (The σ-independent estimates for $\partial_t^a \breve{\eta}$, $a = 0, \dots, 7$)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{7} \|\partial_t^a \breve{\eta}(t)\|^2_{4.5-\frac{1}{2}a} \le \int_0^T \ddot{\dddot{\mathcal{R}}}.$$

Via Proposition 2.1, the estimates stated in Lemma 7.2 and Proposition 7.2 establish

Proposition 7.7 (The σ-independent estimate for $\sqrt{\sigma}\partial_t^5 \breve{v}$)**.**

$$\|\sqrt{\sigma}\partial_t^5 \breve{v}(t)\|^2_2 \le \int_0^T \ddot{\dddot{\mathcal{R}}}.$$

Step 5: The σ-independent higher-order estimates via interpolation.

Proposition 7.8 (The σ-independent estimates for $\sqrt{\sigma}\partial_t^a\breve{\eta}$, $a = 0, \ldots, 3$)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{3} \|\sqrt{\sigma}\partial_t^a\breve{\eta}(t)\|^2_{5.5-\frac{1}{2}a} \leq \int_0^T \ddddot{\mathcal{R}}.$$

Proof. We note that by use of interpolation and Young's inequality,

$$\sup_{t\in[0,T]} \|\sqrt{\sigma}\breve{\eta}(t)\|^2_{5.5} \leq C_\delta \sup_{t\in[0,T]} \|\breve{\eta}(t)\|^2_{4.5} + \delta \sup_{t\in[0,T]} \|\sigma\breve{\eta}(t)\|^2_{5.5}.$$

It follows that the estimates given in Proposition 7.6 complete the proof. □

Step 6: The σ-independent higher-order estimates via the Euler equations (7.2a)**.**

Proposition 7.9 (The σ-independent estimates for $\partial_t^a\breve{J}$, $a = 0, \ldots, 5$)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{5} \|\partial_t^a\breve{J}(t)\|^2_{4.5-\frac{1}{2}a} \leq \int_0^T \ddddot{\mathcal{R}}.$$

Proof. We infer from the ath time-derivative of the Euler equations (7.2a) that

$$\sup_{t\in[0,T]} \|\partial_t^a\breve{J}(t)\|^2_{4.5-\frac{1}{2}a} \leq C \sup_{t\in[0,T]} \|\partial_t^a\breve{A}(t)\|^2_{3.5-\frac{1}{2}a} + C \sup_{t\in[0,T]} \|\partial_t^a\breve{v}_t(t)\|^2_{3.5-\frac{1}{2}a} + \int_0^T \ddddot{\mathcal{R}}.$$

The highest-order term of $\partial_t^a\breve{A}$ scales like $\breve{A}\breve{A}D\partial_t^a\breve{\eta}$. Hence, the estimates stated in Proposition 7.6 complete the proof. □

Given Proposition 7.9, we now establish

Proposition 7.10 (The σ-independent estimates for $\sqrt{\sigma}\partial_t^4\breve{v}$ and $\sqrt{\sigma}\breve{v}_{ttt}$ via Proposition 2.1)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{1} \|\sqrt{\sigma}\partial_t^{3+a}\breve{v}(t)\|^2_{3.5-\frac{1}{2}a} \leq \int_0^T \ddddot{\mathcal{R}}.$$

Proof. Via Proposition 2.1, the estimates stated in Lemma 7.2 and Propositions 7.3 and 7.9 establish the estimate for $\sqrt{\sigma}\breve{v}_{ttt}$. For $\sqrt{\sigma}\partial_t^4\breve{v}$, we have that

$$|\sqrt{\sigma}\partial_t^4\breve{v}\cdot\breve{n}(t)|^2_{2.5} \leq C_\delta|\partial_t^4\breve{v}\cdot\breve{n}(t)|^2_{1.5} + \delta|\sigma\partial_t^4\breve{v}\cdot\breve{n}(t)|^2_{3.5} \leq \int_0^T \ddddot{\mathcal{R}},$$

thanks to the estimate for $\partial_t^4\breve{v}$ stated in Proposition 7.6. Hence, we infer via Proposition 2.1 the estimate for $\sqrt{\sigma}\partial_t^4\breve{v}$ from the estimates stated in Lemma 7.2 and Proposition 7.9. □

Step 7: The σ-independent higher-order estimates for $\sigma\partial_t^a\breve{\eta}$, $a = 0, \ldots, 4$.

Lemma 7.5 (The σ-independent normal trace-estimates for $\sigma\partial_t^a\breve{\eta}$, $a = 0, \ldots, 4$)**.**

$$\sup_{t\in[0,T]} \sum_{a=0}^{3} |\sigma\bar{\partial}^2\partial_t^a\breve{\eta}\cdot\breve{n}(t)|^2_{4-\frac{1}{2}a} + \sup_{t\in[0,T]} |\sigma\partial_t^3\breve{v}\cdot\breve{n}(t)|^2_{3.5} \leq \int_0^T \ddddot{\mathcal{R}}.$$

Proof. The estimates for $\partial_t^a\breve{J}$ given in Proposition 7.9 and the fundamental theorem of calculus provide that the ath time-derivative of the Laplace-Young boundary condition (7.2b) yields the desired estimates. □

Proposition 7.11 (The σ-independent estimates for $\sigma\partial_t^a\breve{\eta}$, $a=0,\ldots,4$)**.**

$$\sup_{t\in[0,T]}\sum_{a=0}^{3}\|\sigma\partial_t^a\breve{\eta}(t)\|^2_{6.5-\frac{1}{2}a}+\sup_{t\in[0,T]}\|\sigma\partial_t^3\breve{v}(t)\|^2_4\le\int_0^T\dddot{\mathcal{R}}.$$

Proof. Since $\sigma<\sqrt{\sigma}$, we infer from the proof of Proposition 7.9 that the estimates for $\sqrt{\sigma}\partial_t^a\breve{v}$, $a=3,4,5$, stated in Propositions 7.7 and 7.10 establish the divergence-estimates for $\sigma\breve{v}_t$, $\sigma\breve{v}_{tt}$ and $\sigma\breve{v}_{ttt}$. Similarly, the estimates for $\sqrt{\sigma}\partial_t^a\breve{v}$, $a=1,2$, stated in Proposition 7.8 establish the divergence-estimates for $\sigma\breve{v}$ and $\sigma\breve{\eta}$. Via Proposition 2.1, the estimates stated in Lemmas 7.2 and 7.5 therefore complete the proof. ☐

Step 8: The σ-independent improved boundary-regularity estimates. Considering the action of ∂_t^a, $a=4,5,6$, in the Laplace-Young boundary condition (7.2b), we notice that the estimates stated in Propositions 7.9 and 7.11 establish

Proposition 7.12 (The σ-independent estimates for $\sigma\partial_t^a\breve{v}\cdot\breve{n}$, $a=3,4,5$)**.**

$$\sup_{t\in[0,T]}\sum_{a=0}^{1}|\sigma\partial_t^{3+a}\breve{v}\cdot\breve{n}(t)|^2_{4-\frac{1}{2}a}+\sup_{t\in[0,T]}|\sigma\partial_t^5\breve{v}\cdot\breve{n}(t)|^2_{2.5}\le\int_0^T\dddot{\mathcal{R}}.$$

Step 9: Concluding the proof of Lemma 7.1. The sum of the estimates given in Propositions 7.1–7.12 completes the proof of Lemma 7.1. Taking δ sufficiently small in the inequality (7.8) yields a polynomial-type inequality of the form (2.19). Hence, there exists $T>0$ that is independent of $\sigma>0$ and verifies

$$\sup_{t\in[0,T]}E^\sigma(t)\le 2\dddot{\mathcal{N}}_0. \tag{7.43}$$

7.4. **The proof of Theorem 1.2.**

7.4.1. *Existence.* Assuming that the initial data (ρ_0,u_0,Ω) is of C^∞-class, as in Section 7.1, we have that $M_0=P(E(0))<\infty$ where the higher-order energy function $E(t)$ is defined by (1.9). Since the difference of the compatibility conditions (1.14) and (7.4) is in $C^\infty(\Gamma)$, we repeat the proof of Theorem 1.1 to obtain the existence of a solution $\breve{v}$ to the σ-problem (7.2). The σ-independent estimate (7.43) and standard compactness arguments establish the strong convergence, as σ tends to zero,

$$\breve{\eta}\to\eta \text{ in } L^2(0,T;H^{3.5}(\Omega)),$$
$$\breve{v}_t\to v_t \text{ in } L^2(0,T;H^{2.5}(\Omega)).$$

Letting $\phi\in L^2(0,T;H^1(\Omega))$, we have that the variational form of (7.2) is

$$\int_0^T\int_\Omega\rho_0\breve{v}_t\cdot\phi-\int_0^T\int_\Omega\rho_0^2\breve{J}^{-2}\breve{a}_i^k\phi^i,_k+\int_0^T\int_\Gamma\beta_\sigma(t)\sqrt{\breve{g}}\phi\cdot\breve{n}=\sigma\int_0^T\int_\Gamma\sqrt{\breve{g}}\breve{g}^{\mu\nu}\breve{\eta},_{\mu\nu}\cdot\breve{n}\,\phi\cdot\breve{n}.$$

The strong convergence of the sequences $(\breve{\eta},\breve{v}_t)$ and the pointwise convergence $\beta_\sigma(t)\to\beta$ provide that the limit (η,v_t,β) satisfies

$$\int_0^T\int_\Omega\rho_0v_t\cdot\phi-\int_0^T\int_\Omega\rho_0^2J^{-2}a_i^k\phi^i,_k+\int_0^T\int_\Gamma\beta\sqrt{g}\phi\cdot n=0.$$

Thus, v is a solution of the zero surface tension limit of (1.7) on a nonempty time-interval $[0,T]$. Standard arguments provide that $v(0)=u_0$ and $\eta(0)=e$. Furthermore, letting $\sigma=0$ in the a priori estimates in Section 7.3, we conclude that the right-hand side of inequality

(7.43) depends only on $\mathcal{M}_0 = P(\mathcal{E}(0))$. Hence, for sufficiently small $T > 0$ the higher-order energy function $\mathcal{E}(t)$ defined in (1.11) satisfies

$$\sup_{t\in[0,T]} \mathcal{E}(t) \leq 2\mathcal{M}_0. \tag{7.44}$$

The bounds (7.7) remain valid by taking $T > 0$ even smaller if necessary. By the arguments in Section 6.1, the boundedness of J in assumption (7.7a) implies that

$$\rho(t) \geq \lambda \ \text{ in } \overline{\Omega}(t).$$

Similarly, the lower-bound (7.7b) provides that $\rho(t) = f \circ \eta^{-1}(t)$ satisfies

$$0 < \nu \leq -\frac{\partial \rho^2(t)}{\partial n(t)} \ \text{ on } \Gamma(t).$$

7.4.2. *Optimal regularity for the initial data.* In order to obtain the $H^{4.5}(\Omega)$-regularity of our existence theory, we assumed that the given initial data is of C^∞-class in Section 7.1. By virtue of the estimate (7.44), it in fact suffices for the regularity of the initial data to be such that $\mathcal{E}(0) < \infty$.

7.4.3. *Uniqueness.* We infer the uniqueness of the solution to the zero surface tension limit of (1.7) by repeating the arguments given in Section 6.3.

Acknowledgements. DC was supported by the Centre for Analysis and Nonlinear PDEs funded by the UK EPSRC grant EP/E03635X and the Scottish Funding Council. SS and JH were supported by the National Science Foundation under grant DMS-1001850.

Appendix A. Constructing C^∞-class initial data

We demonstrate how given initial data (ρ_0, u_0, Ω) of optimal regularity satisfying the conditions (1.13) and (1.14) (or the conditions (1.13), (1.15) and (1.16)) we are able to produce asymptotically consistent C^∞-class data $(\boldsymbol{\rho}_0, \boldsymbol{v}_0, \boldsymbol{\Omega})$ which satisfy similar statements of the conditions (1.13) and (1.14) (or the conditions (1.13), (1.15) and (1.16)).

A.1. **The C^∞-class data for the surface tension problem** (1.7)**.** We suppose the initial data (ρ_0, u_0, Ω) is of the optimal regularity stated in Theorem 1.1 and satisfy the conditions (1.13) and (1.14). Letting

$$\boldsymbol{J}_a = \partial_t^a (J^{-2})|_{t=0} \quad \text{and} \quad \boldsymbol{H}_a = \partial_t^a H(\eta)|_{t=0}, \tag{A.1}$$

we will construct C^∞-class data $(\boldsymbol{\rho}_0, \boldsymbol{v}_0, \boldsymbol{\Omega})$ which satisfy

$$\sigma \boldsymbol{H}_0 \geq 4\boldsymbol{\lambda}^2 - \beta \qquad \text{for } \sigma, \beta > 0, \tag{A.2a}$$

$$\boldsymbol{\rho}_0 \geq 2\boldsymbol{\lambda} > 0 \qquad \text{in } \overline{\boldsymbol{\Omega}}, \tag{A.2b}$$

$$\boldsymbol{\rho}_0^2 \boldsymbol{J}_a = \partial_t^a \beta + \sigma \boldsymbol{H}_a \quad \text{on } \boldsymbol{\Gamma} \ \text{ for } a = 0, 1, 2, 3. \tag{A.2c}$$

A.1.1. *Defining the C^∞-class initial domain $\boldsymbol{\Omega}$.* For $\boldsymbol{\epsilon} > 0$, we let $0 \leq \varphi^{\boldsymbol{\epsilon}} \in C_0^\infty(\mathbb{R}^3)$ denote the standard family of mollifiers with $\operatorname{spt}(\varphi^{\boldsymbol{\epsilon}}) \subset \overline{B(0,\boldsymbol{\epsilon})}$. Recalling the local coordinates near Γ defined in Section 2.1.5, we let $\boldsymbol{\Omega} \subset \mathbb{R}^3$ denote the C^∞-class domain defined by the C^∞-class charts

$$\boldsymbol{\theta}_l = \varphi^{\boldsymbol{\epsilon}} * [\mathcal{E}_{\mathcal{B}_l^{\boldsymbol{\epsilon}}}(\theta_l)] \;\text{ in } \mathcal{B}_l \;\text{ for } 1 \leq l \leq L,$$

where for $X \subset \mathbb{R}^3$ the operator $\mathcal{E}_X$ denotes a Sobolev extension operator mapping $H^s(X)$ to $H^s(\mathbb{R}^3)$ and where $\operatorname{spt}(\xi_l \circ \theta_l) \subset \mathcal{B}_l^{\boldsymbol{\epsilon}} \subset \mathcal{B}_l$. We assume that $\boldsymbol{\epsilon} > 0$ is taken sufficiently small so that the collection of open set $\{U_l\}_{l=1}^L$ introduced in Section 2.1.5 is an open covering of $\boldsymbol{\Omega}$. Setting $\boldsymbol{\Gamma} = \partial\boldsymbol{\Omega}$ defines a C^∞-class surface. We let $\boldsymbol{\eta}$ denote the $C^\infty(\boldsymbol{\Gamma})$-vector describing the geometry of $\boldsymbol{\Gamma}$. In other words, $\bar{\partial}\boldsymbol{\eta}$ spans the tangent space of $\boldsymbol{\Gamma}$ and, by letting $\boldsymbol{N}$ denote the outward-pointing unit normal vector to the surface $\boldsymbol{\Gamma}$, we have that $\boldsymbol{N} = \boldsymbol{\eta},_1 \times \boldsymbol{\eta},_2 / |\boldsymbol{\eta},_1 \times \boldsymbol{\eta},_2|$. We let $\mathbf{g}$ denote the surface metric induced by $\boldsymbol{\eta}$ and let $\boldsymbol{H}_0$ denote twice the mean curvature of $\boldsymbol{\Gamma}$. Thus,

$$\boldsymbol{H}_0 = -\mathbf{g}^{\alpha\beta}\boldsymbol{\eta},_{\alpha\beta} \cdot \boldsymbol{N} \text{ in } C^\infty(\boldsymbol{\Gamma}).$$

We assume that the given H^5-class domain $\Omega \subset \mathbb{R}^3$ is such that (1.14) is satisfied. Thus, for $\boldsymbol{\epsilon} > 0$ taken sufficiently small, standard properties of convolution provide that

$$\beta + \sigma\boldsymbol{H}_0 \geq 2\lambda^2 \;\text{ for } \sigma, \beta > 0. \tag{A.3}$$

Hence, (A.2a) is satisfied.

A.1.2. *Defining the C^∞-class data $(\boldsymbol{\varrho}_0, \boldsymbol{V}_0, \boldsymbol{\Omega})$ to satsify* (A.2) *for $a = 0$.* Given $u_0 \in H^4(\Omega)$, we define u_0 in the C^∞-class domain $\boldsymbol{\Omega}$ via the equation

$$\mathrm{u}_0 = \sum_{l=1}^{L} [\xi_l u_0 \circ \theta_l] \circ \boldsymbol{\theta}_l^{-1} \quad \text{ in } H^4(\boldsymbol{\Omega}). \tag{A.4}$$

Using the operator Λ_ϵ defined in Section 5.1, for $\boldsymbol{\epsilon} > 0$ we define the vector field $\boldsymbol{V}_0 \in C^\infty(\boldsymbol{\Omega})$ as the solution of the following elliptic Dirichlet problem:

$$\boldsymbol{V}_0 - \boldsymbol{\epsilon}\Delta\boldsymbol{V}_0 = \varphi^{\boldsymbol{\epsilon}} * \mathcal{E}_{\boldsymbol{\Omega}}(\mathrm{u}_0) \qquad \text{ in } \boldsymbol{\Omega}, \tag{A.5a}$$

$$\boldsymbol{V}_0 = \sum_{l=1}^{K} \Lambda_{\boldsymbol{\epsilon}}\big[\xi_l \mathrm{u}_0 \circ \boldsymbol{\theta}_l\big] \circ \boldsymbol{\theta}_l^{-1} \quad \text{ on } \boldsymbol{\Gamma}. \tag{A.5b}$$

Given $\rho_0 \in H^4(\Omega)$, we define ϱ_0 in the C^∞-class domain $\boldsymbol{\Omega}$ via the equation

$$\varrho_0 = \sum_{l=1}^{L} [\xi_l \rho_0 \circ \theta_l] \circ \boldsymbol{\theta}_l^{-1} \quad \text{ in } H^4(\boldsymbol{\Omega}). \tag{A.6}$$

Assuming that $\rho_0 \geq 2\lambda > 0$ in $\overline{\Omega}$, we let $\boldsymbol{\lambda} > 0$ be such that $2\lambda^2 \geq 9\boldsymbol{\lambda}^2$. It follows for ϱ_0 defined by (A.6) that

$$\varrho_0 \geq 3\boldsymbol{\lambda} \;\text{ in } \overline{\boldsymbol{\Omega}}. \tag{A.7}$$

For $\boldsymbol{\mu} > 0$ and $\boldsymbol{\epsilon} > 0$, we define $\boldsymbol{\varrho}_0 \in C^\infty(\boldsymbol{\Omega})$ to be the solution of

$$\boldsymbol{\varrho}_0 - \boldsymbol{\mu}\Delta\boldsymbol{\varrho}_0 = \varphi^{\boldsymbol{\mu}} * \mathcal{E}_{\boldsymbol{\Omega}}(\varrho_0) \quad \text{ in } \boldsymbol{\Omega}, \tag{A.8a}$$

$$\boldsymbol{\varrho}_0 = \sqrt{\beta + \sigma\boldsymbol{H}_0} \quad \text{ on } \boldsymbol{\Gamma}. \tag{A.8b}$$

The boundary condition (A.8b) implies for $a = 0$ that (A.2c) is satisfied, since (A.8b) is equivalently stated as

$$\boldsymbol{\varrho}_0^2 \boldsymbol{J}_0 = \beta + \sigma \boldsymbol{H}_0 \quad \text{on } \boldsymbol{\Gamma}. \tag{A.9}$$

Elliptic regularity provides for $s \geq 2$ and a positive constant C independent of $\boldsymbol{\mu}$ that

$$\|\sqrt{\boldsymbol{\mu}}\boldsymbol{\varrho}_0\|^2_{H^s(\boldsymbol{\Omega})} \leq C\Big(\|\varphi^{\boldsymbol{\mu}} * \mathcal{E}_{\boldsymbol{\Omega}}(\varrho_0)\|^2_{H^{s-2}(\boldsymbol{\Omega})} + \|\boldsymbol{\varrho}_0\|^2_{H^{s-\frac{1}{2}}(\boldsymbol{\Gamma})}\Big).$$

The boundary forcing function used in defining $\boldsymbol{\varrho}_0$ is in $C^\infty(\boldsymbol{\Gamma})$. Since the function ϱ_0 appearing in the right-hand side of (A.8a) is in $H^4(\boldsymbol{\Omega})$, there exists a constant C which is independent of $\boldsymbol{\mu}$ verifying

$$\|\sqrt{\boldsymbol{\mu}}\boldsymbol{\varrho}_0\|^2_{H^6(\boldsymbol{\Omega})} \leq C.$$

The equation (A.8a) provides that $\boldsymbol{\varrho}_0 = \boldsymbol{\mu}\Delta\boldsymbol{\varrho}_0 + \varphi^{\boldsymbol{\mu}} * \mathcal{E}_{\boldsymbol{\Omega}}(\varrho_0)$ in $\boldsymbol{\Omega}$. Hence, by standard properties of convolution and taking $\boldsymbol{\mu}$ sufficiently small, we conclude that

$$\|\boldsymbol{\varrho}_0 - \varrho_0\|_{L^\infty(\boldsymbol{\Omega})} \leq \sqrt{\boldsymbol{\mu}}\|\sqrt{\boldsymbol{\mu}}\Delta\boldsymbol{\varrho}_0\|_{H^2(\boldsymbol{\Omega})} + \|\varphi^{\boldsymbol{\mu}} * \mathcal{E}_{\boldsymbol{\Omega}}(\varrho_0) - \varrho_0\|_{L^\infty(\boldsymbol{\Omega})} \leq \boldsymbol{\lambda}.$$

Recalling the lower-bound (A.7), we conclude that $\boldsymbol{\varrho}_0 \geq 2\boldsymbol{\lambda} > 0$ in $\boldsymbol{\Omega}$. Letting $\boldsymbol{\epsilon}$ be sufficiently small, the boundary condition (A.8b) provides that

$$\boldsymbol{\varrho}_0 \geq 2\boldsymbol{\lambda} > 0 \ \text{ in } \overline{\boldsymbol{\Omega}}. \tag{A.10}$$

By (A.10) and (A.3), the boundary condition (A.8b) provides that

$$\beta + \sigma\boldsymbol{H}_0 \geq 4\boldsymbol{\lambda}^2 \ \text{ for } \sigma, \beta > 0. \tag{A.11}$$

Hence, the C^∞-class data $(\boldsymbol{\varrho}_0, \boldsymbol{V}_0, \boldsymbol{\Omega})$ satisfy the conditions (A.2) for $a = 0$, as verified by (A.9), (A.10) and (A.11).

A.1.3. *Formal definitions.* We formally set $\boldsymbol{\rho}_0$ equal to $\boldsymbol{\varrho}_0$ defined by (A.8). We define

$$\boldsymbol{v}_{a+1} = -2\partial_t^a\big(A^k_\cdot(\boldsymbol{\rho}_0 J^{-1})\big)|_{t=0} \quad \text{for } a = 0, \dots, 5, \tag{A.12}$$

and assume that (A.12) yields $\boldsymbol{v}_0 = \boldsymbol{V}_0$ defined by (A.5). We make the following definitions:

$$\boldsymbol{j}_a = \operatorname{div}\boldsymbol{v}_a \quad \text{and} \quad \boldsymbol{k}_a = \partial_t^a(A^s_r v^r,_s)|_{t=0} \quad \text{for } a = 0, \dots, 6. \tag{A.13}$$

Using the notation

$$[\boldsymbol{A}_a]^s_r = \partial_t^a A^s_r|_{t=0} \quad \text{for } a = 0, \dots, 5, \tag{A.14}$$

it follows that $\boldsymbol{k}_a$ defined in (A.13) satisfies

$$\boldsymbol{k}_0 = \boldsymbol{j}_0, \quad \boldsymbol{k}_1 = [\boldsymbol{A}_1]^s_r \boldsymbol{v}^r_0,_s + \boldsymbol{j}_1 \ \text{ and } \ \boldsymbol{k}_2 = [\boldsymbol{A}_2]^s_r \boldsymbol{v}^r_0,_s + 2[\boldsymbol{A}_1]^s_r \boldsymbol{v}^r_1,_s + \boldsymbol{j}_2. \tag{A.15}$$

We also have that $\boldsymbol{J}_a$ defined by (A.1) is equivalently given by

$$\boldsymbol{J}_0 = 1, \quad \boldsymbol{J}_1 = -2\boldsymbol{k}_0, \quad \boldsymbol{J}_2 = 4\boldsymbol{k}_0^2 - 2\boldsymbol{k}_1 \ \text{ and } \ \boldsymbol{J}_3 = -8\boldsymbol{k}_0^3 + 12\boldsymbol{k}_0\boldsymbol{k}_1 - 2\boldsymbol{k}_2. \tag{A.16}$$

Using (A.15) and (A.16), the condition (A.2c) that $\boldsymbol{\rho}_0^2\boldsymbol{J}_a = \sigma\boldsymbol{H}_a$ is equivalently stated as

$$\operatorname{div}\boldsymbol{v}_{a-1} = \boldsymbol{j}_{a-1} = \underbrace{-\frac{1}{2}\big[\sigma\boldsymbol{H}_a - \boldsymbol{\rho}_0^2\boldsymbol{J} - 2\boldsymbol{k}_{a-1}\big] + \boldsymbol{d}_{a-1}}_{\boldsymbol{f}_{a-1}}, \quad \text{for } a = 1, 2, 3, \tag{A.17}$$

where $\boldsymbol{d}_{a-1}$ represents the lower-order terms defining $\boldsymbol{k}_{a-1}$. We notice that $\boldsymbol{v}_a$, $a = 1, 2$, defined by (A.12) is equivalently given by $\boldsymbol{v}_1 = -2D\boldsymbol{\rho}_0$ and $\boldsymbol{v}_2 = -2[\boldsymbol{A}_1]^k_\cdot \boldsymbol{\rho}_0,_k - 2D(\boldsymbol{\rho}_0\boldsymbol{j}_0)$. Thus,

$$\operatorname{div}\boldsymbol{v}_0 = \boldsymbol{f}_0, \quad \operatorname{div}\boldsymbol{v}_1 = \boldsymbol{f}_1 = -2\Delta\boldsymbol{\rho}_0, \quad \text{and} \quad \operatorname{div}\boldsymbol{v}_2 = \boldsymbol{f}_2 \approx -2\Delta\operatorname{div}\boldsymbol{v}_0. \tag{A.18}$$

A.1.4. *Defining* $\boldsymbol{v}_0$ *so that the* C^∞*-class data* $(\boldsymbol{\varrho}_0, \boldsymbol{v}_0, \boldsymbol{\Omega})$ *satisfy* (A.2) *for* $a = 0, 1$. Extending $\boldsymbol{N}$ and $\boldsymbol{\tau}_\alpha = \boldsymbol{\eta},_\alpha / |\boldsymbol{\eta},_\alpha|$, for $\alpha = 1, 2$, into $\boldsymbol{\Omega}$, we decompose any vector $\xi \in \mathbb{R}^3$ into normal and tangential components as

$$\xi = \xi^\alpha \boldsymbol{\tau}_\alpha + \xi^3 \boldsymbol{N} \quad \text{in } \boldsymbol{\Omega}.$$

It follows for sufficiently regular ξ that $\operatorname{div} \xi = \xi^i,_i$ is equivalently written as

$$\operatorname{div} \xi = \underbrace{\xi^\alpha,_\alpha + \xi^3,_3 + \xi^\alpha \operatorname{div} \boldsymbol{\tau}_\alpha + \xi^3 \operatorname{div} \boldsymbol{N}}_{j_0} \quad \text{on } \boldsymbol{\Gamma}. \tag{A.19}$$

This provides the following identity for $(\boldsymbol{N} \cdot D)\xi^3 = \xi^3,_3$:

$$\frac{\partial \xi^3}{\partial \boldsymbol{N}} = \underbrace{j_0 - \left[\xi^\alpha,_\alpha + \xi^\alpha \operatorname{div} \boldsymbol{\tau}_\alpha + \xi^3 \operatorname{div} \boldsymbol{N}\right]}_{\varphi_0} \quad \text{on } \boldsymbol{\Gamma}. \tag{A.20}$$

We define $\boldsymbol{v}_0^3 \in C^\infty(\boldsymbol{\Omega})$ by

$$\boldsymbol{v}_0^3 + \boldsymbol{\epsilon} \Delta^2 \boldsymbol{v}_0^3 = \varphi^{\boldsymbol{\epsilon}} * \mathcal{E}_{\boldsymbol{\Omega}}(\mathrm{u}_0^3) \quad \text{in } \boldsymbol{\Omega}, \tag{A.21a}$$

$$\frac{\partial \boldsymbol{v}_0^3}{\partial \boldsymbol{N}} = \boldsymbol{\varphi}_0 \quad \text{on } \boldsymbol{\Gamma}, \tag{A.21b}$$

$$\boldsymbol{v}_0^3 = \boldsymbol{V}_0^3 \quad \text{on } \boldsymbol{\Gamma}, \tag{A.21c}$$

where the boundary forcing function $\boldsymbol{\varphi}_0$ is defined as

$$\boldsymbol{\varphi}_0 = \boldsymbol{f}_0 - [\boldsymbol{V}_0^\alpha,_\alpha + \boldsymbol{V}_0^\alpha \operatorname{div} \boldsymbol{\tau}_\alpha + \boldsymbol{V}_0^3 \operatorname{div} \boldsymbol{N}], \tag{A.22}$$

with the function $\boldsymbol{f}_0$ appearing in the right-hand side of (A.22) defined by (A.17). With $\boldsymbol{V}_0^\alpha$ denoting the tangential component of the vector defined by (A.5), we define the vector $\boldsymbol{v}_0 \in C^\infty(\Omega)$ as

$$\boldsymbol{v}_0 = \boldsymbol{V}_0^\alpha \boldsymbol{\tau}_\alpha + \boldsymbol{v}_0^3 \boldsymbol{N} \quad \text{in } \boldsymbol{\Omega}. \tag{A.23}$$

The boundary condition (A.21c) and the definition (A.23) imply that

$$\boldsymbol{v}_0 = \boldsymbol{V}_0 \quad \text{on } \boldsymbol{\Gamma}. \tag{A.24}$$

Thanks to (A.19), the boundary condition (A.21b) and the definition (A.22) yield

$$\operatorname{div} \boldsymbol{v}_0 = \boldsymbol{f}_0. \tag{A.25}$$

Thanks to the definition (A.17) of $\boldsymbol{f}_0$, we equivalently write (A.25) as

$$\boldsymbol{\varrho}_0^2 \boldsymbol{J}_1 = \sigma \boldsymbol{H}_1 \quad \text{on } \boldsymbol{\Gamma}. \tag{A.26}$$

Hence, the C^∞-class data $(\boldsymbol{\varrho}_0, \boldsymbol{v}_0, \boldsymbol{\Omega})$ satisfy the conditions (A.2) for $a = 0, 1$, as verified by (A.11), (A.9), (A.10) and (A.26).

A.1.5. *Defining the* C^∞*-class data* $(\boldsymbol{\rho}_0, \boldsymbol{v}_0, \boldsymbol{\Omega})$ *to satisfy* (A.2) *for* $a = 0, 1, 2$. According to (A.18), the condition (A.2c) for $a = 2$ may be imposed by prescribing a Dirichlet boundary condition for $\Delta \boldsymbol{\rho}_0$. We define $\boldsymbol{\rho}_0 \in C^\infty(\boldsymbol{\Omega})$ to be the solution of the polyharmonic problem

$$\boldsymbol{\rho}_0 - \boldsymbol{\mu} \Delta^3 \boldsymbol{\rho}_0 = \varphi^{\boldsymbol{\mu}} * \mathcal{E}_{\boldsymbol{\Omega}}(\varrho_0) \quad \text{in } \boldsymbol{\Omega}, \tag{A.27a}$$

$$-2\Delta \boldsymbol{\rho}_0 = \boldsymbol{f}_1 \quad \text{on } \boldsymbol{\Gamma}, \tag{A.27b}$$

$$-2\frac{\partial \boldsymbol{\rho}_0}{\partial \boldsymbol{N}} = \boldsymbol{V}_1^3 \quad \text{on } \boldsymbol{\Gamma}, \tag{A.27c}$$

$$\boldsymbol{\rho}_0 = \boldsymbol{\varrho}_0 \quad \text{on } \boldsymbol{\Gamma}. \tag{A.27d}$$

The boundary-forcing function $\boldsymbol{f}_1$ appearing in the right-hand side of (A.27b) is defined as

$$\boldsymbol{f}_1 = -\frac{\sigma \boldsymbol{H}_2}{2\boldsymbol{\varrho}_0^2} + 2\boldsymbol{k}_0^2 - [\boldsymbol{\mathcal{A}}_1]^s_r \boldsymbol{v}^r_{0},_s \,. \tag{A.28}$$

Using the boundary conditions of the polyharmonic problem (A.21), we may express the right-hand side of (A.28) in terms of $\boldsymbol{V}_0$, $\boldsymbol{V}_1$ and $\boldsymbol{\varphi}_0$, where

$$\boldsymbol{V}_1 = -2D\boldsymbol{\varrho}_0 \quad \text{on } \boldsymbol{\Gamma}. \tag{A.29}$$

The boundary-forcing function $\boldsymbol{V}_1^3$ appearing in (A.27c) is the normal component of $\boldsymbol{V}_1$ and the function $\boldsymbol{\varrho}_0$ is the solution of the elliptic Dirichlet problem (A.8). The boundary conditions (A.27c) and (A.27d) imply that $\boldsymbol{v}_1 = -2D\boldsymbol{\rho}_0$ defined by (A.12) satisfies

$$\boldsymbol{v}_1 = \boldsymbol{V}_1 \quad \text{on } \boldsymbol{\Gamma}. \tag{A.30}$$

Polyharmonic regularity provides for $s \geq 0$ a positive constant C independent of $\boldsymbol{\mu}$ that

$$\begin{aligned} &\|\sqrt{\boldsymbol{\mu}}\boldsymbol{\rho}_0\|^2_{H^{s+6}(\boldsymbol{\Omega})} \\ &\quad \leq C\Big(\|\varphi^{\boldsymbol{\mu}} * \boldsymbol{\mathcal{E}}_{\boldsymbol{\Omega}}(\varrho_0)\|^2_{H^s(\boldsymbol{\Omega})} + \|\Delta\boldsymbol{\rho}_0\|^2_{H^{s+3.5}(\boldsymbol{\Gamma})} + \left\|\frac{\partial\boldsymbol{\rho}_0}{\partial\boldsymbol{N}}\right\|^2_{H^{s+4.5}(\boldsymbol{\Gamma})} + \|\boldsymbol{\rho}_0\|^2_{H^{s+5.5}(\boldsymbol{\Gamma})}\Big). \end{aligned}$$

Recalling Section A.1.2, the arguments establishing the lower-bound (A.10) provide that

$$\boldsymbol{\rho}_0 \geq 2\boldsymbol{\lambda} > 0 \ \text{ in } \overline{\boldsymbol{\Omega}}. \tag{A.31}$$

Similarly, we infer from the arguments given in Section A.1.4 that the boundary condition (A.27b) and the identity (A.30) establish that

$$\boldsymbol{\rho}_0^2\boldsymbol{J}_2 = \sigma\boldsymbol{H}_2 \quad \text{on } \boldsymbol{\Gamma}. \tag{A.32}$$

Hence, the C^∞-class data $(\boldsymbol{\rho}_0, \boldsymbol{v}_0, \boldsymbol{\Omega})$, where $\boldsymbol{\rho}_0$ is the solution of (A.27) and $\boldsymbol{v}_0$ is given by (A.23), satisfy the conditions (A.2) for $a = 0, 1, 2$, as verified by (A.11), (A.9), (A.26), (A.31) and (A.32).

Remark 22. According to (A.18), the condition (A.2c) for $a = 3$ may be imposed by prescribing a Dirichlet boundary condition for $\frac{\partial}{\partial \boldsymbol{N}}\Delta\boldsymbol{v}_0^3$ in a polyharmonic problem for $\boldsymbol{v}_0^3$ satisfying $\boldsymbol{v}_0^3 + \boldsymbol{\epsilon}\Delta^4\boldsymbol{v}_0^3 = \varphi^{\boldsymbol{\epsilon}} * \boldsymbol{\mathcal{E}}_{\boldsymbol{\Omega}}(\mathrm{u}_0^3)$.

A.2. The C^∞-class data for the zero surface tension limit of (1.7).

We suppose the initial data (ρ_0, u_0, Ω) is of the optimal regularity stated in Theorem 1.2 and satisfy the conditions (1.13), (1.15) and (1.16). Then setting $\sigma = 0$ in the construction of the C^∞-class data of Section A.1, $\boldsymbol{J}_a = \partial_t^a(J^{-2})|_{t=0}$ satisfies

$$\boldsymbol{J}_0 = \beta, \quad \boldsymbol{J}_1 = 0, \quad \boldsymbol{J}_2 = 0 \ \text{ and } \ \boldsymbol{J}_3 = 0 \quad \text{on } \boldsymbol{\Gamma}.$$

For $\boldsymbol{\mu} > 0$, and with ϱ_0 defined by (A.6), $\boldsymbol{\varrho}_0$ solving (A.8), we define $\boldsymbol{\rho}_0 \in C^\infty(\boldsymbol{\Omega})$ to be the solution of the polyharmonic problem

$$\begin{aligned} \boldsymbol{\rho}_0 - \boldsymbol{\mu}\Delta^5\boldsymbol{\rho}_0 &= \varphi^{\boldsymbol{\mu}} * \boldsymbol{\mathcal{E}}_{\boldsymbol{\Omega}}(\varrho_0) && \text{in } \boldsymbol{\Omega}, && \text{(A.33a)} \\ \Delta^2\boldsymbol{\rho}_0 &= \boldsymbol{f}_3 && \text{on } \boldsymbol{\Gamma}, && \text{(A.33b)} \\ \frac{\partial\Delta\boldsymbol{\rho}_0}{\partial\boldsymbol{N}} &= \frac{\partial\Delta\boldsymbol{\varrho}_0}{\partial\boldsymbol{N}} && \text{on } \boldsymbol{\Gamma}, && \text{(A.33c)} \\ -2\Delta\boldsymbol{\rho}_0 &= \boldsymbol{f}_1 && \text{on } \boldsymbol{\Gamma}, && \text{(A.33d)} \\ -2\frac{\partial\boldsymbol{\rho}_0}{\partial\boldsymbol{N}} &= \boldsymbol{V}_1^3 && \text{on } \boldsymbol{\Gamma}, && \text{(A.33e)} \\ \boldsymbol{\rho}_0 &= \sqrt{\beta} && \text{on } \boldsymbol{\Gamma}. && \text{(A.33f)} \end{aligned}$$

where the boundary-forcing function $\boldsymbol{f}_3$ is defined below in (A.36). By Section A.1.5,

$$\boldsymbol{\rho}_0 \geq \boldsymbol{\lambda} > 0 \quad \text{in } \overline{\boldsymbol{\Omega}}. \tag{A.34}$$

According to the definition (A.12) of $\boldsymbol{v}_a$,

$$\boldsymbol{v}_3 = -2[\boldsymbol{\mathcal{A}}_2]^k_\cdot \boldsymbol{\rho}_0,_k -4[\boldsymbol{\mathcal{A}}_1]^k_\cdot \big(\boldsymbol{\rho}_0 \boldsymbol{K}_0\big),_k -2D\big(\boldsymbol{\rho}_0 \boldsymbol{K}_1\big) \quad \text{in } \boldsymbol{\Omega},$$

where $\boldsymbol{K}_a = \partial_t^a\big(J^{-1}(J^{-1}J_t)\big)|_{t=0} = \partial_t^a\big(J^{-1}(A^s_r v^r,_s)\big)|_{t=0}$. We compute

$$\boldsymbol{J}_4 = 16\boldsymbol{k}_0^4 - 48\boldsymbol{k}_0^2\boldsymbol{k}_1 + 12\boldsymbol{k}_1^2 + 16\boldsymbol{k}_0\boldsymbol{k}_2 - 2\boldsymbol{k}_3.$$

Setting $\boldsymbol{J}_4 = 0$ yields an equation for $\boldsymbol{k}_3$. Since $\boldsymbol{k}_3 = [\boldsymbol{\mathcal{A}}_2]^s_r \boldsymbol{v}^r_0,_s + 3[\boldsymbol{\mathcal{A}}_2]^s_r \boldsymbol{v}^r_1,_s + 3[\boldsymbol{\mathcal{A}}_1]^s_r \boldsymbol{v}^r_2,_s + \boldsymbol{j}_3$,

$$\boldsymbol{j}_3 = \frac{1}{2}\Big[16\boldsymbol{k}_0^4 - 48\boldsymbol{k}_0^2\boldsymbol{k}_1 + 12\boldsymbol{k}_1^2 + 16\boldsymbol{k}_0\boldsymbol{k}_2\Big] - \Big[[\boldsymbol{\mathcal{A}}_2]^s_r \boldsymbol{v}^r_0,_s + 3[\boldsymbol{\mathcal{A}}_2]^s_r \boldsymbol{v}^r_1,_s + 3[\boldsymbol{\mathcal{A}}_1]^s_r \boldsymbol{v}^r_2,_s\Big].$$

On the other hand, the divergence of $\boldsymbol{v}_3 = -2[\boldsymbol{\mathcal{A}}_2]^k_\cdot \boldsymbol{\rho}_0,_k -4[\boldsymbol{\mathcal{A}}_1]^k_\cdot \big(\boldsymbol{\rho}_0 \boldsymbol{K}_0\big),_k -2D\big(\boldsymbol{\rho}_0 \boldsymbol{K}_1\big)$ yields

$$\operatorname{div} \boldsymbol{v}_3 = \operatorname{div}\big[-2[\boldsymbol{\mathcal{A}}_2]^k_\cdot \boldsymbol{\rho}_0,_k -4[\boldsymbol{\mathcal{A}}_1]^k_\cdot \big(\boldsymbol{\rho}_0 \boldsymbol{K}_0\big),_k -2D\boldsymbol{\rho}_0 \boldsymbol{K}_1\big] - 2\boldsymbol{\rho}_0,_i \boldsymbol{K}_1,_i - 2\boldsymbol{\rho}_0 \Delta \boldsymbol{K}_1.$$

We solve for $\Delta \boldsymbol{K}_1$ to write

$$\Delta \boldsymbol{K}_1 = \underbrace{\frac{1}{2\boldsymbol{\rho}_0}\Big[-\boldsymbol{j}_3 + \operatorname{div}\big[-2[\boldsymbol{\mathcal{A}}_2]^k_\cdot \boldsymbol{\rho}_0,_k -4[\boldsymbol{\mathcal{A}}_1]^k_\cdot \big(\boldsymbol{\rho}_0 \boldsymbol{K}_0\big),_k -2D\boldsymbol{\rho}_0 \boldsymbol{K}_1\big] - 2\boldsymbol{\rho}_0,_i \boldsymbol{K}_1,_i\Big]}_{\mathfrak{J}_3}.$$

Using that $\boldsymbol{K}_1 = -\boldsymbol{k}_0^2 + \boldsymbol{k}_1$ and $\boldsymbol{k}_1 = [\boldsymbol{\mathcal{A}}_1]^s_r \boldsymbol{v}^r_0,_s + \boldsymbol{j}_1$, we find that

$$\Delta \boldsymbol{j}_1 = \mathfrak{J}_3 + \Delta \boldsymbol{k}_0^2 - \Delta([\boldsymbol{\mathcal{A}}_1]^s_r \boldsymbol{v}^r_0,_s). \tag{A.35}$$

Since $\boldsymbol{v}_1 = -2D\boldsymbol{\rho}_0$ implies that $\boldsymbol{j}_1 = -2\Delta\boldsymbol{\rho}_0$ we equivalently write (A.35) as

$$\Delta^2 \boldsymbol{\rho}_0 = \underbrace{-\frac{1}{2}\big[\mathfrak{J}_3 + \Delta \boldsymbol{k}_0^2 - \Delta([\boldsymbol{\mathcal{A}}_1]^s_r \boldsymbol{v}^r_0,_s)\big]}_{\boldsymbol{f}_3} \quad \text{on } \boldsymbol{\Gamma}. \tag{A.36}$$

The quantity $D\boldsymbol{K}_1$ may be expressed using the boundary conditions (A.33c)–(A.33f) and $D^a \boldsymbol{v}_0$, $a = 0, 1, 2, 3$ may be expressed using the boundary conditions of the polyharmonic problems defining $\boldsymbol{v}_0$. It follows from (A.12) and (A.36) that $\boldsymbol{J}_a = \partial_t^a(J^{-2})|_{t=0}$ satisfies

$$\boldsymbol{J}_a = \partial_t^a \beta \quad \text{on } \boldsymbol{\Gamma} \text{ for } a = 0, 1, 2, 3, 4. \tag{A.37}$$

With $0 < 2\nu \leq -\frac{\partial \rho_0^2}{\partial N}$, we may take $\boldsymbol{\epsilon}$ sufficiently small so that $\boldsymbol{\varrho}_0$ defined in (A.8) satisfies

$$0 < \nu \leq -\frac{\partial \boldsymbol{\varrho}_0^2}{\partial \boldsymbol{N}} \quad \text{on } \boldsymbol{\Gamma}.$$

We then have that (A.29) defining the vector field $\boldsymbol{V}_1 \in C^\infty(\boldsymbol{\Gamma})$ is

$$\boldsymbol{V}_1 = -\frac{\partial \boldsymbol{\varrho}_0^2}{\partial \boldsymbol{N}} \boldsymbol{N} \quad \text{on } \boldsymbol{\Gamma}.$$

The boundary condition (A.33e) of the polyharmonic problem (A.33) defining $\boldsymbol{\rho}_0$ provides that

$$0 < \nu \leq -\frac{\partial \boldsymbol{\rho}_0^2}{\partial \boldsymbol{N}} \quad \text{on } \boldsymbol{\Gamma}.$$

The conditions (A.33e), (A.34) and (A.37) establish that the C^∞-class initial data $(\boldsymbol{\rho}_0, \boldsymbol{v}_0, \boldsymbol{\Omega})$ satisfy

$$\boldsymbol{\rho}_0 \geq \boldsymbol{\lambda} > 0 \quad \text{in } \overline{\boldsymbol{\Omega}}, \tag{A.38a}$$

$$-\frac{\partial \boldsymbol{\rho}_0^2}{\partial \boldsymbol{N}} \geq \nu > 0 \quad \text{on } \boldsymbol{\Gamma}, \tag{A.38b}$$

$$\boldsymbol{\rho}_0^2 \boldsymbol{J}_a = \partial_t^a \beta \qquad \text{on } \boldsymbol{\Gamma} \quad \text{for } a = 0, 1, 2, 3, 4. \tag{A.38c}$$

The conditions (A.38) are analogous statements of the conditions (1.13), (1.15) and (1.16).

Remark 23. According to (A.12),

$$\operatorname{div} \boldsymbol{v}_4 = \operatorname{div} \big[-2[\boldsymbol{\mathcal{A}}_3]_\cdot^k \boldsymbol{\rho}_0,_k -6[\boldsymbol{\mathcal{A}}_2]_\cdot^k (\boldsymbol{\rho}_0 \boldsymbol{K}_0),_k -2[\boldsymbol{\mathcal{A}}_1]_\cdot^k (\boldsymbol{\rho}_0 \boldsymbol{K}_1),_k -2D\boldsymbol{\rho}_0 \boldsymbol{K}_2 \big] - 2\boldsymbol{\rho}_0,_i \boldsymbol{K}_2,_i \\ -2\boldsymbol{\rho}_0 \Delta \boldsymbol{K}_2.$$

Since $\Delta \boldsymbol{K}_2$ is approximately $\Delta^2 \operatorname{div} \boldsymbol{v}_0$, the boundary condition $\boldsymbol{J}_5 = 0$ may be imposed by prescribing a Dirichlet boundary condition for $\frac{\partial}{\partial \boldsymbol{N}} \Delta^2 \boldsymbol{v}_0^3$ in a polyharmonic problem for $\boldsymbol{v}_0^3$ satisfying $\boldsymbol{v}_0^3 + \boldsymbol{\epsilon} \Delta^6 \boldsymbol{v}_0^3 = \varphi^{\boldsymbol{\epsilon}} * \mathcal{E}_{\boldsymbol{\Omega}}(\mathrm{u}_0^3)$.

The boundary condition $\boldsymbol{J}_6 = 0$ is obtained by defining a Dirichlet boundary condition for $\Delta^3 \boldsymbol{\rho}_0$ in a polyharmonic problem for $\boldsymbol{\rho}_0$ satisfying $\boldsymbol{\rho}_0 - \boldsymbol{\mu} \Delta^7 \boldsymbol{\rho}_0 = \varphi^{\boldsymbol{\mu}} * \mathcal{E}_{\boldsymbol{\Omega}}(\varrho_0)$.

Appendix B. Solutions to the μ-problem (5.52)

In this appendix, we construct a fixed-point solution $\mathring{v}$ to the μ-problem (5.52):

Proposition B.1 (Solutions to the μ-problem)**.** *For C^∞-class initial data (ρ_0, u_0, Ω) satisfying the conditions* (1.13) *and* (1.14)*, and for some $T = T_\kappa(\epsilon\mu) > 0$, there exists a unique $\mathring{v} \in L^2(0,T;H^9(\Omega))$ solving the μ-problem* (5.52) *on a time-interval $[0,T]$, with $\partial_t^a \mathring{v} \in L^2(0,T;H^{9-2a}(\Omega))$ for $a = 1,2,3,4$, $\mathring{v}_{tttt} \in L^\infty(0,T;L^2(\Omega))$ and $(\mathring{v}, \mathring{v}_t, \ldots, \mathring{v}_{tttt})|_{t=0} = (u_0, \mathbf{v}_1, \ldots, \mathbf{v}_4)$.*

Remark 24. We recall that the initial data $\mathbf{v}_a$, $a = 1,2,3,4$, is defined in Section 5.2.1.

B.1. **The functional framework for the fixed-point scheme.** To establish the existence and uniqueness of solutions to the μ-problem (5.52), we define for $T > 0$ the Hilbert space

$$\mathbf{X}_T = \{ v \in L^2(0,T;H^5(\Omega)) \mid \partial_t^a v \in L^2(0,T;H^{5-2a}(\Omega)) \text{ for } a = 1,2 \},$$

endowed with the natural Hilbert norm

$$\|v\|_{\mathbf{X}_T}^2 = \sum_{a=0}^{2} \|\partial_t^a v\|_{L^2(0,T;H^{5-2a}(\Omega))}^2.$$

For $\mathcal{M} > 0$ (where the particular value of $\mathcal{M}$ is specified later), we define the closed, bounded, convex subset $\mathcal{C}_T(\mathcal{M})$ of $\mathbf{X}_T$

$$v \in \mathcal{C}_T(\mathcal{M}) \subset \mathbf{X}_T \tag{B.1}$$

to be all $v \in \mathbf{X}_T$ that satisfy each of the following conditions:

- (X1) $(v, v_t, v_{tt})|_{t=0} = (u_0, \mathbf{v}_1, \mathbf{v}_2)$, and
- (X2) $\|v\|_{\mathbf{X}_T}^2 \leq \mathcal{M}$.

Lemma B.1 (Solutions to the μ-problem in $\mathbf{X}_T$)**.** *For C^∞-class initial data (ρ_0, u_0, Ω) satisfying the conditions* (1.13) *and* (1.14)*, and for some $T = T_\kappa(\epsilon\mu) > 0$, there exists a unique $\mathring{v} \in \mathbf{X}_T$ that solves the μ-problem* (5.52) *on a time-interval $[0,T]$ and verifies $(\mathring{v}, \mathring{v}_t, \mathring{v}_{tt})|_{t=0} = (u_0, \mathbf{v}_1, \mathbf{v}_2)$.*

B.2. **Linearizing the μ-problem** (5.52)**.** Letting $\bar{v} \in \mathcal{C}_T(\mathcal{M})$, we set $\bar{\eta} = e + \int_0^t \bar{v}$ and $\bar{\eta}_\epsilon = e + \int_0^t \bar{v}_\epsilon$ on Γ. We define $\bar{\zeta}_\epsilon$ to be the solution of the following time-dependent elliptic Dirichlet problem:

$$\Delta \bar{\zeta}_\epsilon = \Delta \bar{\eta} \quad \text{in } \Omega, \tag{B.2a}$$

$$\bar{\zeta}_\epsilon = \bar{\eta}_\epsilon \quad \text{on } \Gamma. \tag{B.2b}$$

We define the following ϵ-approximate Lagrangian variables:

$$\bar{A}_\epsilon = [D\bar{\zeta}_\epsilon]^{-1}, \quad \bar{J}_\epsilon = \det D\bar{\zeta}_\epsilon, \quad \bar{a}_\epsilon = \bar{J}_\epsilon \bar{A}_\epsilon, \ [\bar{g}_\epsilon]_{\alpha\beta} = \bar{\zeta}_\epsilon,_\alpha \cdot \bar{\zeta}_\epsilon,_\beta \,, \text{ and } \sqrt{\bar{g}_\epsilon}\bar{n}_\epsilon = [\bar{a}_\epsilon]^T N.$$

We assume that $T > 0$ is given such that independently of the choice of $\bar{v} \in \mathcal{C}_T(\mathcal{M})$, the ϵ-approximate Lagrangian map $\bar{\zeta}_\epsilon$ is injective for all $t \in [0, T]$, and that

$$\frac{1}{2} \le \bar{J}(t) \le \frac{3}{2} \ \text{ and } \ \frac{1}{2} \le \bar{J}_\epsilon(t) \le \frac{3}{2} \ \text{ for all } t \in [0, T] \text{ and } x \in \overline{\Omega}. \tag{B.3}$$

This is possible by the inequality (4.4) as $\bar{v} \in \mathcal{C}_T(\mathcal{M})$ satisfies $\|\bar{v}\|^2_{\mathbf{X}_T} \le \mathcal{M}$.

Definition B.1 (The system of linear heat-equations for v)**.** *For $\bar{v} \in \mathcal{C}_T(\mathcal{M})$, $\kappa > 0$, $\epsilon > 0$ and $\mu > 0$ given, we define v to be the solution of the system of linear equations*

$$v_t - \bar{\varrho}[\bar{A}_\epsilon]^j_r \big([\bar{A}_\epsilon]^k_r v,_k\big),_j = \bar{\mathcal{K}} \qquad \textit{in } \Omega \times (0, T_\kappa(\epsilon\mu)), \tag{B.4a}$$

$$\bar{\varrho} N^j [\bar{A}_\epsilon]^j_r [\bar{A}_\epsilon]^s_r v,_s = \bar{h}^\mu + c^\mu(t) \quad \textit{on } \Gamma \times (0, T_\kappa(\epsilon\mu)), \tag{B.4b}$$

$$v|_{t=0} = u_0 \qquad \textit{on } \Omega. \tag{B.4c}$$

The bounded, nonnegative function $\bar{\varrho}$ is defined as

$$\bar{\varrho} = \kappa \rho_0 \bar{J}_\epsilon. \tag{B.5}$$

The vector field $\bar{\mathcal{K}}$ appearing in the right-hand side of (B.4a) *is given by*

$$\bar{\mathcal{K}} = \bar{\varrho}\, \operatorname{curl}_{\bar{\zeta}_\epsilon} (\operatorname{curl} u_0 + \varepsilon_{\cdot ji} \int_0^t \partial_t [\bar{A}_\epsilon]^s_j \bar{v}^i,_s) + \operatorname{div}_{\bar{\zeta}_\epsilon} \bar{v} [\bar{A}_\epsilon]^k_\cdot \bar{\varrho},_k - 2[\bar{A}_\epsilon]^k_\cdot (\rho_0 \bar{J}^{-1}),_k \,. \tag{B.6}$$

The vector field $\bar{h}^\mu$ appearing in the right-hand side of (B.4b) *is given by*

$$\bar{h}^\mu = \bar{h}_{\text{curl}} + \bar{h}_{\text{div}} + \sum_{l=1}^K \sqrt{\xi_l} \Big(\Lambda_\mu \Big[\bar{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l} \Lambda_\mu \big[(\sqrt{\xi_l} \bar{v},_{\alpha\beta} \cdot \bar{n}_\epsilon) \circ \theta_l \big] \Big] \Big) \circ \theta_l^{-1} \bar{n}_\epsilon + \bar{\varrho} \Big[\bar{b}^\mu_{\text{curl}} + \bar{b}^\mu_{\text{div}} \Big], \tag{B.7}$$

where

$$\bar{\mathfrak{g}}^{\alpha\beta}_{\epsilon,l} = \kappa \frac{\sqrt{\bar{g}_\epsilon}}{\rho_0} \bar{g}^{\alpha\beta}_\epsilon \circ \theta_l, \tag{B.8}$$

and the vectors $\bar{h}_{\text{curl}}$, $\bar{h}_{\text{div}}$, $\bar{b}^\mu_{\text{curl}}$ and $\bar{b}^\mu_{\text{div}}$ are defined as

$$\bar{h}_{\text{curl}} = \bar{\varrho} \sqrt{\bar{g}_\epsilon} (\bar{J}_\epsilon)^{-1} \big(\operatorname{curl} u_0 + \varepsilon_{\cdot ji} \int_0^t \partial_t [\bar{A}_\epsilon]^s_j \bar{v}^i,_s \big) \times \bar{n}_\epsilon, \tag{B.9a}$$

$$\bar{h}_{\text{div}} = \frac{\sqrt{\bar{g}_\epsilon}}{\rho_0} \Big[\rho_0^2 (\bar{J}_\epsilon)^{-2} - \beta_\epsilon(t) + \sigma \bar{g}^{\alpha\beta}_\epsilon \bar{\zeta}_\epsilon,_{\alpha\beta} \cdot \bar{n}_\epsilon \Big] \bar{n}_\epsilon, \tag{B.9b}$$

$$\bar{b}^\mu_{\text{curl}} = \frac{\sqrt{\bar{g}_\epsilon}}{\bar{J}_\epsilon} \Big[\sum_{l=1}^K \bar{g}^{\alpha\beta}_\epsilon \sqrt{\xi_l} \Lambda_\mu [(\sqrt{\xi_l} \bar{v} \cdot \bar{n}_\epsilon),_\beta \circ \theta_l] \circ \theta_l^{-1} + \bar{v}^\gamma \bar{g}^{\gamma\delta}_\epsilon \bar{g}^{\alpha\beta}_\epsilon \bar{\zeta}_\epsilon,_{\delta\beta} \cdot \bar{n}_\epsilon \Big] \bar{\zeta}_\epsilon,_\alpha \,, \tag{B.9c}$$

$$\bar{b}^\mu_{\text{div}} = - \sum_{l=1}^K \sqrt{\xi_l} \Big(\Lambda_\mu \Big[\sqrt{\xi_l} \frac{\sqrt{\bar{g}_\epsilon}}{\bar{J}_\epsilon} \big[\bar{g}^{\alpha\beta}_\epsilon \bar{v}^\alpha,_\beta + \bar{v}^\alpha (\sqrt{\bar{g}_\epsilon} \bar{g}^{\alpha\beta}_\epsilon),_\beta + \bar{v}^3 H(\bar{\zeta}_\epsilon) \big] \circ \theta_l \Big] \Big) \circ \theta_l^{-1} \bar{n}_\epsilon. \tag{B.9d}$$

The vector field $c^\mu(t)$ *appearing in the right-hand side of* (B.4b) *is given by* (5.58)*.*

B.3. **Implementation of the fixed-point scheme to solve the μ-problem** (5.52)**.** We will allow constants to depend on $1/\delta\kappa\epsilon\mu$ in our fixed-point scheme.

Definition B.2 (Notational convention for constants depending on $1/\delta\kappa\epsilon\mu > 0$)**.** *Given* $\bar v \in \mathcal{C}_T(\mathcal{M})$*, we let* $\ddddot{\mathcal{P}}$ *denote a generic polynomial with constant and coefficients depending on* $1/\delta\kappa\epsilon\mu > 0$*. We define the constant* $\ddddot{\mathcal{N}}_0 > 0$ *by*

$$\ddddot{\mathcal{N}}_0 = \ddddot{\mathcal{P}}(\|u_0\|_{100}, \|\rho_0\|_{100}).$$

We let $\ddddot{\mathcal{R}}$ *denote generic lower-order terms satisfying*

$$\int_0^T \ddddot{\mathcal{R}} \le \ddddot{\mathcal{N}}_0 + \delta\|\bar v\|^2_{\mathbf{X}_T} + T\,\ddddot{\mathcal{P}}(\|\bar v\|^2_{\mathbf{X}_T}).$$

Lemma B.2. *Given* $\bar v \in \mathcal{C}_T(\mathcal{M})$*,* $\kappa > 0$*,* $\epsilon > 0$ *and* $\mu > 0$*, there exists a unique* $v \in \mathbf{X}_T$ *solving* (B.4) *and satisfying* (X1)*. Furthermore,*

$$\|v\|^2_{\mathbf{X}_T} \le \int_0^T \ddddot{\mathcal{R}}. \tag{B.10}$$

Proof. Standard parabolic theory provides for the existence and uniqueness of $v \in \mathbf{X}_T$ solving (B.4) and satisfying (X1). For the purpose of establishing the estimate (B.10), it is useful to note the scaling relations

$$\bar{\mathcal{K}} \sim D\bar v + \int_0^t D^2\bar v \quad \text{and} \quad \bar h^\mu \sim \bar v + \int_0^t D\bar v + \Lambda_\mu \bar\partial^2 \bar v + \bar h_\epsilon,$$

where $\bar h_\epsilon \in C^\infty(\Gamma)$ with $|\bar h_\epsilon|_s \le C_\epsilon |\bar v|_s$ thanks to the inequality (5.2). We will establish the inequality (B.10) in the following three steps:

Step 1: Parabolic estimates for v_{tt}. Testing two time-derivatives of the equations (B.4a) against v_{tt} in the $L^2(\Omega)$-inner product and integrating by parts with respect to ∂_j in the integral $-\int_\Omega \partial_t^2\big(\bar\varrho[\bar A_\epsilon]^j_r\big([\bar A_\epsilon]^k_r v,_k\big),_j\big)v_{tt}$ yields

$$\frac{1}{2}\frac{d}{dt}\int_\Omega |v_{tt}|^2 + \int_\Omega \bar\varrho|[\bar A_\epsilon]^k_\cdot v_{tt},_k|^2 = \underbrace{\int_\Omega \bar{\mathcal{K}}_{tt} v_{tt}}_{\ddddot{\mathcal{R}}} + \underbrace{\int_\Gamma \bar h^\mu_{tt} v_{tt}}_{\mathfrak{i}} + \ddddot{\mathcal{R}}. \tag{B.11}$$

We used the boundary condition (B.4b) in writing the boundary integral $\mathfrak{i}$. Thanks to Lemma 2.1, we have that $|v_{tt}|^2_0 = \ddddot{\mathcal{R}}$. Hence, $\mathfrak{i} = \ddddot{\mathcal{R}}$. Taking the time-integral of (B.11) and using that $\bar\varrho \ge \lambda_\epsilon > 0$,

$$\sup_{t\in[0,T]} \|v_{tt}(t)\|^2_0 + \int_0^T \|v_{tt}\|^2_1 \le \int_0^T \ddddot{\mathcal{R}}. \tag{B.12}$$

Step 2: Elliptic estimates for v_t. A time-derivative of the equations (B.4) yields the following linear Neumann-type elliptic problem for v_t:

$$-\bar\varrho[\bar A_\epsilon]^j_r\big([\bar A_\epsilon]^k_r v_t,_k\big),_j = \bar{\mathcal{G}}_1 \text{ in } \Omega, \tag{B.13a}$$

$$\bar\varrho N^j[\bar A_\epsilon]^j_r[\bar A_\epsilon]^s_r v_t,_s = \bar\jmath_1 \text{ on } \Gamma, \tag{B.13b}$$

where the forcing functions $\bar{\mathcal{G}}_1$ and $\bar\jmath_1$ are defined as

$$\begin{aligned}
\bar{\mathcal{G}}_1 &= \bar{\mathcal{K}}_t - v_{tt} + (\bar\varrho[\bar A_\epsilon]^j_r)_t\big([\bar A_\epsilon]^k_r v,_k\big),_j + \bar\varrho[\bar A_\epsilon]^j_r\big(\partial_t[\bar A_\epsilon]^k_r v,_k\big),_j\,,\\
\bar\jmath_1 &= \partial_t\big[\bar h^\mu + c^\mu(t)\big] - \big(\bar\varrho N^j[\bar A_\epsilon]^j_r[\bar A_\epsilon]^s_r\big)_t v,_s\,.
\end{aligned}$$

Since $\bar{\varrho}[\bar{A}_\epsilon]^j_r \partial_j [[\bar{A}_\epsilon]^k_r \partial_k]$ is a uniformly elliptic operator, standard elliptic regularity theory provides that for $s \geq 2$

$$\|v_t\|^2_s \leq C\big(\|v_t\|^2_0 + \|\bar{\mathcal{G}}_1\|^2_{s-2} + |\bar{\jmath}_1|^2_{s-\frac{3}{2}}\big), \tag{B.14}$$

with the constant C depending on the coefficients of the elliptic and Neumann-type operators. The fundamental theorem of calculus provides that

$$\|v_t\|^2_0 \leq \ddddot{\mathcal{N}}_0 + C\,T \sup_{t\in[0,T]} \|v_{tt}(t)\|^2_0 \leq \int_0^T \ddddot{\mathcal{R}}.$$

Thanks to the estimate (B.12), we conclude that

$$\sup_{t\in[0,T]} \|\bar{\mathcal{G}}_1(t)\|^2_0 + \int_0^T \|\bar{\mathcal{G}}_1\|^2_1 \leq \int_0^T \ddddot{\mathcal{R}}.$$

Since $\bar{\jmath}_1$ scales like $Dv + D\bar{v} + \bar{v}_t$, the fundamental theorem of calculus provides that

$$\sup_{t\in[0,T]} |\bar{\jmath}_1(t)|^2_{0.5} + \int_0^T |\bar{\jmath}_1|^2_{1.5} \leq \int_0^T \ddddot{\mathcal{R}} + \sup_{t\in[0,T]} \|\bar{v}_t(t)\|^2_1.$$

Since interpolation and Young's inequality provide that $\|\bar{v}_t\|^2_1 \leq C_\delta \|\bar{v}_t\|^2_0 + \delta \|\bar{v}_t\|^2_2$, we conclude by use of the fundamental theorem of calculus that

$$\sup_{t\in[0,T]} |\bar{\jmath}_1(t)|^2_{0.5} + \int_0^T |\bar{\jmath}_1|^2_{1.5} \leq \int_0^T \ddddot{\mathcal{R}}.$$

It therefore follows from the inequality (B.14) that

$$\sup_{t\in[0,T]} \|v_t(t)\|^2_2 + \int_0^T \|v_t\|^2_3 \leq \int_0^T \ddddot{\mathcal{R}}. \tag{B.15}$$

Step 3: Concluding the proof of Lemma B.2. By repeating Step 2, we infer the following elliptic estimate for v:

$$\sup_{t\in[0,T]} \|v(t)\|^2_4 + \int_0^T \|v\|^2_5 \leq \int_0^T \ddddot{\mathcal{R}}. \tag{B.16}$$

The sum of the inequalities (B.12), (B.15) and (B.16) establishes the inequality (B.10). □

Lemma B.3. *Let v be the solution of* (B.4) *determined by $\bar{v} \in \mathcal{C}_T(\mathcal{M})$. The solutions map*

$$\mathfrak{S} \colon \mathcal{C}_T(\mathcal{M}) \to \mathcal{C}_T(\mathcal{M}) \colon \bar{v} \mapsto v$$

is a well-defined map for some $T = T_\kappa(\epsilon\mu)$ and possesses a unique fixed-point.

Proof. We recall that $v \in \mathbf{X}_T$ is unique and satisfies (X1) by Lemma B.2 . Setting $\mathcal{M} = \ddddot{\mathcal{N}}_0 + 1$, we take δ so that $\delta\|\bar{v}\|^2_{\mathbf{X}_T} < \frac{1}{2}$ and let $T = T_\kappa(\epsilon\mu)$ be sufficiently small so that the inequality given in Lemma B.2 reads

$$\|v\|^2_{\mathbf{X}_T} \leq \mathcal{M}.$$

Hence, for some $T = T_\kappa(\epsilon\mu) > 0$, the solutions map $\mathfrak{S}$ is well-defined.

Letting $\bar{v}_l \in \mathcal{C}_T(\mathcal{M})$ for $l = 1, 2$, we set $\bar{\eta}_l = e + \int_0^t \bar{v}_l$ in Ω and $\bar{\eta}_{\epsilon l} = e + \int_0^t \bar{v}_{\epsilon l}$ on Γ. We define $\bar{\zeta}_{\epsilon l}$ to be the solution of the following time-dependent elliptic Dirichlet problem:

$$\Delta\bar{\zeta}_{\epsilon l} = \Delta\bar{\eta}_l \quad \text{in } \Omega, \tag{B.17a}$$

$$\bar{\zeta}_{\epsilon l} = \bar{\eta}_{\epsilon l} \quad \text{on } \Gamma. \tag{B.17b}$$

We define the following ϵ-approximate Lagrangian variables:

$$\bar{A}_{\epsilon l} = [D\bar{\zeta}_{\epsilon l}]^{-1},\ \bar{J}_l = \det D\bar{\zeta}_{\epsilon l},\ \bar{a}_{\epsilon l} = \bar{J}_{\epsilon l}\bar{A}_{\epsilon l},\ [\bar{g}_{\epsilon l}]_{\alpha\beta} = \bar{\zeta}_{\epsilon l},_\alpha \cdot \bar{\zeta}_{\epsilon l},_\beta\ ,\ \sqrt{\bar{g}_{\epsilon l}}\bar{n}_{\epsilon l} = [\bar{a}_{\epsilon l}]^T N.$$

For $l = 1, 2$, we also take the quantities $\bar{\varrho}_l$, $\bar{\mathcal{K}}_l$, $\bar{h}^\mu_l$ to be respectively formed via the definitions (B.5), (B.6), (B.7) with $\bar{v} = \bar{v}_l$. Letting v_l denote the solution of (B.4) formed using $\bar{v} = \bar{v}_l$, we have that the difference

$$w = v_1 - v_2$$

satisfies

$$w_t - \bar{\varrho}_1[\bar{A}_{\epsilon 1}]^j_r\big([\bar{A}_{\epsilon 1}]^k_r w,_k\big),_j = F \quad \text{in } \Omega \times (0, T_\kappa(\epsilon\mu)], \tag{B.18a}$$

$$\bar{\varrho}_1 N^j [\bar{A}_{\epsilon 1}]^j_r [\bar{A}_{\epsilon 1}]^s_r w,_s = G \quad \text{on } \Gamma \times (0, T_\kappa(\epsilon\mu)], \tag{B.18b}$$

$$w|_{t=0} = 0 \quad \text{on } \Omega, \tag{B.18c}$$

where the vector field F appearing in the right-hand side of (B.18a) is given by

$$F = \bar{\mathcal{K}}_1 - \bar{\mathcal{K}}_2 + \underbrace{\bar{\varrho}_1[\bar{A}_{\epsilon 1}]^j_r\big([\bar{A}_{\epsilon 1}]^k_r v_2,_k\big),_j - \bar{\varrho}_2[\bar{A}_{\epsilon 2}]^j_r\big([\bar{A}_{\epsilon 2}]^k_r v_2,_k\big),_j}_{\mathfrak{I}}, \tag{B.19}$$

and the vector field G appearing in the right-hand side of (B.18b) is given by

$$G = \bar{h}^\mu_1 - \bar{h}^\mu_2 - \underbrace{\big[\bar{\varrho}_1 N^j [\bar{A}_{\epsilon 1}]^j_r [\bar{A}_{\epsilon 1}]^k_r v_2,_k + \bar{\varrho}_2 N^j [\bar{A}_{\epsilon 2}]^j_r [\bar{A}_{\epsilon 2}]^k_r v_2,_k\big]}_{\mathfrak{j}}. \tag{B.20}$$

Testing two time-derivatives of (B.18a) against w_{tt} in the $L^2(\Omega)$-inner product and integrating by parts in the integrals $-\int_\Omega \partial_t^2\big(\bar{\varrho}_1[\bar{A}_{\epsilon 1}]^j_r\big([\bar{A}_{\epsilon 1}]^k_r w,_k\big),_j\big) w_{tt}$ and $\int_\Omega \mathfrak{I}_{tt} w_{tt}$ yields

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\int_\Omega |w_{tt}|^2 + \int_\Omega \partial_t^2\big(\bar{\varrho}_1[\bar{A}_{\epsilon 1}]^j_r[\bar{A}_{\epsilon 1}]^k_r w,_k\big) w_{tt},_j + \int_\Omega \partial_t^2\big((\bar{\varrho}_1[\bar{A}_{\epsilon 1}]^j_r),_j [\bar{A}_{\epsilon 1}]^k_r w,_k\big) w_{tt} \\
&\qquad\qquad - \int_\Gamma (\bar{h}^\mu_1 - \bar{h}^\mu_2)_{tt} w_{tt} \\
&= \int_\Omega \partial_t^2\Big[\bar{\mathcal{K}}_1 - \bar{\mathcal{K}}_2 - \Big((\bar{\varrho}_1[\bar{A}_{\epsilon 1}]^j_r),_j [\bar{A}_{\epsilon 1}]^k_r - (\bar{\varrho}_2[\bar{A}_{\epsilon 2}]^j_r),_j [\bar{A}_{\epsilon 2}]^k_r\Big) v_2,_k\Big] w_{tt} \\
&\qquad\qquad - \int_\Omega \partial_t^2\big[\big(\bar{\varrho}_1[\bar{A}_{\epsilon 1}]^j_r[\bar{A}_{\epsilon 1}]^k_r - \bar{\varrho}_2[\bar{A}_{\epsilon 2}]^j_r[\bar{A}_{\epsilon 2}]^k_r\big) v_2,_k\big] w_{tt},_j\,.
\end{aligned} \tag{B.21}$$

We have used that two time-derivatives of the boundary condition (B.18b) is equivalent to

$$\partial_t^2\big[\bar{\varrho}_1 N^j[\bar{A}_{\epsilon 1}]^j_r[\bar{A}_{\epsilon 1}]^s_r w,_s + \mathfrak{j}\big] = (\bar{h}^\mu_1 - \bar{h}^\mu_2)_{tt}.$$

Since $\partial_t^2(\bar{\varrho}_1 - \bar{\varrho}_2) = \int_0^t \partial_t^3(\bar{\varrho}_1 - \bar{\varrho}_2)$ scales like $\int_0^T D(\bar{v}_1 - \bar{v}_2)_{tt}$, we infer that the time-integral of (B.21) yields

$$\sup_{t\in[0,T]} \|w_{tt}(t)\|_0^2 + \int_0^T \|w_{tt}\|_1^2 \le \delta \int_0^T \|\bar{v}_1 - \bar{v}_2\|^2_{\mathbf{X}_T} + C_{\delta\kappa\mu} T \|\bar{v}_1 - \bar{v}_2\|^2_{\mathbf{X}_T}. \tag{B.22}$$

We use the estimate (B.22) and follow Steps 2 and 3 of the proof of Lemma B.2 to obtain estimates for w_t in $L^2(0, T; H^3(\Omega))$ and w in $L^2(0, T; H^5(\Omega))$. Thus,

$$\|w\|^2_{\mathbf{X}_T} \le \delta \int_0^T \|\bar{v}_1 - \bar{v}_2\|^2_{\mathbf{X}_T} + C_{\delta\kappa\mu} T \|\bar{v}_1 - \bar{v}_2\|^2_{\mathbf{X}_T}. \tag{B.23}$$

We recall that $w = v_1 - v_2$. The inequality (B.23) therefore provides that $\mathfrak{S}$ is a contraction mapping for sufficiently small $\delta > 0$ and $T = T_\kappa(\epsilon\mu) > 0$. Hence, the solutions mapping $\mathfrak{S}$ possess a unique fixed-point $\mathring{v} \in \mathcal{C}_T(\mathcal{M})$. □

B.4. **The proof of Lemma B.1.** The proof of Lemma B.1 is complete since the fixed-point $\mathring{v} \in \mathcal{C}_T(\mathcal{M})$ of Lemma B.3 verifies (X1) and is the unique solution of the μ-problem (5.52).

B.5. **The proof of Proposition B.1.** We infer from the proof of Lemma B.1 that considering two more time-derivatives in the functional framework of our fixed-point scheme establishes Proposition B.1.

Appendix C. Equivalence of the $\kappa\epsilon$-problem (5.6) and the heat-type $\kappa\epsilon$-problem (5.49)

In this appendix, we prove the following

Lemma C.1. *The $\kappa\epsilon$-problem* (5.6) *and the heat-type $\kappa\epsilon$-problem* (5.49) *are equivalent.*

Proof. Sections 5.3.1 and 5.3.2 provide that a solution of the $\kappa\epsilon$-problem (5.6) satisfies the heat-type $\kappa\epsilon$-problem (5.49). We now establish the converse. Using the identity (5.29) stating $-[\check{A}_\epsilon]^j_r[[\check{A}_\epsilon]^k_r\check{v},_k],_j = \operatorname{curl}_{\check{\zeta}_\epsilon}\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v} - [\check{A}_\epsilon]^s_{\cdot}(\operatorname{div}_{\check{\zeta}_\epsilon}\check{v}),_s$, the nonlinear heat-type equations (5.49a) are equivalently written as

$$\check{v}_t + \check{\varrho}\operatorname{curl}_{\check{\zeta}_\epsilon}\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v} - \check{\varrho}[\check{A}_\epsilon]^s_{\cdot}(\operatorname{div}_{\check{\zeta}_\epsilon}\check{v}),_s = \check{\mathcal{K}}.$$

The identity

$$-\check{\varrho}[\check{A}_\epsilon]^s_{\cdot}(\operatorname{div}_{\check{\zeta}_\epsilon}\check{v}),_s = -[\check{A}_\epsilon]^k_{\cdot}(\check{\varrho}\operatorname{div}_{\check{\zeta}_\epsilon}\check{v}),_k + \operatorname{div}_{\check{\zeta}_\epsilon}\check{v}[\check{A}_\epsilon]^k_{\cdot}\check{\varrho},_k\,,$$

and the definition (5.32) of $\check{\mathcal{K}}$ imply that the equations (5.49a) are equivalently written as

$$\check{v}_t - [\check{A}_\epsilon]^k_{\cdot}(\check{\varrho}\operatorname{div}_{\check{\zeta}_\epsilon}\check{v} - 2\rho_0\check{J}_\epsilon^{-1}),_k = -\check{\varrho}\operatorname{curl}_{\check{\zeta}_\epsilon}\big[\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v} - (\operatorname{curl}u_0 + \varepsilon_{\cdot ji}\int_0^t\partial_t[\check{A}_\epsilon]^s_j\check{v}^i,_s)\big]. \tag{C.1}$$

Using the identity $\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v} = \operatorname{curl}u_0 + \int_0^t\partial_t(\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v}) = \operatorname{curl}u_0 + \int_0^t\partial_t(\varepsilon_{\cdot ji}[\check{A}_\epsilon]^s_j\check{v}^i,_s)$,

$$\int_0^t\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v}_t = \operatorname{curl}_{\check{\zeta}_\epsilon}\check{v} - (\operatorname{curl}u_0 + \varepsilon_{\cdot ji}\int_0^t\partial_t[\check{A}_\epsilon]^s_j\check{v}^i,_s). \tag{C.2}$$

The identity (C.2) implies that the equations (C.1) are equivalently written as

$$\check{v}_t - [\check{A}_\epsilon]^k_{\cdot}(\check{\varrho}\operatorname{div}_{\check{\zeta}_\epsilon}\check{v} - 2\rho_0\check{J}_\epsilon^{-1}),_k = -\check{\varrho}\operatorname{curl}_{\check{\zeta}_\epsilon}\big[\int_0^t\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v}_t\big]. \tag{C.3}$$

Applying the ϵ-approximate Lagrangian curl operator $\operatorname{curl}_{\check{\zeta}_\epsilon}$ to (C.3) yields

$$\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v}_t + \operatorname{curl}_{\check{\zeta}_\epsilon}\big(\check{\varrho}\operatorname{curl}_{\check{\zeta}_\epsilon}\int_0^t\operatorname{curl}_{\check{\zeta}_\epsilon}\check{v}_t\big) = 0 \quad \text{in } \Omega. \tag{C.4}$$

We recall $c(t)$ defined in (5.50). Using the definition (5.46) of h, we equivalently have that

$$\begin{aligned} c(t) = \sum_{a=0}^{2}\frac{t^a}{a!}\partial_t^a[\check{\varrho}N^j[\check{A}_\epsilon]^j_r[\check{A}_\epsilon]^s_r\check{v},_s - \check{\varrho}[\check{b}_{\text{curl}} + \check{b}_{\text{div}}]]|_{t=0} \\ - \sum_{a=0}^{2}\frac{t^a}{a!}\partial_t^a[\check{h}_{\text{curl}} + \check{h}_{\text{div}} + \check{\mathfrak{g}}_\epsilon^{\alpha\beta}[\check{v},_{\alpha\beta}\cdot\check{n}_\epsilon]\check{n}_\epsilon]|_{t=0}. \end{aligned}$$

By the identities (5.39) and (5.48), we have that

$$
\begin{aligned}
c(t) &= \sum_{a=0}^{2} \frac{t^a}{a!}\partial_t^a [\underbrace{\check\varrho\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}(\operatorname{curl}_{\check\zeta_\epsilon} \check v)\times \check n_\epsilon + \check\varrho\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}(\operatorname{div}_{\check\zeta_\epsilon}\check v)\check n_\epsilon}_{\check\varrho N^j[\check A_\epsilon]^j_r[\check A_\epsilon]^s_r \check v,_s - \check\varrho\check b}]|_{t=0} \\
&\qquad - \sum_{a=0}^{2} \frac{t^a}{a!}\partial_t^a [\check h_{\rm curl} + \underbrace{\check\varrho \check J_\epsilon^{-1}\sqrt{\check g_\epsilon}(\operatorname{div}_{\check\zeta_\epsilon}\check v)n}_{\check h_{\rm div} + \check{\mathfrak{g}}_\epsilon^{\alpha\beta}[v,_{\alpha\beta}\cdot \check n_\epsilon]\check n_\epsilon}]|_{t=0} \\
&= \sum_{a=0}^{2} \frac{t^a}{a!}\partial_t^a [\check\varrho\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}(\operatorname{curl}_{\check\zeta_\epsilon} \check v)\times \check n_\epsilon - \check h_{\rm curl}]|_{t=0}.
\end{aligned}
$$

Recalling that $\check h_{\rm curl} = \check\varrho\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}\big(\operatorname{curl} u_0 + \varepsilon_{\cdot ji}\int_0^t \partial_t[\check A_\epsilon]^s_j \check v^i,_s\big)\times \check n_\epsilon$, we conclude that

$$
c(t) = \sum_{a=0}^{2} \frac{t^a}{a!}\partial_t^a [\check\varrho\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}(\operatorname{curl}_{\check\zeta_\epsilon}\check v - \operatorname{curl} u_0 - \varepsilon_{\cdot ji}\int_0^t \partial_t[\check A_\epsilon]^s_j\check v^i,_s\,)\times\check n_\epsilon]|_{t=0}.
$$

By the identity (C.2) for $\int_0^t \operatorname{curl}_{\check\zeta_\epsilon}\check v_t$, we have established that

$$
c(t) = \sum_{a=0}^{2} \frac{t^a}{a!}\partial_t^a [\check\varrho\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}(\int_0^t \operatorname{curl}_{\check\zeta_\epsilon}\check v_t)\times\check n_\epsilon]|_{t=0}. \tag{C.5}
$$

We will now verify that $c(t)=0$. With $(\int_0^t \operatorname{curl}_{\check\zeta_\epsilon}\check v_t)|_{t=0}=0$, we equivalently write (C.5) as

$$
c(t) = t\kappa\rho_0\sqrt{\check g_\epsilon}\operatorname{curl}\mathbf{v}_1\times N + \frac{t^2}{2}\kappa\rho_0\sqrt{\check g_\epsilon}[\operatorname{curl}\mathbf{v}_2 + \varepsilon_{\cdot ji}\partial_t[\check A_\epsilon]^s_j(0)\mathbf{v}_1^i,_s\,]\times N.
$$

According to (5.10), we have that $\mathbf{v}_1 = D(\kappa\rho_0\operatorname{div}u_0 - 2\rho_0)$. Hence, $\operatorname{curl}\mathbf{v}_1 = 0$. Using the identity $\partial_t[\check A_\epsilon]^k_i|_{t=0} = -([\check A_\epsilon]^s_i[\check A_\epsilon]^k_r\check v^r,_s)|_{t=0} = -u_0^k,_j$, we conclude that

$$
c(t) = \frac{t^2}{2}\kappa\rho_0\sqrt{\check g_\epsilon}[\operatorname{curl}\mathbf{v}_2 - \varepsilon_{\cdot ji}u_0^s,_j\,\mathbf{v}_1^i,_s\,]\times N.
$$

According to (5.10), $\mathbf{v}_2 = D\partial_t(\kappa\rho_0\check{\mathcal{J}}_t - 2\rho_0\check J_\epsilon^{-1})|_{t=0} + \partial_t[\check A_\epsilon]^k_\cdot|_{t=0}\partial_k(\kappa\rho_0\operatorname{div}u_0 - 2\rho_0)$. Thus, $\operatorname{curl}\mathbf{v}_2 = -\varepsilon_{\cdot ji}\big[u_0^k,_i\,(\kappa\rho_0\operatorname{div}u_0-2\rho_0),_k\big],_j = -\varepsilon_{\cdot ji}u_0^k,_i\,(\kappa\rho_0\operatorname{div}u_0 - 2\rho_0),_{jk} = -\varepsilon_{\cdot ji}u_0^k,_i\,\mathbf{v}_1^j,_k\,$. That is,

$$
\operatorname{curl}\mathbf{v}_2 = \varepsilon_{\cdot ji}u_0^s,_j\,\mathbf{v}_1^i,_s\,,
$$

so that the identity $c(t) = \frac{t^2}{2}\kappa\rho_0\sqrt{\check g_\epsilon}[\operatorname{curl}\mathbf{v}_2 - \varepsilon_{\cdot ji}u_0^s,_j\,\mathrm{v}_1^i,_s\,]\times N$ implies that

$$
c(t) = 0.
$$

The boundary condition (5.49b) is therefore equivalently written as

$$
\begin{aligned}
&\check\varrho N^j[\check A_\epsilon]^j_r[\check A_\epsilon]^s_r\check v,_s = \\
&\qquad \check h_{\rm curl} + \check\varrho\check b + \frac{\sqrt{\check g_\epsilon}}{\rho_0}\Big[\rho_0^2\check J_\epsilon^{-2} - \beta_\epsilon(t) + \sigma\check g_\epsilon^{\alpha\beta}\check\zeta_\epsilon,_{\alpha\beta}\cdot\check n_\epsilon + \check{\mathfrak{g}}_\epsilon^{\alpha\beta}\check v,_{\alpha\beta}\cdot\check n_\epsilon\Big]\check n_\epsilon.
\end{aligned} \tag{C.6}
$$

Taking the scalar product of (C.6) with $\check\varrho^{-1}\bar\partial\check\zeta_\epsilon$ yields

$$
\underbrace{\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}[(\operatorname{curl}_{\check\zeta_\epsilon}\check v)\times\check n_\epsilon]\cdot\bar\partial\check\zeta_\epsilon}_{N^j[\check A_\epsilon]^j_r[\check A_\epsilon]^k_r\check v,_k\cdot\bar\partial\check\zeta_\epsilon} = \underbrace{\sqrt{\check g_\epsilon}\check J_\epsilon^{-1}[\big(\operatorname{curl}u_0 + \varepsilon_{\cdot ji}\int_0^t\partial_t[\check A_\epsilon]^s_j\check v^i,_s\big)\times\check n_\epsilon]\cdot\bar\partial\check\zeta_\epsilon}_{\check\varrho^{-1}\check h_{\rm curl}\cdot\bar\partial\check\zeta_\epsilon}.
$$

Using that $[(\operatorname{curl}_{\check\zeta_\epsilon} \check v) \times \check n_\epsilon] \cdot \check n_\epsilon = 0$ and $\check h_{\text{curl}} \cdot \check n_\epsilon = 0$, we have that

$$(\operatorname{curl}_{\check\zeta_\epsilon} \check v) \times \check n_\epsilon = \big(\operatorname{curl} u_0 + \varepsilon_{\cdot ji} \int_0^t \partial_t [\check A_\epsilon]^s_j \check v^i,_s \big) \times \check n_\epsilon \quad \text{on } \Gamma. \tag{C.7}$$

We note that (C.7) is equivalently stated as $(\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t) \times \check n_\epsilon = 0$. Hence, by the equation (C.4), we record that $\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t$ satisfies

$$\operatorname{curl}_{\check\zeta_\epsilon} \check v_t + \operatorname{curl}_{\check\zeta_\epsilon} (\check\varrho \operatorname{curl}_{\check\zeta_\epsilon} \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t) = 0 \quad \text{in } \Omega \times [0,T], \tag{C.8a}$$

$$(\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t) \times \check n_\epsilon = 0 \quad \text{on } \Gamma \times [0,T], \tag{C.8b}$$

$$(\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t)|_{t=0} = 0 \quad \text{on } \Omega. \tag{C.8c}$$

Testing (C.8a) against $\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t$ in the $L^2(\Omega)$-inner product and integrating by parts with respect to the operator $\operatorname{curl}_{\check\zeta_\epsilon}$ in $\int_\Omega \operatorname{curl}_{\check\zeta_\epsilon} \big(\check\varrho \operatorname{curl}_{\check\zeta_\epsilon} \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t\big) \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t$ yields

$$\begin{aligned} &\frac{1}{2}\frac{d}{dt} \| \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t \|_0^2 + \| \sqrt{\check\varrho} \operatorname{curl}_{\check\zeta_\epsilon} \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t \|_0^2 \\ &\qquad + \int_\Gamma N^s [\check A_\epsilon]^s_j \varepsilon_{ijk} \big(\check\varrho \operatorname{curl}_{\check\zeta_\epsilon} \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t\big)^k \big(\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t\big)^i = 0. \end{aligned} \tag{C.9}$$

With the boundary condition (C.8b), $\int_\Gamma N^s [\check A_\epsilon]^s_j \varepsilon_{ijk} (\check\varrho \operatorname{curl}_{\check\zeta_\epsilon} \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t)^k (\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t)^i = \int_\Gamma \sqrt{\check g_\epsilon} \check J_\epsilon^{-1} [(\int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t) \times n] \cdot (\check\varrho \operatorname{curl}_{\check\zeta_\epsilon} \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t) = 0$. Integrating the identity (C.9) in time from 0 to $t \in (0,T]$ produces $\| \int_0^t \operatorname{curl}_{\check\zeta_\epsilon} \check v_t \|_0^2 = 0$. So according to (C.8a), $\| \operatorname{curl}_{\check\zeta_\epsilon} \check v_t \|_0 = 0$. The regularity of $\check v$ stated in Proposition 5.4 then provides that almost everywhere in Ω,

$$\operatorname{curl}_{\check\zeta_\epsilon} \check v_t = 0.$$

The equations (C.3) are thus written as

$$\check v_t - [\check A_\epsilon]^k_{\cdot} (\kappa \rho_0 \check J_\epsilon \operatorname{div}_{\check\zeta_\epsilon} \check v - 2\rho_0 \check J_\epsilon^{-1}),_k = 0. \tag{C.10a}$$

As the solution $\check v$ of the heat-type $\kappa\epsilon$-problem (5.49) verifies the ϵ-approximate Lagrangian vorticity equation (5.15), we infer that the boundary condition (C.6) of the heat-type problem is equivalently written as

$$\kappa \rho_0^2 \operatorname{div}_{\check\zeta_\epsilon} \check v = \rho_0^2 \check J_\epsilon^{-2} - \beta_\epsilon(t) + \sigma \check g_\epsilon^{\alpha\beta} \check\zeta_\epsilon,_{\alpha\beta} \cdot \check n_\epsilon + \kappa \check g_\epsilon^{\alpha\beta} \check v,_{\alpha\beta} \cdot \check n_\epsilon. \tag{C.10b}$$

The equations (C.10) are equivalent to the momentum equations and boundary condition of the $\kappa\epsilon$-problem (5.6). ☐

LIST OF NOTATION

u	Eulerian velocity	2
p	Eulerian pressure	**??**
ρ	Eulerian density	2
u_0	Initial velocity	**??**
ρ_0	Initial density	**??**
σ	Surface tension	**??**
β	A parameter to the equation of state $p = \alpha\rho^\gamma - \beta$ for $\gamma > 1$	2
$H(\eta)$	Twice the mean curvature of the moving surface $\eta(\Gamma)$	**??**
η	The particle flow map	3
v	Lagrangian velocity	3
f	Lagrangian density	3
A	Inverse of the deformation tensor $D\eta$	3
J	Jacobian determinant of the deformation tensor $D\eta$	3
a	Cofactor matrix of the deformation tensor $D\eta$	3
e	The identity function $e(x) = x$	4
D	The three-dimensional gradient operator	7
$\bar\partial$	The surface gradient operator	7
$\operatorname{div}_\eta, \operatorname{curl}_\eta$	The Lagrangian divergence and curl operators	7
n	The outward-pointing unit normal to the moving surface $\eta(\Gamma)$	9
g	The surface metric induced by the moving surface $\eta(\Gamma)$	9
g_0	The surface metric of the initial surface Γ	9
$\sqrt{g}$	The square root of the determinant of the metric g	9
Δ_g	The Laplace-Beltrami operator	9
$\\|\cdot\\|_s$	The norm of the Hilbert space $H^s(\Omega)$	9
$\vert\cdot\vert_s$	The norm of the Hilbert space $H^s(\Gamma)$	9
N	The outward-pointing unit normal to Γ	10
$\tilde v$	A solution of the κ-problem (3.1)	15
$\tilde\eta$	The Lagrangian map of the solution $\tilde v$ to the κ-problem	15
$E^\kappa(t)$	The higher-order energy function for solutions $\tilde v$ to the κ-problem	15
$\mathcal{P}, \mathcal{N}_0, \mathcal{R}$	The generic system of constants used in the κ-independent estimates	15
Λ_ϵ	The horizontal-convolution operator for $\epsilon > 0$	27
$\check v$	A solution of the $\kappa\epsilon$-problem (5.6)	28
$E^\epsilon(t)$	The higher-order energy function for solutions $\check v$ to the $\kappa\epsilon$-problem	28
$\dot{\mathcal{P}}, \dot{\mathcal{N}}_0, \dot{\mathcal{R}}$	The generic system of constants used in the ϵ-independent estimates	28
$\mathring{v}$	A solution of the μ-problem (5.52)	35
$E^\mu(t)$	The higher-order energy function for solutions $\mathring{v}$ to the μ-problem (5.52)	35
$\ddot{\mathcal{P}}, \ddot{\mathcal{N}}_0, \ddot{\mathcal{R}}$	The generic system of constants used in the μ-independent estimates	35
$\breve v$	A solution of the σ-problem (7.2)	42
$E^\sigma(t)$	The higher-order energy function for solutions $\breve v$ to the surface tension problem	42
$\dddot{\mathcal{P}}, \dddot{\mathcal{N}}_0, \dddot{\mathcal{R}}$	The generic system of constants used in the σ-independent estimates	42
$\mathbf{X}_T$	The Hilbert space used in the fixed-point scheme	63
$\mathcal{C}_T(\mathcal{M})$	A closed, bounded, convex subset of $\mathbf{X}_T$	63
$\bar v$	An arbitrary vector in $\mathcal{C}_T(\mathcal{M})$	64
$\ddddot{\mathcal{P}}, \ddddot{\mathcal{N}}_0, \ddddot{\mathcal{R}}$	The generic system of constants used in the fixed-point scheme	65

References

[1] Adams, R. A. *Sobolev spaces.* Academic Press, 1975.

[2] D. Ambrose and N. Masmoudi. The zero surface tension limit of two-dimensional water waves. *Comm. Pure Appl. Math.* **58**(10):1287-1315, 2005.

[3] D. Ambrose and N. Masmoudi. The zero surface tension limit of three-dimensional water waves. *Indiana Univ. Math. J.*, **58**:479–521, 2009.

[4] G.-Q. Chen and Y.-G. Wang. Existence and stability of compressible current-vortex sheets in three-dimensional magnetohydrodynamics. *Arch. Rational Mech. Anal.* **187**:369–408, 2008.

[5] C.-H. A. Cheng, D. Coutand, and S. Shkoller. Navier-Stokes equations interacting with nonlinear elastic biofluid shell. *SIAM J. Math. Anal.*, **39**(3):742–800 (electronic), 2007.

[6] J.-F. Coulombel and P. Secchi. Nonlinear compressible vortex sheets in two space dimensions. *Ann. Sci. Ecole Norm. Sup.* **41**:85–139, 2008.

[7] J.-F. Coulombel and P. Secchi. Uniqueness of 2-D compressible vortex sheets. *Commun. Pure Appl. Anal.* **8**:1439–1450, 2009.

[8] R. Courant and K. O. Friedrichs. *Supersonic flow and shock waves. Reprinting of the 1948 original. Applied Mathematical Sciences, Vol. 21.* Springer-Verlag, New York-Heidelberg, 1976. xvi+464 pp.

[9] D. Coutand, H. Lindblad and S. Shkoller. A priori estimates for the free-boundary 3-D compressible Euler equations in physical vacuum, *Comm. Math. Phys.*, **296**(2):559–587, 2010.

[10] D. Coutand and S. Shkoller. The interaction between Quasilinear Elastodynamics and the Navier-Stokes equations. *Arch. Rational Mech. Anal.*, **179**:303–352, 2006.

[11] D. Coutand and S. Shkoller. Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J. Amer. Math. Soc.*, **20**(3):829-930 (electronic), 2007.

[12] D. Coutand and S. Shkoller. Well-posedness in smooth function spaces for the moving-boundary 1-D compressible Euler equations in physical vacuum. *Comm. Pure Appl. Math.*, **64**:328–366, 2011.

[13] D. Coutand and S. Shkoller. Well-posedness in smooth function spaces for the moving-boundary 3-D compressible Euler equations in physical vacuum. *Arch. Rational Mech. Anal.*, 2012, DOI: 10.1007/s00205-012-0536-1.

[14] D. Coutand and S. Shkoller. On the finite-time splash and splat singularities for the 3-D free-surface Euler equations. arXiv:1201.4919.

[15] I. Denisova and V. Solonnikov. Classical solvability of the problem of the motion of an isolated mass of compressible fluid. (Russian) *Algebra i Analiz* **14**(1):71–98, 2002.

[16] J. Francheteau and G. Métivier. Existence de chocs faibles pour des systemes quasilinéaires hyperboliques multidimensionnels. *Astérisque* **268**:1–198, 2000.

[17] J. Glimm and A. Majda. Multidimensional hyperbolic problems and computations. *The IMA Volumes in Mathematics and its Applications*, Vol. **29**. Springer, 1991.

[18] O. Gués, G. Métivier, M. Williams and K. Zumbrun. Existence and stability of multidimensional shock fronts in the vanishing viscosity limit. *Arch. Rational Mech. Anal.* **175**:151–244, 2005.

[19] J. Jang and N. Masmoudi. Well-posedness for compressible Euler equations in a physical vacuum singularity. *Comm. Pure Appl. Math.* **62**(10):1327-1385, 2009.

[20] J. Jang and N. Masmoudi. Vacuum in gas and fluid dynamics. *Nonlinear conservation laws and applications*, 315329, IMA Vol. Math. Appl., 153, Springer, 2011.

[21] D. Lannes. Well-posedness of the water-waves equations. *J. Amer. Math. Soc.*, 18:605–654, 2005.

[22] H. Lindblad. Well-posedness for the motion of an incompressible liquid with free surface boundary. *Annals of Math.*, **162**:109–194, 2005.

[23] H. Lindblad. Well posedness for the motion of a compressible liquid with free surface boundary. *Commun. Math. Phys.* **260**:319–392, 2005.

[24] A. Majda. *Compressible Fluid Flow and Systems of Conservation Laws in Several Space Variables.* Springer, 1984.

[25] G. Métivier. Stability of multidimensional shocks. Advances in the Theory of Shock Waves. *Progress in Nonlinear Differential Equations and Their Applications*, Vol. 47. Birkhäuser, 25103, 2001.

[26] L. Rayleigh, *On the instability of jets*, Proceedings of the London Mathematical Society, **s1-10**(1):4–13, 1878.

[27] J. Shatah and C. Zeng. Geometry and a priori estimates for free boundary problems of the Euler equation. *Comm. Pure Appl. Math.*, **61**:698–744, 2008.

[28] V. Solonnikov and A. Tani. *Free boundary problem for a viscous compressible flow with a surface tension. Constantin Carathodory: an international tribute, Vol. I, II*, 1270–1303, World Sci. Publ., Teaneck, NJ, 1991.

[29] N. Tanaka and A. Tani. Surface waves for a compressible viscous fluid. (English summary) *J. Math. Fluid Mech.*, **5**(4):303–363, 2003.

[30] G. Taylor, *The instability of liquid surfaces when accelerated in a direction perpendicular to their planes I.*, Proc. Roy. Soc. London. Ser. A., **201**:192–196, 1950.

[31] R. Temam. *Navier-Stokes equations. Theory and numerical analysis.* North-Holland Publishing Co., Amsterdam, 1977. Studies in Mathematics and its Applications, Vol. 2.

[32] Y. Trakhinin. Existence of compressible current-vortex sheets: variable coefficients linear analysis. *Arch. Rational Mech. Anal.* **177** 331–366, 2005.

[33] Y. Trakhinin. Local existence for the free boundary problem for the non-relativistic and relativistic compressible Euler equations with a vacuum boundary condition. *Comm. Pure Appl. Math*, **62**:1551–1594, 2009.

[34] Y. Trakhinin. The existence of current-vortex sheets in ideal compressible magnetohydrodynamics. *Arch. Rational Mech. Anal.* **191**:245–310, 2009.

[35] S. Wu. Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.*, **130**:39–72, 1997.

[36] S. Wu. Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Amer. Math. Soc.*, **12**:445–495, 1999.

[37] C.-J. Xu and T. Yang. Local existence with physical vacuum boundary condition to Euler equations with damping. *J. Differential Equations*, **210**:217–231, 2005.

[38] W. M. Zajaczkowski. On nonstationary motion of a compressible barotropic viscous capillary fluid bounded by a free surface. *SIAM J. Math. Anal.*, **25**(1):1-84, 1994.

[39] P. Zhang and Z. Zhang. On the free boundary problem of three-dimensional incompressible Euler equations. *Comm. Pure Appl. Math.*, **61**:877–940, 2008.

CANPDE, Maxwell Institute for Mathematical Sciences and department of Mathematics, Heriot-Watt University, Edinburgh, EH14 4AS, UK
E-mail address: `D.Coutand@ma.hw.ac.uk`

Department of Mathematics, University of California, Davis, CA 95616, USA
E-mail address: `jhole@math.ucdavis.edu`
E-mail address: `shkoller@math.ucdavis.edu`